

04-03-00

03/31/00

00544795-033100

Please type a plus sign (+) inside this box → ☐

PTO/SB/05 (4/98)

Approved for use through 09/30/2000. OMB 0651-0032
Patent and Trademark Office, U.S. DEPARTMENT OF COMMERCE
Under the Paperwork Reduction Act of 1995, no persons are required to respond to a collection of information unless it displays a valid OMB control number.UTILITY
PATENT APPLICATION
TRANSMITTAL

(Only for new nonprovisional applications under 37 C.F.R. § 1.53(b))

Attorney Docket No. 6446.US.P2

First Inventor or Application Identifier James Link

Title See 1 in Addendum

Express Mail Label No. EL384169956US

Assistant Commissioner for Patents
Box Patent Application
Washington, DC 20231

APPLICATION ELEMENTS

See MPEP chapter 600 concerning utility patent application contents.

ADDRESS TO:

1. ☒ * Fee Transmittal Form (e.g., PTO/SB/17)
(Submit an original and a duplicate for fee processing)
2. ☒ Specification [Total Pages 472]
(preferred arrangement set forth below)
- Descriptive title of the Invention
 - Cross References to Related Applications
 - Statement Regarding Fed sponsored R & D
 - Reference to Microfiche Appendix
 - Background of the Invention
 - Brief Summary of the Invention
 - Brief Description of the Drawings (if filed)
 - Detailed Description
 - Claim(s)
 - Abstract of the Disclosure
3. ☐ Drawing(s) (35 U.S.C. 113) [Total Sheets]
4. ☐ Oath or Declaration [Total Pages 5]
- a. ☐ Newly executed (original or copy)
 - b. ☐ Copy from a prior application (37 C.F.R. § 1.63(d))
(for continuation/divisional with Box 16 completed)
 - i. ☐ DELETION OF INVENTOR(S)
Signed statement attached deleting inventor(s) named in the prior application, see 37 C.F.R. §§ 1.63(d)(2) and 1.33(b).

* NOTE FOR ITEMS 1 & 13 IN ORDER TO BE ENTITLED TO PAY SMALL ENTITY FEES, A SMALL ENTITY STATEMENT IS REQUIRED (37 C.F.R. § 1.27), EXCEPT IF ONE FILED IN A PRIOR APPLICATION IS RELIED UPON (37 C.F.R. § 1.28).

5. ☐ Microfiche Computer Program (Appendix)
6. Nucleotide and/or Amino Acid Sequence Submission (if applicable, all necessary)
- a. ☐ Computer Readable Copy
 - b. ☐ Paper Copy (identical to computer copy)
 - c. ☐ Statement verifying identity of above copies

ACCOMPANYING APPLICATION PARTS

7. ☐ Assignment Papers (cover sheet & document(s))
8. ☐ 37 C.F.R. § 3.73(b) Statement of Power of Attorney (when there is an assignee)
9. ☐ English Translation Document (if applicable)
10. ☐ Information Disclosure Statement (IDS)/PTO-1449 ☐ Copies of IDS Citations
11. ☐ Preliminary Amendment
12. ☒ Return Receipt Postcard (MPEP 503)
(Should be specifically itemized)
- * Small Entity Statement filed in prior application (PTO/SB/09-12)
 - 13. ☐ Statement(s) Status still proper and desired
 - 14. ☐ Certified Copy of Priority Document(s) (if foreign priority is claimed)
 - 15. ☒ Other: Certificate of Mailing

16. If a CONTINUING APPLICATION, check appropriate box, and supply the requisite information below and in a preliminary amendment:
- ☐ Continuation ☐ Divisional ☒ Continuation-in-part (CIP) of prior application No. 09/474,517
- Prior application information: Examiner not yet assigned Group / Art Unit: 1614

For CONTINUATION or DIVISIONAL APPS only: The entire disclosure of the prior application, from which an oath or declaration is supplied under Box 4b, is considered a part of the disclosure of the accompanying continuation or divisional application and is hereby incorporated by reference. The incorporation can only be relied upon when a portion has been inadvertently omitted from the submitted application parts.

17. CORRESPONDENCE ADDRESS

☐ Customer Number or Bar Code Label

(Insert Customer No. or Attach bar code label here)

or ☒ Correspondence address below

Name	Steven F. Weinstock				
	Abbott Laboratories				
Address	D377/AP6-D				
	100 Abbott Park Road				
City	Abbott Park		State	IL	Zip Code
Country		Telephone	(847) 937-4395	Fax	(847) 938-2623

Name (Print/Type)	Janelle D. Strode	Registration No. (Attorney/Agent)	34,738
Signature	<i>Janelle D. Strode</i>	Date	March 31, 2000

Burden Hour Statement: This form is estimated to take 0.2 hours to complete. Time will vary depending upon the needs of the individual case. Any comments on the amount of time you are required to complete this form should be sent to the Chief Information Officer, Patent and Trademark Office, Washington, DC 20231. DO NOT SEND FEES OR COMPLETED FORMS TO THIS ADDRESS. SEND TO: Assistant Commissioner for Patents, Box Patent Application, Washington, DC 20231.

IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

In re application of: J. Link, *et al.*

Application No.: (not yet assigned)

Group No.: (not yet assigned)

Filed:

Examiner: (not yet assigned)

For: CELL ADHESION-INHIBITING ANTIINFLAMMATORY AND IMMUNE-SUPPRESSIVE COMPOUNDS

Assistant Commissioner for Patents

Washington, D.C. 20231

EXPRESS MAIL CERTIFICATE

"Express Mail" label number EL384169845US


Date of Deposit: March 31, 2000

I hereby state that the following *attached* papers

1. Utility Patent Application Transmittal (in duplicate)
2. Fee Transmittal Form PTO SB/17 (in duplicate)
3. Invention specification including:
 - Description of invention (416 pages)
 - 26 Claims (55 pages)
 - One (1) page abstract
4. Unexecuted Declaration and Power of Attorney (in duplicate)
5. Return Receipt Postcard

are being deposited with the United States Postal Service "Express Mail Post Office to Addressee" service under 37 CFR 1.10, on the date indicated above and is addressed to the Assistant Commissioner for Patents, Washington, D.C. 20231.

Katrina Williams



Signature of person mailing paper or fee

09541795-033100

CELL ADHESION-INHIBITING ANTIINFLAMMATORY
AND IMMUNE-SUPPRESSIVE COMPOUNDS

This application is a continuation-in-part of Application Serial Number 09/474,517, filed December 29, 1999, which is a continuation-in-part of provisional Application Serial Number 60/114,097, filed December 29, 1998.

5

Technical Field

The present invention relates to compounds that are useful for treating
10 inflammatory and immune diseases, to pharmaceutical compositions comprising these compounds, and to methods of inhibiting inflammation or suppressing immune response in a mammal.

Background

15 Inflammation results from a cascade of events that includes vasodilation accompanied by increased vascular permeability and exudation of fluid and plasma proteins. This disruption of vascular integrity precedes or coincides with an infiltration of inflammatory cells. Inflammatory mediators generated at the site of the initial lesion serve to recruit inflammatory cells to the site of injury. These mediators

001220.56214560

(chemokines such as IL-8, MCP-1, MIP-1, and RANTES, complement fragments and lipid mediators) have chemotactic activity for leukocytes and attract the inflammatory cells to the inflamed lesion. These chemotactic mediators which cause circulating leukocytes to localize at the site of inflammation require the cells to cross the vascular endothelium at a precise location. This leukocyte recruitment is accomplished by a process called cell adhesion.

- Cell adhesion occurs through a coordinately regulated series of steps that allow the leukocytes to first adhere to a specific region of the vascular endothelium and then cross the endothelial barrier to migrate to the inflamed tissue (Springer, T.A., 1994, Traffic Signals for Lymphocyte Recirculation and Leukocyte Emigration: The Multistep Paradigm, Cell 76: 301-314; Lawrence, M. B., and Springer, T. A., 1991, Leukocytes' Roll on a Selectin at Physiologic Flow Rates: Distinction from and Prerequisite for Adhesion Through Integrins, Cell.65: 859-873; von Adrian, U., Chambers, J. D., McEnvoy, L.M., Bargatze, R.F., Arfos, K.E, and Butcher, E.C., 1991, Two-Step Model of Leukocyte-Endothelial Cell Interactions in Inflammation, Proc. Natl. Acad. Sci. USA 88: 7538-7542; and Ley, K., Gaehtgens, P., Fennie, C., Singer, M.S., Lasky, L.H. and Rosen, S.D.,1991, Lectin-Like Cell Adhesion Molecule 1 Mediates Rolling in Mesenteric Venules *in vivo*, Blood 77: 2553-2555). These steps are mediated by families of adhesion molecules such as integrins, Ig supergene family members, and selectins which are expressed on the surface of the circulating leukocytes and on the vascular endothelial cells. The first step consists of leukocytes rolling along the vascular endothelial cell lining in the region of inflammation. The

rolling step is mediated by an interaction between a leukocyte surface oligosaccharide, such as Sialylated Lewis-X antigen (SLe^x), and a selectin molecule expressed on the surface of the endothelial cell in the region of inflammation. The selectin molecule is not normally expressed on the surface of endothelial cells but rather is induced by the

5 action of inflammatory mediators such as TNF- α and interleukin-1. Rolling decreases the velocity of the circulating leukocyte in the region of inflammation and allows the cells to more firmly adhere to the endothelial cell. The firm adhesion is accomplished by the interaction of integrin molecules that are present on the surface of the rolling leukocytes and their counter-receptors (the Ig superfamily molecules) on the surface

10 of the endothelial cell. The Ig superfamily molecules or CAMs (Cell Adhesion Molecules) are either not expressed or are expressed at low levels on normal vascular endothelial cells. The CAM's, like the selectins, are induced by the action of inflammatory mediators like TNF-alpha and IL-1. The final event in the adhesion process is the extravasation of leukocytes through the endothelial cell barrier and their

15 migration along a chemotactic gradient to the site of inflammation. This transmigration is mediated by the conversion of the leukocyte integrin from a low avidity state to a high avidity state. The adhesion process relies on the induced expression of selectins and CAM's on the surface of vascular endothelial cells to mediate the rolling and firm adhesion of leukocytes to the vascular endothelium.

20 The interaction of the intercellular adhesion molecule ICAM-1 (cd54) on endothelial cells with the integrin LFA-1 on leukocytes plays an important role in endothelial-leukocyte contact. Leukocytes bearing high-affinity LFA-1 adhere to

endothelial cells through interaction with ICAM-1, initiating the process of extravasation from the vasculature into the surrounding tissues. Thus, an agent which blocks the ICAM-1/LFA-1 interaction suppresses these early steps in the inflammatory response. Consistent with this background, ICAM-1 knockout mice

5 have numerous abnormalities in their inflammatory responses.

The present invention discloses compounds which bind to the interaction-domain (I-domain) of LFA-1, thus interrupting endothelial cell-leukocyte adhesion by blocking the interaction of LFA-1 with ICAM-1, ICAM-3, and other adhesion molecules. These compounds are useful for the treatment or prophylaxis of diseases

10 in which leukocyte trafficking plays a role, notably acute and chronic inflammatory diseases, autoimmune diseases, tumor metastasis, allograft rejection, and reperfusion injury. The compounds of this invention are diaryl sulfides, which are substituted with a cinnamide moiety. The cinnamide functionality may be placed either ortho- or para- to the linking sulfur atom, although para-substitution is preferable. Appropriate


15 substitution of both aromatic rings is tolerated, and can be used to modulate a variety of biochemical, physicochemical and pharmacokinetic properties. In particular the amide moiety is readily modified; a variety of secondary and tertiary amides are active, and alternatively a heterocyclic ring may be attached at this position. Modifications of this amide functionality are particularly useful in modulating

20 physicochemical and pharmacokinetic properties.

00544795.033100
DATE: 05/27/2000

Summary of The Invention

The present invention provides compounds of formula **I**, below,


**I**

or a pharmaceutically-acceptable salt or prodrug thereof,


wherein R₁, R₂, R₃, R₄, and R₅ are independently selected from

- a. hydrogen,
- b. halogen,
- c. alkyl,
- d. haloalkyl,
- e. alkoxy,
- f. cyano,
- g. nitro,
- h. carboxaldehyde, and

with the proviso that at least one of R_1 or R_3 is a “*cis*-cinnamide” or a “*trans*-cinnamide”, defined as



“*cis*-cinnamide”



“*trans*-cinnamide”,

wherein R_8 and R_9 are independently selected from

- a. hydrogen, and
- b. alkyl,
- c. carboxy alkyl,
- d. alkylaminocarbonyl alkyl, and
- e. dialkylaminocarbonyl alkyl,

and R_{10} and R_{11} are independently selected from

- a. hydrogen,
- b. alkyl,
- c. cycloalkyl,
- d. alkoxy carbonyl alkyl,
- e. hydroxyalkyl,

- f. substituted aryl,
g. heterocyclyl,
h. heterocyclylalkyl,
i. heterocyclylamino,
5 j. substituted heterocyclyl, and
k. substituted heterocyclylalkyl,

or where $\text{NR}_{10}\text{R}_{11}$ is heterocyclyl or substituted heterocyclyl, where
substituents are independently selected from

10

- 1) alkyl,
2) alkoxy,
3) alkoxyalkyl,
4) cycloalkyl,
15 5) aryl,
6) heterocyclyl,
7) heterocyclylcarbonyl,
8) heterocyclylalkylaminocarbonyl,
9) hydroxy,
20 10) hydroxyalkyl,
11) hydroxyalkoxyalkyl,
12) carboxy,

- 13) carboxyalkyl,
14) carboxycarbonyl,
15) carboxaldehyde,
16) alkoxycarbonyl,
5 17) arylalkoxycarbonyl,
18) aminoalkyl,
19) aminoalkanoyl,
20) carboxamido,
21) alkoxycarbonylalkyl,
10 22) carboxamidoalkyl,
23) cyano,
24) tetrazolyl,
25) substituted tetrazolyl,
26) alkanoyl,
15 27) hydroxyalkanoyl,
28) alkanoyloxy,
29) alkanoylamino,
30) alkanoyloxyalkyl,
31) alkanoylaminoalkyl,
20 32) sulfonate,
33) alkylsulfonyl,
34) alkylsulfonylaminocarbonyl,

03544795-03310
01110-5621450

35) arylsulfonylaminocarbonyl, and

36) heterocyclylsulfonylaminocarbonyl,

and wherein **Ar** is a substituted aryl or substituted heteroaryl group, where

5 substitutions are independently selected from

a. hydrogen,

b. halogen,

c. alkyl,

10 d. aryl,

e. haloalkyl,

f. hydroxy,

g. alkoxy,

h. alkoxyalkyl,

15 i. alkoxycarbonyl,

j. alkoxyalkoxy,

k. hydroxyalkyl,

l. aminoalkyl,

m. aminocarbonyl,

20 n. alkyl(alkoxycarbonylalkyl)aminoalkyl,

o. heterocyclyl,

p. substituted heterocyclyl,

00120-5621-560

- q. heterocyclalkyl,
- r. substituted heterocyclalkyl,
- s. carboxaldehyde,
- t. carboxaldehyde hydrazone,
- 5 u. carboxamide,
- v. alkoxycarbonylalkyl,
- w. carboxy,
- x. carboxyalkyl,
- y. carboxy alkoxy,
- 10 z. carboxythioalkoxy,
- aa. carboxycycloalkoxy,
- bb. thioalkyl,
- cc. hydroxycarbonylalkyl (carboxyalkyl),
- dd. hydroxyalkylaminocarbonyl,
- 15 ee. cyano,
- ff. amino,
- gg. heterocyclalkylamino,
- hh. carboxyalkylamino,
- ii. heterocyclalkylaminocarbonyl, and
- 20 jj. "*trans*-cinnamide".

00000-5674560

Additionally provided are methods of treatment or prophylaxis in which the inhibition of inflammation or suppression of immune response is desired, comprising administering an effective amount of a compound of formula I.

Still further provided are pharmaceutical compositions containing compounds
5 of formula I.

00130-562450

Detailed Description

The term "alkanoyl" as used herein refers to an alkyl group attached to the parent molecular group through a carbonyl group.

The term "alkanoylamino" as used herein refers to an alkanoyl group attached
5 to the parent molecular group through an amino group.

The term "alkanoylaminoalkyl" as used herein refers to an alkanoylamino group attached to the parent molecular group through an alkyl group.

The term "alkanoyloxy" as used herein refers to an alkanoyl group attached to the parent molecular group through an oxygen radical.

10 The term "alkanoyloxyalkyl" as used herein refers to an alkanoyloxy group attached to the parent molecular group through an alkyl group.

The term "alkoxy" as used herein refers to an alkyl group attached to the parent molecular group through an oxygen atom.

15 The term "alkoxyalkoxy" as used herein refers to an alkoxy group attached to the parent molecular group through an alkoxy group.

The term "alkoxyalkyl" as used herein refers to an alkoxy group attached to the parent molecular group through an alkyl group.

The term "alkoxycarbonyl" as used herein refers to an alkoxy group attached to the parent molecular group through a carbonyl group.

20 The term "alkoxycarbonylalkyl" as used herein refers to an alkoxycarbonyl group attached to the parent molecular group through an alkyl group.

09541795-023100

The term "alkyl" as used herein refers to a saturated straight or branched chain group of 1-10 carbon atoms derived from an alkane by the removal of one hydrogen atom.

The term "alkyl(alkoxycarbonylalkyl)amino" as used herein refers to an amino group substituted with one alkyl group and one alkoxycarbonylalkyl group.

The term "alkyl(alkoxycarbonylalkyl)aminoalkyl" as used herein refers to an alkyl(alkoxycarbonylalkyl)amino group attached to the parent molecular group through an alkyl group.

The term "alkylene" as used herein refers to a divalent group of 1-10 carbon atoms derived from a straight or branched chain alkane by the removal of two hydrogen atoms.

The term "alkylsulfonyl" as used herein refers to an alkyl radical attached to the parent molecular group through an $-SO_2-$ group.

The term "alkylsulfonylaminocarbonyl" as used herein refers to an alkylsulfonyl group attached to the parent molecular group through an aminocarbonyl group.

The term "amino" as used herein refers to a radical of the form $-NR_{18}R_{19}$, or to a radical of the form $-NR_{18}-$, where R_{18} and R_{19} are independently selected from hydrogen, alkyl or cycloalkyl.

The term "aminoalkanoyl" as used herein refers to an amino group attached to the parent molecular group through an alkanoyl group.

00544795.033140

The term "aminoalkyl" as used herein refers to an amino group attached to the parent molecular group through an alkyl group.

The term "aminocarbonyl" as used herein refers to an amino group attached to the parent molecular group through a carbonyl group.

5 The term "aryl" as used herein refers to a mono- or bicyclic carbocyclic ring system having one or two aromatic rings. The aryl group can also be fused to a cyclohexane, cyclohexene, cyclopentane or cyclopentene ring. The aryl groups of this invention can be optionally substituted with alkyl, halogen, hydroxy, carboxy, or alkoxy substituents.

10 The term "aryloalkoxy" as used herein refers to an aryl group attached to the parent molecular group through an alkoxy group.

The term "aryloalkoxycarbonyl" as used herein refers to an aryloalkoxy group attached to the parent molecular group through a carbonyl group.

15 The term "arylsulfonyl" as used herein refers to an aryl radical attached to the parent molecular group through an $-SO_2-$ group.

The term "arylsulfonylaminoalkyl" as used herein refers to an arylsulfonyl group attached to the parent molecular group through an aminocarbonyl group.

The term "carboxaldehyde" as used herein refers to the radical $-CHO$.

20 The term "carboxaldehyde hydrazone" as used herein refers to the radical $-CH=N-NR_{20}R_{21}$, where R_{20} and R_{21} are independently selected from hydrogen, alkyl or cycloalkyl.

09:44:35 - 03:10:00

The terms "carboxamide" or "carboxamido" as used herein refer to an amino group attached to the parent molecular group through a carbonyl group.

The term "carboxamidoalkyl" as used herein refers to a carboxamido group attached to the parent molecular group through an alkyl group.

5 The term "carboxy" as used herein refers to the radical -COOH.

The term "carboxyalkyl" as used herein refers to a carboxy group attached to the parent molecular group through an alkyl group.

The term "carboxycarbonyl" as used herein refers to a carboxy group attached to the parent molecular group through a carbonyl group.

10 The term "cyano" as used herein refers to the radical -CN.

The term "cycloalkyl" as used herein refers to a monovalent saturated cyclic or bicyclic hydrocarbon group of 3-12 carbons derived from a cycloalkane by the removal of a single hydrogen atom. Cycloalkyl groups may be optionally substituted with alkyl, alkoxy, halo, or hydroxy substituents.

15 The terms "halo" or "halogen" as used herein refers to F, Cl, Br, or I.

The term "haloalkyl" as used herein refers to an alkyl group substituted with one or more halogen atoms.


The terms "heterocycle" or "heterocyclyl" represent a 4-, 5-, 6- or 7-membered ring containing one, two or three heteroatoms independently selected from the group consisting of nitrogen, oxygen and sulfur. The 4- and 5-membered rings have zero to two double bonds and the 6- and 7-membered rings have zero to three double bonds. The term "heterocycle" or "heterocyclic" as used herein additionally refers to bicyclic,

20

- tricyclic and tetracyclic groups in which any of the above heterocyclic rings is fused to one or two rings independently selected from an aryl ring, a cyclohexane ring, a cyclohexene ring, a cyclopentane ring, a cyclopentene ring or another monocyclic heterocyclic ring. Heterocycles include acridinyl, benzimidazolyl, benzofuryl,
- 5 benzothiazolyl, benzothienyl, benzoxazolyl, biotinyl, cinnoliny, dihydrofuryl, dihydroindolyl, dihydropyranyl, dihydrothienyl, dithiazolyl, furyl, homopiperidinyl, imidazolidinyl, imidazoliny, imidazolyl, indolyl, isoquinolyl, isothiazolidinyl, isothiazolyl, isoxazolidinyl, isoxazolyl, morpholiny, oxadiazolyl, oxazolidinyl, oxazolyl, piperazinyl, piperidinyl, pyranyl, pyrazolidinyl, pyrazinyl, pyrazolyl,
- 10 pyrazoliny, pyridazinyl, pyridyl, pyrimidinyl, pyrimidyl, pyrrolidinyl, pyrrolidin-2-onyl, pyrroliny, pyrrolyl, quinoliny, quinoxaloyl, tetrahydrofuryl, tetrahydroisoquinolyl, tetrahydroquinolyl, tetrazolyl, thiadiazolyl, thiazolidinyl, thiazolyl, thienyl, thiomorpholiny, triazolyl, and the like.

Heterocyclics also include bridged bicyclic groups where a monocyclic

- 15 heterocyclic group is bridged by an alkylene group such as



Heterocyclics also include compounds of the formula



where X*

and Z* are independently selected from -CH₂-, -CH₂NH-, -CH₂O-, -NH- and -O-,

- 20 with the proviso that at least one of X* and Z* is not -CH₂-, and Y* is selected from

-C(O)- and -(C(R'')₂)_v -, where R'' is hydrogen or alkyl of one to four carbons, and v is 1-3. These heterocycles include 1,3-benzodioxolyl, 1,4-benzodioxanyl, 1,3-benzimidazol-2-one and the like. The heterocycle groups of this invention can be optionally substituted with alkoxy, alkyl, halogen, hydroxy, carboxy, carboxyalkyl, or alkoxy carbonyl substituents.

The term "heterocyclalalkyl" as used herein refers to an heterocyclic group attached to the parent molecular group through an alkyl group.

The term "heterocyclalalkylamino" as used herein refers to an heterocyclalalkyl group attached to the parent molecular group through an amino group.

The term "heterocyclalalkylaminocarbonyl" as used herein refers to a heterocyclalalkylamino group attached to the parent molecular group through a carbonyl group.

The term "heterocyclalamino" as used herein refers to a heterocyclal group attached to the parent molecular group through a amino group.

The term "heterocyclalcarbonyl" as used herein refers to a heterocyclal group attached to the parent molecular group through a carbonyl group.

The term "heterocyclalsulfonyl" as used herein refers to a heterocyclal radical attached to the parent molecular group through an -SO₂- group.

The term "heterocyclalsulfonylaminocarbonyl" as used herein refers to a heterocyclalsulfonyl group attached to the parent molecular group through an aminocarbonyl group.

The term "hydroxyalkanoyl" as used herein refers to an hydroxy radical attached to the parent molecular group through an alkanoyl group.

The term "hydroxyalkoxy" as used herein refers to an hydroxy radical attached to the parent molecular group through an alkoxy group.

- 5 The term "hydroxyalkoxyalkyl" as used herein refers to an hydroxyalkoxy group attached to the parent molecular group through an alkyl group.

The term "hydroxyalkyl" as used herein refers to an hydroxy radical attached to the parent molecular group through an alkyl group.

- 10 The term "hydroxyalkylaminocarbonyl" as used herein refers to an hydroxyalkyl group attached to the parent molecular group through an aminocarbonyl group.

The term "perfluoroalkyl" as used herein refers to an alkyl group in which all of the hydrogen atoms have been replaced by fluoride atoms.

- 15 The term "phenyl" as used herein refers to a monocyclic carbocyclic ring system having one aromatic ring. The phenyl group can also be fused to a cyclohexane or cyclopentane ring. The phenyl groups of this invention can be optionally substituted with alkyl, halogen, hydroxy or alkoxy substituents.

- 20 The term "pharmaceutically-acceptable prodrugs" as used herein represents those prodrugs of the compounds of the present invention which are, within the scope of sound medical judgment, suitable for use in contact with the tissues of humans and lower animals with undue toxicity, irritation, allergic response, and the like,

commensurate with a reasonable benefit/risk ratio, and effective for their intended use, as well as the zwitterionic forms, where possible, of the compounds of the invention.

The term "prodrug," as used herein, represents compounds which are rapidly transformed *in vivo* to the parent compound of the above formula, for example, by hydrolysis in blood. A thorough discussion is provided in T. Higuchi and V. Stella, Pro-drugs as Novel Delivery Systems, Vol. 14 of the A.C.S. Symposium Series, and in Edward B. Roche, ed., Bioreversible Carriers in Drug Design, American Pharmaceutical Association and Pergamon Press, 1987, both of which are incorporated herein by reference.

The term "sulfonate" as used herein refers to the radical $-\text{SO}_3\text{H}$

The term "tetrazole" or "tetrazolyl" as used herein refers to the heterocyclic radical $-\text{CN}_4\text{H}$.

The term "thioalkoxy" as used herein refers to an alkyl group attached to the parent molecular group through a sulfur atom.

Compounds of the present invention can exist as stereoisomers wherein asymmetric or chiral centers are present. These compounds are designated by the symbols "R" or "S," depending on the configuration of substituents around the chiral carbon atom. The present invention contemplates various stereoisomers and mixtures thereof. Stereoisomers include enantiomers and diastereomers, and mixtures of enantiomers or diastereomers are designated (\pm). Individual stereoisomers of compounds of the present invention can be prepared synthetically from commercially available starting materials which contain asymmetric or chiral centers or by

preparation of racemic mixtures followed by resolution well-known to those of ordinary skill in the art. These methods of resolution are exemplified by (1) attachment of a mixture of enantiomers to a chiral auxiliary, separation of the resulting mixture of diastereomers by recrystallization or chromatography and

5 liberation of the optically pure product from the auxiliary, (2) salt formation employing an optically active resolving agent, or (3) direct separation of the mixture of optical enantiomers on chiral chromatographic columns.

Geometric isomers can also exist in the compounds of the present invention. The present invention contemplates the various geometric isomers and mixtures

10 thereof resulting from the arrangement of substituents around a carbon-carbon double bond or arrangement of substituents around a carbocyclic ring. Substituents around a carbon-carbon double bond are designated as being in the Z or E configuration wherein the term "Z" represents substituents on the same side of the carbon-carbon double bond and the term "E" represents substituents on opposite sides of the carbon-

15 carbon double bond. The arrangement of substituents around a carbocyclic ring are designated as cis or trans wherein the term "cis" represents substituents on the same side of the plane of the ring and the term "trans" represents substituents on opposite sides of the plane of the ring. Mixtures of compounds wherein the substituents are disposed on both the same and opposite sides of plane of the ring are designated

20 cis/trans.

As is apparent from the foregoing descriptions, the compounds of Formula 1 are useful in a variety of forms, i.e., with various substitutions as identified.

Examples of particularly desirable compounds are quite diverse, and many are mentioned herein. Included are compounds in which R₁ is a "cis-cinnamide" or a "trans-cinnamide", and R₃ is hydrogen; or where R₃ is a "cis-cinnamide" or a "trans-cinnamide", and R₁ is hydrogen, or R₁, R₂, and R₄ are each independently hydrogen or alkyl, and R₅ is halogen, haloalkyl or nitro. Further preferred compounds include those as above wherein R₁₀ and R₁₁ are each independently hydrogen, alkyl, cycloalkyl, alkoxy-carbonyl-alkyl, hydroxy-alkyl, or heterocyclyl-alkyl, or where NR₁₀R₁₁ is heterocyclyl or substituted heterocyclyl, and where Ar is aryl, substituted aryl, heteroaryl, or substituted heteroaryl.

Compounds of the present invention include:

(2,4-Dichlorophenyl)[2-(*E*-((6-hydroxyhexylamino)carbonyl)ethenyl)phenyl] sulfide;
 (2,4-Dichlorophenyl)[2-(*E*-((3-(1-imidazolyl)propylamino)carbonyl)ethenyl)phenyl] sulfide;

(2,4-Dichlorophenyl)[2-chloro-4-(*E*-((2-

hydroxyethylamino)carbonyl)ethenyl)phenyl]

sulfide;

(2,4-Dichlorophenyl)[2-chloro-4-(*E*-((6hydroxyhexylamino)carbonyl)ethenyl)phenyl]

sulfide;

(2,4-Dichlorophenyl)[2-chloro-4-(*E*-((bis-(2-hydroxyethyl)amino)carbonyl)ethenyl)

phenyl] sulfide;

(2,4-Dichlorophenyl)[2-chloro-4-(*E*-((3-(1-pyrrolidin-2-onyl)propylamino)carbonyl)

ethenyl)phenyl] sulfide;

(2,4-Dichlorophenyl)[2-chloro-4-(*E*-((1-morpholinyl)carbonyl)ethenyl)phenyl] sulfide;

(2,4-Dichlorophenyl)[2-chloro-4-(*E*-((4-methylpiperazin-1-yl)carbonyl)ethenyl)phenyl] sulfide;

- 5 (2,4-Dichlorophenyl)[2-chloro-4-(*E*-((4-acetyl) piperazin-1-yl)carbonyl)ethenyl)phenyl] sulfide;

(2,4-Dichlorophenyl)[2-chloro-4-(*E*-((4-(2-pyridyl)piperazin-1-yl)carbonyl)ethenyl)phenyl] sulfide;

- 10 (2-(Hydroxymethyl)phenyl)[2-chloro-4-(*E*-((1-morpholinyl)carbonyl)ethenyl)phenyl] sulfide;

(2-Bromophenyl)[2-chloro-4-(*E*-((1-morpholinyl)carbonyl)ethenyl)phenyl] sulfide;

(2,4-Dichlorophenyl)[2-chloro-4-(*E*-((4-(2-hydroxyethyl)piperazin-1-yl)carbonyl)ethenyl)phenyl] sulfide;

- 15 (2,4-Dichlorophenyl)[2-chloro-4-(*E*-((4-(2-hydroxyethoxyethyl)piperazin-1-yl)carbonyl)ethenyl)phenyl] sulfide;

(2-Bromophenyl)[2-chloro-4-(*E*-((3-(hydroxymethyl)piperidin-1-yl)carbonyl)ethenyl)phenyl] sulfide;

(2-Bromophenyl)[2-chloro-4-(*E*-((2-(hydroxymethyl)piperidin-1-yl)carbonyl)

- 20 ethenyl)phenyl] sulfide;

(2-Bromophenyl)[2-chloro-4-(*E*-((3-acetamidopyrrolidin-1-yl)carbonyl)ethenyl)phenyl] sulfide;

- (2-Bromophenyl)[2-chloro-4-(*E*-((4-hydroxypiperidin-1-yl)carbonyl)ethenyl)phenyl] sulfide;
- (2-Bromophenyl)[2-chloro-4-(*E*-((piperidin-1-yl)carbonyl) ethenyl)phenyl] sulfide;
- (2,4-Dichlorophenyl)[2-chloro-4-(*E*-((3-carboxypiperidin-1-yl)carbonyl)ethenyl)phenyl] sulfide;
- 5 (2,4-Dichlorophenyl)[2-chloro-4-(*E*-((4-carboxypiperidin-1-yl)carbonyl)ethenyl)phenyl] sulfide;
- (2-Bromophenyl)[2-chloro-4-(*E*-((4-acetylhomopiperazin-1-yl)carbonyl)ethenyl)phenyl] sulfide;
- 10 (2-Bromophenyl)[2-chloro-4-(*E*-((thiomorpholin-1-yl)carbonyl)ethenyl)phenyl] sulfide;
- (2-Bromophenyl)[2-chloro-4-(*E*-((4-(1-benzimidazol-2-onyl)piperidin-1-yl)carbonyl)ethenyl)phenyl] sulfide;
- (2-Bromophenyl)[2-chloro-4-(*E*-((2-tetrahydroisoquinoliny)carbonyl)ethenyl)phenyl] sulfide;
- 15 (2-Methylphenyl)[2-trifluoromethyl-4-(*E*-((4-acetylperazin-1-yl)carbonyl)ethenyl)phenyl] sulfide;
- (2-Methylphenyl)[2-trifluoromethyl-4-(*E*-((1-morpholiny)carbonyl)ethenyl)phenyl] sulfide;
- 20 (2-Methylphenyl)[2-trifluoromethyl-4-(*E*-((2-(1-morpholiny)ethylamino)carbonyl)ethenyl)phenyl] sulfide;
- (2-Methylphenyl)[2-trifluoromethyl-4-(*E*-((4-phenylpiperazin-1-yl)carbonyl)

00160 5274560

ethenyl)phenyl] sulfide;

(2-Methylphenyl)[2-trifluoromethyl-4-(*E*-((3-(1-pyrrolidin-2-onyl)propylamino)carbonyl) ethenyl)phenyl] sulfide;

(2-Methylphenyl)[2-trifluoromethyl-4-(*E*-((cyclopropylamino)carbonyl)ethenyl)

5 phenyl] sulfide;

(2,4-Dichlorophenyl)[2-nitro-4-(*E*-((4-acetylpiperazin-1-yl)carbonyl)ethenyl)phenyl] sulfide;

(2,4-Dichlorophenyl)[2-nitro-4-(*E*-((3-(1-pyrrolidin-2-onyl)propylamino)carbonyl) ethenyl)phenyl] sulfide;

10 (2,3-Dichlorophenyl)[2-nitro-4-(*E*-((4-acetylpiperazin-1-yl)carbonyl)ethenyl)phenyl] sulfide;

(4-Bromophenyl)[2-nitro-4-(*E*-((4-acetylpiperazin-1-yl)carbonyl)ethenyl)phenyl] sulfide;

(4-Methylphenyl)[2-nitro-4-(*E*-((4-acetylpiperazin-1-yl)carbonyl)ethenyl)phenyl]

15 sulfide;

(2,4-Dichlorophenyl)[2-nitro-4-(*E*-((4-(*tert*-butoxycarbonyl)piperazin-1-yl)carbonyl) ethenyl)phenyl] sulfide;

(2,4-Dichlorophenyl)[2-nitro-4-(*E*-((4-(2-furoylcarbonyl)piperazin-1-yl)carbonyl) ethenyl)phenyl] sulfide;

20 (2,4-Dichlorophenyl)[2-nitro-4-(*E*-((4-(methanesulfonyl)piperazin-1-yl)carbonyl) ethenyl)phenyl] sulfide;

(2,4-Dichlorophenyl)[2-nitro-4-(*E*-((4-(diethylaminocarbonyl)methyl)piperazin-1-

yl)carbonyl) ethenyl)phenyl] sulfide;

(2,4-Dichlorophenyl)[2-nitro-4-(*E*-((4-(diethylaminocarbonyl)piperazin-1-yl)carbonyl) ethenyl)phenyl] sulfide;

(2,4-Dichlorophenyl)[2-nitro-4-(*E*-((4-(*tert*-butoxycarbonylmethyl)piperazin-1-

5 yl)carbonyl) ethenyl)phenyl] sulfide;

(2,4-Dichlorophenyl)[2-nitro-4-(*E*-((4-(carboxycarbonyl)piperazin-1-yl)carbonyl) ethenyl)phenyl] sulfide;

(2,4-Dichlorophenyl)[2-nitro-4-(*E*-((4-(carboxymethyl)piperazin-1-yl)carbonyl) ethenyl)phenyl] sulfide;

10 (2-Methylphenyl)[2-nitro-4-(*E*-((4-acetylpiperazin-1-yl)carbonyl)ethenyl)phenyl] sulfide;

(2-Chlorophenyl)[2-nitro-4-(*E*-((4-acetylpiperazin-1-yl)carbonyl)ethenyl)phenyl] sulfide;

(2-Aminophenyl)[2-nitro-4-(*E*-((4-acetylpiperazin-1-yl)carbonyl)ethenyl)phenyl] sulfide;

15

(2-Hydroxymethylphenyl)[2-nitro-4-(*E*-((4-acetylpiperazin-1-yl)carbonyl)ethenyl)phenyl] sulfide;

(2-Ethylphenyl)[2-nitro-4-(*E*-((4-acetylpiperazin-1-yl)carbonyl)ethenyl)phenyl] sulfide;

20 (2-*iso*-Propylphenyl)[2-nitro-4-(*E*-((4-acetylpiperazin-1-yl)carbonyl)ethenyl)phenyl] sulfide;

(2-*tert*-Butylphenyl)[2-nitro-4-(*E*-((4-acetylpiperazin-1-yl)carbonyl)ethenyl)phenyl]

0344795 003100
DATE: 06/27/90

sulfide;

(2-Chlorophenyl)[2-chloro-4-(*E*-((4-acetylpiperazin-1-yl)carbonyl))2-propenyl]phenyl] sulfide;

(2-(1-Morpholinylmethyl)phenyl)[2-chloro-4-(*E*-((1-morpholinyl)carbonyl) ethenyl)

5 phenyl] sulfide;

(2-(4-(1,3-Benzodioxolyl-5-methyl)piperazin-1-ylmethyl)phenyl)[2-chloro-4-(*E*-((1-morpholinyl)carbonyl) ethenyl)phenyl] sulfide;

(2-(4-(*iso*-Propylaminocarbonylmethyl)piperazin-1-ylmethyl)phenyl)[2-chloro-4-(*E*-((1-morpholinyl)carbonyl) ethenyl)phenyl] sulfide;

10 (2-((*N*-Ethoxycarbonylmethyl-*N*-methyl)aminomethyl)phenyl)[2-chloro-4-(*E*-((1-morpholinyl)carbonyl) ethenyl)phenyl] sulfide;

(2-Formylphenyl)[2-chloro-4-(*E*-((1-morpholinyl)carbonyl)ethenyl)phenyl] sulfide;

(2-(4-Formylpiperazin-1-ylmethyl)phenyl)[2-chloro-4-(*E*-((1-morpholinyl)carbonyl) ethenyl)phenyl] sulfide;

15 (2-(*E*-((1-Morpholinyl)carbonyl)ethenyl)phenyl)[2-chloro-4-(*E*-((1-morpholinyl)carbonyl)ethenyl)phenyl] sulfide;

(2-Formylphenyl)[2-nitro-4-(*E*-((4-acetylpiperazin-1-yl)carbonyl)ethenyl)phenyl] sulfide;

(2-Formylphenyl)[2-chloro-4-(*E*-((1-morpholinyl)carbonyl)ethenyl)phenyl] sulfide,

20 *N,N*-dimethyl hydrazone;

(2-((3-(1-Morpholinyl)propyl)-1-amino)phenyl)[2-chloro-4-(*E*-((1-morpholinyl)carbonyl) ethenyl)phenyl] sulfide;

(2,4-Dichlorophenyl)[2-bromo-4-(*E*-((3-(1-pyrrolidin-2-yl)propylamino)carbonyl)ethenyl)phenyl] sulfide;

(2,4-Dichlorophenyl)[2-formyl-4-(*E*-((1-morpholinyl)carbonyl)ethenyl)phenyl] sulfide;

- 5 (2-Chloro-6-formylphenyl)[2-chloro-4-(*E*-((4-acetylpiperazin-1-yl)carbonyl)ethenyl)phenyl] sulfide;

(2-Cyanophenyl)[2-chloro-4-(*E*-((4-acetylpiperazin-1-yl)carbonyl)ethenyl)phenyl] sulfide;

(2-Isopropylphenyl)[2-cyano-4-(*E*-((morpholin-1-yl)carbonyl)

- 10 ethenyl)phenyl] sulfide;

(2-Bromophenyl)[2-nitro-4-(*E*-((4-acetylpiperazin-1-yl)carbonyl)

ethenyl)phenyl] sulfide;

(2-(Pyrrolidin-1-yl)phenyl)[2-chloro-4-(*E*-((morpholin-1-yl)carbonyl)

ethenyl)phenyl] sulfide;

- 15 (2-Methoxyphenyl)-[2-chloro-4(*E*-[(morpholin-1-yl)carbonyl]ethenyl)phenyl]sulfide;

(2-Isopropylphenyl)[2-nitro-4-(*E*-((3-carbomethoxypiperazin-1-yl)carbonyl)

ethenyl)phenyl] sulfide;

(2-Methylphenyl)[2-nitro-4-(*E*-((3-carboxamido-4-carbobenzoxypiperazin-1-yl)carbonyl)ethenyl)phenyl] sulfide;

- 20 (2-Isopropylphenyl)[2-nitro-4-(*E*-((2-carbomethoxy-4-*tert*-butoxycarbonylpiperazin-1-yl)carbonyl)ethenyl)phenyl] sulfide;

- (2-Isopropylphenyl)[2-nitro-4-(*E*-((2-carboxy-4-*tert*-butoxycarbonylpiperazin-1-yl)carbonyl)ethenyl) phenyl] sulfide;
- (2-Isopropylphenyl)[2-trifluoromethyl-4-(*E*-((4-acetylpiperazin-1-yl)carbonyl)ethenyl) phenyl] sulfide;
- 5 (2-Isopropylphenyl)[2-trifluoromethyl-4-(*E*-((morpholin-1-yl)carbonyl)ethenyl) phenyl] sulfide;
- (2-Isopropylphenyl)[2-trifluoromethyl-4-(*E*-((3-(pyrrolidin-2-on-1-yl)prop-1-ylamino)carbonyl)ethenyl)phenyl]sulfide;
- (2-Isopropylphenyl)[2-trifluoromethyl-4-(*E*-((cyclobutylamino)carbonyl)ethenyl) phenyl] sulfide;
- 10 (2-Isopropylphenyl)[2-trifluoromethyl-4-(*E*-((cyclopentylamino)carbonyl)ethenyl) phenyl] sulfide;
- (2-Isopropylphenyl)[2-trifluoromethyl-4-(*E*-((5-hydroxypent-1-ylamino)carbonyl)ethenyl) phenyl] sulfide;
- 15 (2-Isopropylphenyl)[2-nitro-4-(*E*-((3-carbomethoxy-4-acetylpiperazin-1-yl)carbonyl)ethenyl) phenyl] sulfide;
- (2-Biphenyl)[2-chloro-4-(*E*-((morpholin-1-yl)carbonyl)ethenyl) phenyl] sulfide;
- (3,4-Dimethylphenyl)[2-nitro-4-(*E*-((4-acetylpiperazin-1-yl)carbonyl)ethenyl)phenyl]sulfide;
- 20 (2-Bromophenyl)[2-trifluoromethyl-4-(*E*-((4-acetylpiperazin-1-yl)carbonyl)ethenyl) phenyl] sulfide;

001100-56274560

(5-Indolyl)[2-chloro-4-(*E*-((4-acetylpiperazin-1-yl)carbonyl)ethenyl) phenyl] sulfide;

(5-Benzodioxolyl)[2-chloro-4-(*E*-((4-acetylpiperazin-1-yl)carbonyl)ethenyl) phenyl] sulfide;

5 (2-Isopropylphenyl)[2-nitro-4-(*E*-((2-carbomethoxypiperazin-1-yl)carbonyl)ethenyl) phenyl] sulfide;

(2,3-Dimethoxyphenyl)-[2-chloro-4(*E*-[(morpholin-1-yl)carbonyl]ethenyl)phenyl] sulfide;

(2-Fluorophenyl)[2-nitro-4-(*E*-((4-acetylpiperazin-1-yl)carbonyl)ethenyl)phenyl]sulfide;

10 (2-Bromophenyl)[2-trifluoromethyl-4-(*E*-((4-*tert*-butoxycarbonyl)piperazin-1-yl)carbonyl)ethenyl) phenyl] sulfide;

(2-(Pyrrolidin-1-yl)phenyl)[2-trifluoromethyl-4-(*E*-((4-*tert*-butoxycarbonyl)piperazin-1-yl)carbonyl)ethenyl) phenyl] sulfide;

15 (3-Carboxamidophenyl)[2-nitro-4-(*E*-((4-acetylpiperazin-1-yl)carbonyl)ethenyl) phenyl] sulfide;

(3-(Hydroxymethyl)phenyl)[2-nitro-4-(*E*-((4-acetylpiperazin-1-yl)carbonyl)ethenyl) phenyl] sulfide;

Phenyl[2-trifluoromethyl-4-(*E*-((4-*tert*-butoxycarbonyl)piperazin-1-yl)carbonyl)ethenyl) phenyl] sulfide;

(2-Isopropylphenyl)[2-trifluoromethyl-4-(*E*-((2-carbomethoxy-4-*tert*-butoxycarbonyl)piperazin-1-yl)carbonyl)ethenyl) phenyl] sulfide;

(2-Isopropylphenyl)[2-nitro-4-(*E*-((3-(pyridine-4-methylaminocarbonyl)-4-*tert*-butoxycarbonyl)piperazin-1-yl)carbonyl)ethenyl) phenyl] sulfide;

(2-Ethoxyphenyl)-[2-chloro-4(E-[(morpholin-1-yl)carbonyl]ethenyl)phenyl]sulfide;

(2-Methoxyphenyl)[2-nitro-4-(*E*-((4-acetylpiperazin-1-

5 yl)carbonyl)ethenyl)phenyl]sulfide;

(2-(Azetidin-1-yl)phenyl)[2-trifluoromethyl-4-(*E*-((4-(*tert*-

butoxycarbonyl)piperazin-1-yl)carbonyl)ethenyl) phenyl] sulfide;

(2-(Piperidin-1-yl)phenyl)[2-trifluoromethyl-4-(*E*-((4-(*tert*-butoxycarbonyl)piperazin-1-yl)carbonyl)ethenyl) phenyl] sulfide;

10 (3-Chloro-2-formylphenyl)[2-chloro-4-(*E*-((4-acetylpiperazin-1-yl)carbonyl)ethenyl) phenyl] sulfide;

(2-Trifluoromethylphenyl)[2-trifluoromethyl-4-(*E*-((4-acetylpiperazin-1-yl)carbonyl)ethenyl) phenyl] sulfide;

(3-Bromophenyl)[2-trifluoromethyl-4-(*E*-((4-acetylpiperazin-1-yl)carbonyl)

15 etenyl) phenyl] sulfide;

(3,5-Dimethylphenyl)[2-trifluoromethyl-4-(*E*-((4-acetylpiperazin-1-yl)carbonyl)ethenyl) phenyl] sulfide;

(2-Isopropylphenyl)[2-nitro-4-(*E*-((3-dimethylaminocarbonyl)-4-(pyridine-4-carbonyl)piperazin-1-yl)carbonyl)ethenyl) phenyl] sulfide;

20 (2-Isopropylphenyl)[2-nitro-4-(*E*-((3-dimethylaminocarbonyl-4-carbomethoxypiperazin-1-yl)carbonyl)ethenyl) phenyl] sulfide;

- (2-Isopropylphenyl)[2-nitro-4-(*E*-((3-dimethylaminocarbonyl-4-acetylpiperazin-1-yl)carbonyl)ethenyl) phenyl] sulfide;
- (2-Isopropylphenyl)[2-nitro-4-(*E*-((3-(1-morpholinocarbonyl)-4-*tert*-butoxycarbonylpiperazin-1-yl)carbonyl)ethenyl) phenyl] sulfide;
- 5 (2-Isopropylphenyl)[2-nitro-4-(*E*-((3-(pyridine-4-methylaminocarbonyl)piperazin-1-yl)carbonyl)ethenyl) phenyl] sulfide;
- (2-Isopropylphenyl)[2-nitro-4-(*E*-((3-dimethylaminocarbonyl)piperazin-1-yl)carbonyl)ethenyl) phenyl] sulfide;
- (2-Isopropylphenyl)[2-nitro-4-(*E*-((3-(benzylaminocarbonyl)-4-*tert*-butoxycarbonylpiperazin-1-yl)carbonyl)ethenyl) phenyl] sulfide;
- 10 (2-Isopropylphenyl)[2-nitro-4-(*E*-((3-(dimethylaminocarbonyl)-4-*tert*-butoxycarbonylpiperazin-1-yl)carbonyl)ethenyl) phenyl] sulfide;
- (2-Bromophenyl)[2-chloro-4-(*E*-((3-(5*S*-hydroxymethyl-pyrrolidin-2-on-1-yl)prop-1-ylamino)carbonyl) ethenyl)phenyl]sulfide;
- 15 (2-Bromophenyl)[2-chloro-4-(*E*-((3-(pyrrolidin-2-on-1-yl)prop-1-ylamino)carbonyl) ethenyl)phenyl]sulfide;
- (2-Bromophenyl)[2-chloro-4-(*E*-(*N*-methyl-*N*-(3-(pyrrolidin-2-on-1-yl)prop-1-ylamino)carbonyl) ethenyl)phenyl]sulfide;
- (2-[2-Methoxy]ethoxyphenyl)-[2-chloro-4(*E*-[(morpholin-1-yl)carbonyl]ethenyl)phenyl] sulfide;
- 20 (2-Isopropylphenyl)[2-nitro-4-(*E*-((3-(morpholinocarbonyl)piperazin-1-yl)carbonyl)ethenyl) phenyl] sulfide;

- (2-Isopropylphenyl)[2-nitro-4-(*E*-((4-*tert*-butoxycarbonyl)piperazin-1-yl)carbonyl)ethenyl] phenyl] sulfide;
- (2-Isopropylphenyl)[2-nitro-4-(*E*-((4-methoxycarbonyl)piperazin-1-yl)carbonyl)ethenyl] phenyl] sulfide;
- 5 (2-Isopropylphenyl)[2-nitro-4-(*E*-(4-(pyridine-4-carbonyl)piperazin-1-yl)carbonyl)ethenyl] phenyl] sulfide;
- (2-Isopropylphenyl)[2-nitro-4-(*E*-((3-(pyridine-3-methylaminocarbonyl)-4-*tert*-butoxycarbonyl)piperazin-1-yl)carbonyl)ethenyl] phenyl] sulfide;
- (2-Isopropylphenyl)[2-nitro-4-(*E*-((3-(pyridine-2-methylaminocarbonyl)piperazin-1-yl)carbonyl)ethenyl] phenyl] sulfide;
- 10 (2-Isopropylphenyl)[2-nitro-4-(*E*-((3-(pyridine-3-methylaminocarbonyl)piperazin-1-yl)carbonyl)ethenyl] phenyl] sulfide;
- (4-Hydroxyphenyl)[2-nitro-4-(*E*-((4-acetyl)piperazin-1-yl)carbonyl)ethenyl]phenyl]sulfide;
- 15 (3,5-Dichlorophenyl)[2-nitro-4-(*E*-((4-acetyl)piperazin-1-yl)carbonyl)ethenyl]phenyl]sulfide;
- (2-Bromophenyl)[2-chloro-4-(*E*-((3-(5*S*-acetoxymethyl-pyrrolidin-2-on-1-yl)prop-1-ylamino)carbonyl) ethenyl]phenyl]sulfide;
- (2-Bromophenyl)[2-chloro-4-(*E*-((3-(5*S*-methoxymethyl-pyrrolidin-2-on-1-yl)prop-1-ylamino)carbonyl) ethenyl]phenyl]sulfide;
- 20 (2-Bromophenyl)[2-chloro-4-(*E*-((3-(4*R*-hydroxymethyl-pyrrolidin-2-on-1-yl)prop-1-ylamino)carbonyl) ethenyl]phenyl]sulfide;

- Phenyl[2-nitro-4-(*E*-((4-acetylpiperazin-1-yl)carbonyl)ethenyl)phenyl]sulfide;
 (2-Dimethylaminophenyl)[2-nitro-4-(*E*-((4-acetylpiperazin-1-yl)carbonyl)
 ethenyl) phenyl] sulfide;
 (3-((2-Hydroxyethyl)aminocarbonyl)phenyl)[2-nitro-4-(*E*-((4-acetylpiperazin-1-
 5 yl)carbonyl)ethenyl) phenyl] sulfide;
 (3-((3-(1-Imidazolyl)propyl)aminocarbonyl)phenyl)[2-nitro-4-(*E*-((4-
 acetylpiperazin-1-yl)carbonyl)ethenyl) phenyl] sulfide;
 (3-((2-(1-Morpholinyl)ethyl)aminocarbonyl)phenyl)[2-nitro-4-(*E*-((4-
 acetylpiperazin-1-yl)carbonyl)ethenyl) phenyl] sulfide;
 10 (2-Isopropylphenyl)[2-nitro-4-(*E*-((3-hydroxymethyl-4-*tert*-
 butoxycarbonylpiperazin-1-yl)carbonyl)ethenyl) phenyl] sulfide;
 (2-Isopropylphenyl)[2-nitro-4-(*E*-((4-formylpiperazin-1-yl)carbonyl)ethenyl) phenyl]
 sulfide;
 (2-Isopropylphenyl)[2-nitro-4-(*E*-((2-hydroxymethyl-4-*tert*-
 15 butoxycarbonylpiperazin-1-yl)carbonyl)ethenyl) phenyl] sulfide;
 (2-Ethoxyphenyl)-[2-chloro-4-(*E*-[(3-ethoxycarbonylpiperidin-1-yl)carbonyl]ethenyl)
 phenyl]sulfide;
 (3- Aminophenyl)[2-nitro-4-(*E*-((4-acetylpiperazin-1-
 yl)carbonyl)ethenyl)phenyl]sulfide;
 20 (4-Aminophenyl)[2-nitro-4-(*E*-((4-acetylpiperazin-1-
 yl)carbonyl)ethenyl)phenyl]sulfide;

- (2,4-Dimethylphenyl)[2-nitro-4-(*E*-((4-acetylpiperazin-1-yl)carbonyl)ethenyl)phenyl]sulfide;
- (2,5-Dimethylphenyl)[2-nitro-4-(*E*-((4-acetylpiperazin-1-yl)carbonyl)ethenyl)phenyl]sulfide;
- 5 (4-Methoxyphenyl)[2-nitro-4-(*E*-((4-acetylpiperazin-1-yl)carbonyl)ethenyl)phenyl]sulfide;
- (3-Chlorophenyl)[2-nitro-4-(*E*-((4-acetylpiperazin-1-yl)carbonyl)ethenyl)phenyl]sulfide;
- (2-Chloro, 4,5-diaminophenyl)[2-chloro-4-(*E*-((4-acetylpiperazin-1-yl)carbonyl)ethenyl) phenyl] sulfide;
- 10 (3,4-Diaminophenyl)[2-chloro-4-(*E*-((4-acetylpiperazin-1-yl)carbonyl)ethenyl) phenyl]
- (6-Chlorobenzimidazol-2-on-5-yl)[2-chloro-4-(*E*-((4-acetylpiperazin-1-yl)carbonyl)ethenyl) phenyl] sulfide;
- 15 (1-Methylindol-7-yl)[2-chloro-4-(*E*-((4-acetylpiperazin-1-yl)carbonyl)ethenyl) phenyl] sulfide;
- (2-Hydroxy, 4-aminophenyl)[2-chloro-4-(*E*-((4-acetylpiperazin-1-yl)carbonyl)ethenyl) phenyl] sulfide;
- (2-Isopropylphenyl)[2-nitro-4-(*E*-((4-methylpiperazin-1-yl)carbonyl)ethenyl) phenyl] sulfide;
- 20 (2-Isopropylphenyl)[2-nitro-4-(*E*-((4-(pyridine-2-carbonyl)piperazin-1-yl)carbonyl)ethenyl) phenyl] sulfide;

- (2-Isopropylphenyl)[2-nitro-4-(*E*-((4-(pyridine-3-carbonyl)piperazin-1-yl)carbonyl)ethenyl) phenyl] sulfide;
- (2-Isopropylphenyl)[2-nitro-4-(*E*-((2-carbomethoxy-4-methoxycarbonylpiperazin-1-yl)carbonyl)ethenyl) phenyl] sulfide;
- 5 (2-Isopropylphenyl)[2-nitro-4-(*E*-((2-carboxy-4-methoxycarbonylpiperazin-1-yl)carbonyl)ethenyl) phenyl] sulfide;
- (2-Isopropylphenyl)[2-nitro-4-(*E*-((3-carbomethoxy-4-methylpiperazin-1-yl)carbonyl)ethenyl) phenyl] sulfide;
- (2-Ethoxyphenyl)-[2-chloro-4(*E*-[(3-carboxypiperidin-1-yl)carbonyl]ethenyl)phenyl]
- 10 sulfide;
- (2-Ethoxyphenyl)-[2-chloro-4(*E*-[(3-carboxypiperidin-1-yl)carbonyl]ethenyl)phenyl] sulfide;
- (2-Ethoxyphenyl)-[2-chloro-4(*E*-[(2-ethoxycarbonylpiperidin-1-yl)carbonyl]ethenyl)phenyl]sulfide;
- 15 (2-Ethoxyphenyl)[2-trifluoromethyl-4-(*E*-((1-(*tert*-butoxycarbonyl)-4-hydroxypyrrolidin-3-ylamino)carbonyl)ethenyl) phenyl] sulfide;
- (2-Ethoxyphenyl)-[2-chloro-4(*E*-[(2-carboxypiperidin-1-yl)carbonyl]ethenyl)phenyl] sulfide;
- (2-Ethoxyphenyl)[2-trifluoromethyl-4-(*E*-(((pyrrol-3-in-1-yl)carbonyl)ethenyl)
- 20 phenyl] sulfide;
- (2-Ethoxyphenyl)[2-trifluoromethyl-4-(*E*-((3-(pyrrolidin-2-on-1-yl)prop-1-ylamino)carbonyl) ethenyl)phenyl]sulfide;

054175: 03100
00100: 5674460

- (2-Ethoxyphenyl)[2-trifluoromethyl-4-(*E*-((4-acetyl)piperazin-1-yl)carbonyl)ethenyl) phenyl] sulfide;
- (2-Ethoxyphenyl)[2-trifluoromethyl-4-(*E*-((4-(ethoxycarbonyl)piperazin-1-yl)carbonyl)ethenyl) phenyl] sulfide;
- 5 (2-Ethoxyphenyl)[2-trifluoromethyl-4-(*E*-((4-(2-furylcarbonyl)piperazin-1-yl)carbonyl)ethenyl) phenyl] sulfide;
- (2-Ethoxyphenyl)-[2-chloro-4(*E*-[(3-ethoxycarbonylpiperidin-1-yl)carbonyl]ethenyl) phenyl]sulfide;
- (2-Ethoxyphenyl)-[2-chloro-4(*E*-[(4-carboxypiperidin-1-yl)carbonyl]ethenyl)phenyl]
- 10 sulfide;
- (Benzodioxan-6-yl)[2-chloro-4-(*E*-((4-acetyl)piperazin-1-yl)carbonyl)ethenyl) phenyl] sulfide;
- (2-Isopropylphenyl)[2-nitro-4-(*E*-((4-ethoxycarbonyl)piperazin-1-yl)carbonyl)ethenyl) phenyl] sulfide;
- 15 (2-Isopropylphenyl)[2-nitro-4-(*E*-((4-isopropoxycarbonyl)piperazin-1-yl)carbonyl)ethenyl) phenyl] sulfide;
- (2-Isopropylphenyl)[2-nitro-4-(*E*-((4-isobutoxycarbonyl)piperazin-1-yl)carbonyl)ethenyl) phenyl] sulfide;
- (2-Isopropylphenyl)[2-nitro-4-(*E*-((4-(1-propen-2-oxy)carbonyl)piperazin-1-yl)carbonyl)ethenyl) phenyl] sulfide;
- 20 (2-Isopropylphenyl)[2-nitro-4-(*E*-((4-propionylpiperazin-1-yl)carbonyl)ethenyl) phenyl] sulfide;

- (2-Isopropylphenyl)[2-nitro-4-(*E*-((4-carboxamidopiperazin-1-yl)carbonyl)ethenyl)phenyl] sulfide;
- (2-Isopropylphenyl)[2-nitro-4-(*E*-((4-methylaminocarbonyl)piperazin-1-yl)carbonyl)ethenyl) phenyl] sulfide;
- 5 (2-Isopropylphenyl)[2-nitro-4-(*E*-((4-(pyrimidin-2-yl)piperazin-1-yl)carbonyl)ethenyl) phenyl] sulfide;
- (2-Isopropylphenyl)[2-nitro-4-(*E*-((4-hydroxyacetyl)piperazin-1-yl)carbonyl)ethenyl) phenyl] sulfide;
- (2-Isopropylphenyl)[2-nitro-4-(*E*-((4-(pyrazine-2-carbonyl)piperazin-1-yl)carbonyl)ethenyl) phenyl] sulfide;
- 10 (2-Isopropylphenyl)[2-trifluoromethyl-4-(*E*-((2-carboxypyrrol-3-in-1-yl)carbonyl)ethenyl) phenyl] sulfide;
- (2-Isopropylphenyl)[2-trifluoromethyl-4-(*E*-((3-hydroxymethyl-4-methylpiperazin-1-yl)carbonyl)ethenyl) phenyl] sulfide;
- 15 (2-Isopropylphenyl)[2-trifluoromethyl-4-(*E*-(((2-carboxypyrrol-3-in-1-yl)carbonyl)ethenyl) phenyl] sulfide;
- (2-Isopropylphenyl)[2-trifluoromethyl-4-(*E*-(((2-hydroxymethylpyrrolidin-1-yl)carbonyl)ethenyl) phenyl] sulfide;
- (2-Isopropylphenyl)[2-nitro-4-(*E*-((3-methylaminocarbonyl)piperazin-1-yl)carbonyl)ethenyl) phenyl] sulfide;
- 20 (2-Isopropylphenyl)[2-nitro-4-(*E*-(((3-cyclopropylaminocarbonyl)piperazin-1-yl)carbonyl)ethenyl) phenyl] sulfide;

(2-Isopropylphenyl)[2-nitro-4-(*E*-((3-carboxamidopiperazin-1-yl)carbonyl)ethenyl)phenyl] sulfide;

(2-Isopropylphenyl)[2-nitro-4-(*E*-((3-carbomethoxy-4-oxopiperidin-1-yl)carbonyl)ethenyl)phenyl] sulfide;

- 5 (2-Isopropylphenyl)[2-nitro-4-(*E*-((3,5-dimethylpiperazin-1-yl)carbonyl)ethenyl)phenyl] sulfide;

(1-Ethylindol-7-yl)[2-chloro-4-(*E*-((4-acetyl)piperazin-1-yl)carbonyl)ethenyl)phenyl] sulfide;

- (3-[2-Methoxy]ethoxyphenyl)-[2-chloro-4(*E*-[(morpholin-1-yl)carbonyl]ethenyl)phenyl] sulfide;
- 10 (2-Bromophenyl)[2-chloro-4-(*E*-((4,4'-*S*-dioxythiomorpholin-1-yl)carbonyl)ethenyl)phenyl]sulfide;

(2-Bromophenyl)[2-chloro-4-(*E*-(*N*-carbomethoxymethyl-*N*-(3-(pyrrolidin-2-on-1-yl)prop-1-yl)amino)carbonyl)ethenyl)phenyl]sulfide;

- 15 (2-Bromophenyl)[2-chloro-4-(*E*-((4-*S*-oxythiomorpholin-1-yl)-2-pyrrolidinone)carbonyl)ethenyl)phenyl]sulfide;

(2-Methoxy-5-chlorophenyl)[2-nitro-4-(*E*-((4-acetyl)piperazin-1-yl)carbonyl)ethenyl)phenyl] sulfide;

- (2-Isopropylphenyl)[2-nitro-4-(*E*-((3-acetoxymethyl)piperazin-1-yl)carbonyl)ethenyl)phenyl] sulfide;
- 20

(2-Isopropylphenyl)[2-nitro-4-(*E*-((3,5-dimethyl-4-acetyl)piperazin-1-yl)carbonyl)ethenyl)phenyl] sulfide;

- (1-Methylindol-5-yl)[2-chloro-4-(*E*-((4-acetylpiperazin-1-yl)carbonyl)ethenyl)phenyl] sulfide;
- (Benzodioxan-6-yl)[2-nitro-4-(*E*-((4-acetylpiperazin-1-yl)carbonyl)ethenyl) phenyl] sulfide;
- 5 (Benzodioxan-6-yl)[2-nitro-4-(*E*-((3-(pyrrolidin-2-on-1-yl)prop-1-ylamino)carbonyl)ethenyl)phenyl]sulfide;
- (Benzodioxan-6-yl)[2-nitro-4-(*E*-((3-carboethoxypiperidin-1-yl) carbonyl)ethenyl) phenyl] sulfide;
- (Benzodioxan-6-yl)[2-nitro-4-(*E*-((4-carboethoxypiperidin-1-yl) carbonyl)ethenyl) phenyl] sulfide;
- 10 (2-Ethoxyphenyl)[2-trifluoromethyl-4-(*Z*-((4-acetylpiperazin-1-yl)carbonyl)ethenyl) phenyl] sulfide;
- (2-Ethoxyphenyl)[2-trifluoromethyl-4-(*E*-((6-methylpyrid-2-ylamino)carbonyl)ethenyl) phenyl] sulfide;
- 15 (2-Methyl-3-chlorophenyl)[2-nitro-4-(*E*-((4-acetylpiperazin-1-yl)carbonyl)ethenyl) phenyl] sulfide;
- (Benzodioxan-6-yl)[2-nitro-4-(*E*-((3-carboxamidopiperidin-1-yl) carbonyl)ethenyl) phenyl] sulfide;
- (Benzodioxan-6-yl)[2-nitro-4-(*E*-((2-carboethoxypiperidin-1-yl) carbonyl)ethenyl) phenyl] sulfide;
- 20 (Benzodioxan-6-yl)[2-nitro-4-(*E*-((4-carboxamidopiperidin-1-yl) carbonyl)ethenyl) phenyl] sulfide;

00150-5674560 03544795-033100

(Benzodioxan-6-yl)[2-nitro-4-(*E*-((4-*tert*-butoxycarbonyl)piperazin-1-yl) carbonyl)ethenyl) phenyl] sulfide;

(2-Isopropylphenyl)[2-nitro-4-(*E*-((*syn*-3,5-dimethylmorpholin-1-yl)carbonyl)ethenyl) phenyl] sulfide;

- 5 (2-Isopropylphenyl)[2-nitro-4-(*E*-((*anti*-3,5-dimethylmorpholin-1-yl)carbonyl)ethenyl) phenyl] sulfide;

(2-Isopropylphenyl)[2-nitro-4-(*E*-((3-carboethoxypiperazin-1-yl)carbonyl)ethenyl) phenyl] sulfide;

- (2-Isopropylphenyl)[2-nitro-4-(*E*-((3-isopropoxycarbonyl)piperazin-1-yl)carbonyl)ethenyl) phenyl] sulfide;

(2-Isopropylphenyl)[2-nitro-4-(*E*-((3-(dimethylaminocarbonyl)-4-methylpiperazin-1-yl)carbonyl)ethenyl) phenyl] sulfide;

(2-Isopropylphenyl)[2-nitro-4-(*E*-((3-carbomethoxy-4-hydroxypiperidin-1-yl)carbonyl)ethenyl) phenyl] sulfide;

- 15 (2-Isopropylphenyl)[2-nitro-4-(*E*-((3-hydroxymethyl-4-hydroxypiperidin-1-yl)carbonyl)ethenyl) phenyl] sulfide;

(2-Ethoxyphenyl)[2-trifluoromethyl-4-(*E*-((2-carbomethoxy-4-(methoxycarbonyl)piperazin-1-yl)carbonyl)ethenyl) phenyl] sulfide;

(2-Ethoxyphenyl)[2-trifluoromethyl-4-(*E*-((2-carbomethoxy-4-methyl piperazin-1-yl)carbonyl)ethenyl) phenyl] sulfide;

- 20 (2-Ethoxyphenyl)[2-trifluoromethyl-4-(*E*-((2-carboxy-4-(methoxycarbonyl)piperazin-1-yl)carbonyl)ethenyl) phenyl] sulfide;

- (Indol-6-yl)[2-chloro-4-(*E*-((4-acetylpiperazin-1-yl)carbonyl)ethenyl) phenyl] sulfide;
(1-Ethyl,3-(dimethylaminomethyl)indol-7-yl)[2-chloro-4-(*E*-((4-acetylpiperazin-1-yl)carbonyl)ethenyl) phenyl] sulfide;
(5-Ethoxybenzodioxan-6-yl)[2-chloro-4-(*E*-((4-acetylpiperazin-1-yl)carbonyl)ethenyl) phenyl] sulfide;
5 (2-Ethyl-4-bromophenyl)[2-nitro-4-(*E*-((4-acetylpiperazin-1-yl)carbonyl)ethenyl) phenyl] sulfide;
(Benzodioxan-6-yl)[2-nitro-4-(*E*-((2-carboxypiperidin-1-yl) carbonyl)ethenyl) phenyl] sulfide;
10 (Benzodioxan-6-yl)[2-nitro-4-(*E*-((4-carboxymethylpiperazin-1-yl) carbonyl)ethenyl) phenyl] sulfide;
(3-Morpholinophenyl)[2-nitro-4-(*E*-((4-acetylpiperazin-1-yl)carbonyl)ethenyl) phenyl] sulfide;
(5-Ethoxybenzodioxan-8-yl)[2-chloro-4-(*E*-((4-acetylpiperazin-1-yl)carbonyl)ethenyl) phenyl] sulfide;
15 (5-Chloro-8-ethoxyquinolin-7-yl)[2-chloro-4-(*E*-((4-acetylpiperazin-1-yl)carbonyl)ethenyl) phenyl] sulfide;
(2-Isopropylphenyl)[2-nitro-4-(*E*-((3-carboethoxypiperidin-1-yl)carbonyl)ethenyl) phenyl] sulfide;
20 (2-Isopropylphenyl)[2-nitro-4-(*E*-((3-carboxypiperidin-1-yl)carbonyl)ethenyl) phenyl] sulfide;

- (2-Isopropylphenyl)[2-nitro-4-(*E*-((3-ethanesulfonylamino)carbonyl)piperidin-1-yl)carbonyl)ethenyl phenyl] sulfide;
- (2-Isopropylphenyl)[2-nitro-4-(*E*-((3-(4-methylpiperazine)sulfonylamino)carbonyl)piperidin-1-yl)carbonyl)ethenyl phenyl] sulfide;
- 5 (2-Isopropylphenyl)[2-nitro-4-(*E*-((3-*p*-toluenesulfonylamino)carbonyl)piperidin-1-yl)carbonyl)ethenyl phenyl] sulfide;
- (2-Isopropylphenyl)[2-nitro-4-(*E*-((3-methyl-4-acetyl)piperazin-1-yl)carbonyl)ethenyl phenyl] sulfide;
- (2-Hydroxyphenyl)-[2-chloro-4(*E*-[(morpholin-1-yl)carbonyl)ethenyl)phenyl]sulfide
- 10 (1-(Carboxymethyl)indol-5-yl)[2-chloro-4-(*E*-((4-acetyl)piperazin-1-yl)carbonyl)ethenyl phenyl] sulfide;
- (Benzodioxan-6-yl)[2-trifluoromethyl-4-(*E*-((4-acetyl)piperazin-1-yl)carbonyl)ethenyl phenyl] sulfide;
- (2-Isopropylphenyl)[2-nitro-4-(*E*-((3-(1-pyrrolidin-2-onyl)prop-1-ylamino)carbonyl)ethenyl phenyl] sulfide;
- 15 carbonyl)ethenyl phenyl] sulfide;
- (3-(2-Morpholinoethylamino)phenyl)[2-trifluoromethyl-4-(*E*-((4-acetyl)piperazin-1-yl)carbonyl)ethenyl phenyl] sulfide;
- (2-Pyrrolidin-1-ylphenyl)[2-nitro-4-(*E*-((4-acetyl)piperazin-1-yl)carbonyl)ethenyl phenyl] sulfide;
- 20 (3-Bromophenyl)[2-nitro-4-(*E*-((3-carboethoxypyrrolidin-1-yl)carbonyl)ethenyl phenyl] sulfide;

00541795.033100

(3-Bromophenyl)[2-nitro-4-(*E*-((4-carboethoxypyrrolidin-1-yl)carbonyl)ethenyl)phenyl] sulfide;

(2-(Hydroxymethyl)-benzodioxan-6-yl)[2-chloro-4-(*E*-((4-acetyl)piperazin-1-yl)carbonyl)ethenyl] phenyl] sulfide;

- 5 (Benzodioxan-6-yl)[2-trifluoromethyl-4-(*E*-((3-(pyrrolidin-2-on-1-yl)prop-1-ylamino)carbonyl) ethenyl)phenyl]sulfide;

(3-(Dimethylaminomethyl)indol-5-yl)[2-chloro-4-(*E*-((4-acetyl)piperazin-1-yl)carbonyl)ethenyl] phenyl] sulfide;

(2-Isopropylphenyl)[2-nitro-4-(*E*-((2-carboethoxypiperidin-1-yl)carbonyl)ethenyl)phenyl] sulfide;

- 10 (2-Isopropylphenyl)[2-nitro-4-(*E*-((2-carboxypiperidin-1-yl)carbonyl)ethenyl)phenyl] sulfide;

(2-Isopropylphenyl)[2-nitro-4-(*E*-((4-carboethoxypiperidin-1-yl)carbonyl)ethenyl)phenyl] sulfide;

- 15 (2-Isopropylphenyl)[2-nitro-4-(*E*-((4-carboxypiperidin-1-yl)carbonyl)ethenyl)phenyl] sulfide;

(2-Isopropylphenyl)[2-nitro-4-(*E*-(((4-*p*-toluenesulfonylaminocarbonyl)piperidin-1-yl)carbonyl)ethenyl) phenyl] sulfide;

(2-Isopropylphenyl)[2-nitro-4-(*E*-((3-carboxy-4-hydroxypiperidin-1-

- 20 yl)carbonyl)ethenyl) phenyl] sulfide;

(Benzodioxan-6-yl)[2-trifluoromethyl-4-(*E*-((3-carboethoxypiperidin-1-yl)carbonyl)ethenyl) phenyl] sulfide;

(Benzodioxan-6-yl)[2-trifluoromethyl-4-(*E*-((2-carboethoxypiperidin-1-yl) carbonyl)ethenyl) phenyl] sulfide;

(Benzodioxan-6-yl)[2-nitro-4-(*E*-((4-carboxypiperidin-1-yl) carbonyl)ethenyl) phenyl] sulfide;

- 5 (Benzodioxan-6-yl)[2-trifluoromethyl-4-(*E*-((3-carboxypyrrolidin-1-yl)carbonyl)ethenyl) phenyl] sulfide;

(Benzodioxan-6-yl)[2-trifluoromethyl-4-(*E*-((4-carboethoxypiperidin-1-yl) carbonyl)ethenyl) phenyl] sulfide;

- 10 (Benzodioxan-6-yl)[2-trifluoromethyl-4-(*E*-((2-carbomethoxy-4-*tert*-butoxycarbonylpiperazin-1-yl) carbonyl)ethenyl) phenyl] sulfide;

(Benzodioxan-6-yl)[2-trifluoromethyl-4-(*E*-((2-carbomethoxy-4-methoxycarbonylpiperazin-1-yl) carbonyl)ethenyl) phenyl] sulfide;

(Benzodioxan-6-yl)[2-trifluoromethyl-4-(*E*-((2-carbomethoxypiperazin-1-yl) carbonyl)ethenyl) phenyl] sulfide;

- 15 (2-Methyl-3-(carboethoxymethyl)indol-5-yl)[2-trifluoromethyl-4-(*E*-((morpholin-1-yl)carbonyl)ethenyl) phenyl] sulfide;

(1-(2-Methoxyethyl)indol-5-yl)[2-chloro-4-(*E*-((4-acetylpiperazin-1-yl)carbonyl)ethenyl) phenyl] sulfide;

(2-Isopropylphenyl)[2-nitro-4-(*E*-((3-acetoxymethyl-4-hydroxypiperidin-1-

- 20 yl)carbonyl)ethenyl) phenyl] sulfide;

(2-Isopropylphenyl)[2-nitro-4-(*E*-((3-(dimethylaminocarbonyl)-4-hydroxypiperidin-1-yl)carbonyl)ethenyl) phenyl] sulfide;

(2-Isopropylphenyl)[2-nitro-4-(*E*-((3-cyanomorpholin-1-yl)carbonyl)ethenyl) phenyl] sulfide;

(2-Isopropylphenyl)[2-nitro-4-(*E*-((3-carboethoxymorpholin-1-yl)carbonyl)ethenyl) phenyl] sulfide;

- 5 (2-Isopropylphenyl)[2-nitro-4-(*E*-((tetrazol-5-yl)morpholin-1-yl)carbonyl)ethenyl) phenyl] sulfide;

(Benzodioxan-6-yl)[2-trifluoromethyl-4-(*E*-((4-carboxypiperidin-1-yl)carbonyl)ethenyl) phenyl] sulfide;

- (Benzodioxan-6-yl)[2-trifluoromethyl-4-(*E*-((2-carboxypiperidin-1-yl)carbonyl)ethenyl) phenyl] sulfide;

- 10 (Benzodioxan-6-yl)[2-trifluoromethyl-4-(*E*-((4-carbomethoxypiperazin-1-yl)carbonyl)ethenyl) phenyl] sulfide;

(Benzodioxan-6-yl)[2-trifluoromethyl-4-(*E*-((3-aza-6,9-dioxaspiro[5.4]decan-1-yl)carbonyl)ethenyl) phenyl] sulfide;

- 15 (Benzodioxan-6-yl)[2-trifluoro-4-(*E*-((4-(benzimidazol-1-yl)piperidin-1-yl)carbonyl)ethenyl) phenyl] sulfide;

(Benzodioxan-6-yl)[2-trifluoromethyl-4-(*E*-((4-(methylaminocarbonyl)piperidin-1-yl)carbonyl)ethenyl) phenyl] sulfide;

- (Benzodioxan-6-yl)[2-trifluoromethyl-4-(*E*-((3-carbomethoxy-4-methoxycarbonyl)piperazin-1-yl) carbonyl)ethenyl) phenyl] sulfide;

(2-Isopropylphenyl)[2-nitro-4-(*E*-((3-carboxymorpholin-1-yl)carbonyl)ethenyl) phenyl] sulfide;

- (Benzodioxan-6-yl)[2-trifluoromethyl-4-(*E*-((2-carboxy-4-methoxycarbonylpiperazin-1-yl) carbonyl)ethenyl) phenyl] sulfide;
- (Benzodioxan-6-yl)[2-trifluoromethyl-4-(*E*-((morpholin-1-yl)carbonyl)ethenyl) phenyl] sulfide;
- 5 (Benzodioxan-6-yl)[2-trifluoromethyl-4-(*E*-((4-(pyrrolidin-1-yl)piperidin-1-yl)carbonyl)ethenyl) phenyl] sulfide;
- (2-Isopropylphenyl)[2-nitro-4-(*E*-((3-aza-6,9-dioxaspiro[5.4]decan-1-yl)carbonyl)ethenyl) phenyl] sulfide;
- (2-Isopropylphenyl)[2-nitro-4-(*E*-((2-(dimethylaminomethyl)piperidin-1-yl)carbonyl)ethenyl) phenyl] sulfide;
- 10 (2-Isopropylphenyl)[2-nitro-4-(*E*-((piperidin-1-ylamino)carbonyl)ethenyl) phenyl] sulfide;
- (Benzodioxan-6-yl)[2-trifluoromethyl-4-(*E*-((3-carboxy-4-methoxycarbonylpiperazin-1-yl) carbonyl)ethenyl) phenyl] sulfide;
- 15 (2-(Dimethylaminocarbonyl)-benzodioxan-6-yl)[2-chloro-4-(*E*-((4-acetyl)piperazin-1-yl)carbonyl)ethenyl) phenyl] sulfide;;
- (2-Isopropylphenyl)[2-nitro-4-(*E*-((3-(2-(methoxymethyl)tetrazol-5-yl) piperidin-1-yl)carbonyl)ethenyl) phenyl] sulfide
- (2-Isopropylphenyl)[2-nitro-4-(*E*-((3-(1-(methoxymethyl)tetrazol-5-yl) piperidin-1-yl)carbonyl)ethenyl) phenyl] sulfide;
- 20 (1-Methylindol-5-yl)[2-chloro-4-(*E*-((3-(1-pyrrolidin-2-onyl)propylamino) carbonyl)ethenyl) phenyl] sulfide;

- (2-Isopropylphenyl)[2-nitro-4-(*E*-((3-(tetrazol-5-yl) piperidin-1-yl)carbonyl)ethenyl) phenyl] sulfide;
- (1-Methylindol-5-yl)[2-chloro-4-(*E*-((3-carboethoxypiperidin-1-yl)carbonyl)ethenyl) phenyl] sulfide
- 5 (1-Methylindol-5-yl)[2-chloro-4-(*E*-((3-carboxypiperidin-1-yl)carbonyl)ethenyl) phenyl] sulfide;
- (1-Methylindol-5-yl)[2-chloro-4-(*E*-((4-carboethoxypiperidin-1-yl)carbonyl)ethenyl) phenyl] sulfide;
- (1-Methylindol-5-yl)[2-chloro-4-(*E*-((3-carboxypiperidin-1-yl)carbonyl)ethenyl) phenyl] sulfide;
- 10 (2-Isopropylphenyl)[2-nitro-4-(*E*-((2-(1-methylpyrrolidin-2-yl)ethylamino)carbonyl)ethenyl) phenyl] sulfide;
- (2-Isopropylphenyl)[2-nitro-4-(*E*-((4-(pyrrolidin-1-yl)piperidin-1-yl)carbonyl)ethenyl) phenyl] sulfide;
- 15 (2-Isopropylphenyl)[2-nitro-4-(*E*-((4-sulfopiperidin-1-yl)carbonyl)ethenyl) phenyl] sulfide;
- (2-Isopropylphenyl)[2-nitro-4-(*E*-((3-hydroxypiperidin-1-yl)carbonyl)ethenyl) phenyl] sulfide;
- (Benzodioxan-6-yl)[2-trifluoromethyl-4-(*E*-((3-((ethanesulfonylamino)carbonyl)piperidin-1-yl) carbonyl)ethenyl) phenyl] sulfide;
- 20 (Benzodioxan-6-yl)[2-trifluoromethyl-4-(*E*-((3-(*p*-toluenesulfonylamino)carbonyl)piperidin-1-yl) carbonyl)ethenyl) phenyl] sulfide;

09544795-033400

- (Benzodioxan-6-yl)[2-trifluoromethyl-4-(*E*-((4-((ethanesulfonylamino)carbonyl)piperidin-1-yl) carbonyl)ethenyl) phenyl] sulfide;
- (Benzodioxan-6-yl)[2-trifluoromethyl-4-(*E*-((2-tetrazol-5-yl)morpholin-1-yl)carbonyl)ethenyl) phenyl] sulfide;
- 5 (2-Isopropylphenyl)[2-nitro-4-(*E*-((2-butyl, 5-(tetrazol-5-yl)morpholin-1-yl)carbonyl)ethenyl) phenyl] sulfide;
- (2-(and 3-)(Hydroxymethyl)-benzodioxan-6-yl)[2-nitro-4-(*E*-((4-acetylpiperazin-1-yl)carbonyl)ethenyl) phenyl] sulfide;
- (2-(and 3-)(Hydroxymethyl)-benzodioxan-6-yl)[2-nitro-4-(*E*-((3-(pyrrolidin-2-on-1-yl)prop-1-ylamino)carbonyl) ethenyl)phenyl]sulfide;
- 10 (2-(and 3-)(Hydroxymethyl)-benzodioxan-6-yl)[2-trifluoromethyl-4-(*E*-((3-(pyrrolidin-2-on-1-yl)prop-1-ylamino)carbonyl) ethenyl)phenyl]sulfide;
- (3-Hydroxymethyl)-benzodioxan-6-yl)[2-nitro-4-(*E*-((3-(pyrrolidin-2-on-1-yl)prop-1-ylamino)carbonyl) ethenyl)phenyl]sulfide;
- 15 (Benzodioxan-6-yl)[2-chloro-4-(*E*-((3-carboxypiperidin-1-yl)carbonyl)ethenyl) phenyl]
- (2-(and 3-)(Aminomethyl)-benzodioxan-6-yl)[2-trifluoromethyl-4-(*E*-((3-(pyrrolidin-2-on-1-yl)prop-1-ylamino)carbonyl) ethenyl)phenyl]sulfide;
- (2-Isopropylphenyl)[2-nitro-4-(*E*-((3-(methylaminocarbonyl)morpholin-1-yl)carbonyl)ethenyl) phenyl] sulfide;
- 20 (2-Isopropylphenyl)[2-nitro-4-(*E*-((3-(hydroxymethyl)morpholin-1-yl)carbonyl)ethenyl) phenyl] sulfide;

001100-96714560

(2-Isopropylphenyl)[2-nitro-4-(*E*-((3-(acetoxymethyl)morpholin-1-yl)carbonyl)ethenyl) phenyl] sulfide;

(2-Isopropylphenyl)[2-nitro-4-(*E*-((3-(aminomethyl)morpholin-1-yl)carbonyl)ethenyl) phenyl] sulfide;

- 5 (2-Isopropylphenyl)[2-nitro-4-(*E*-((3-(acetamidomethyl)morpholin-1-yl)carbonyl)ethenyl) phenyl] sulfide;

(Benzodioxan-6-yl)[2-chloro-4-(*E*-((3-(pyrrolidin-2-on-1-yl)prop-1-ylamino)carbonyl) ethenyl)phenyl]sulfide;

- 10 (Benzodioxan-6-yl)[2-chloro-4-(*E*-((3-carboethoxypiperidin-1-yl) carbonyl)ethenyl) phenyl] sulfide;

(Benzodioxan-6-yl)[2-chloro-4-(*E*-((2-carboethoxypiperidin-1-yl) carbonyl)ethenyl) phenyl] sulfide

(2-Methoxyphenyl)-[2,3-dichloro-4(*E*-[(morpholin-1-yl)carbonyl]ethenyl)phenyl] sulfide;

- 15 (2-Methoxyphenyl)-[2,3-dimethyl-4(*E*-[(morpholin-1-yl)carbonyl]ethenyl)phenyl] sulfide;

(2-Isopropylphenyl)[2-nitro-4-(*E*-((indol-5-ylamino)carbonyl)ethenyl) phenyl] sulfide;

(Benzodioxan-6-yl)[2-chloro-4-(*E*-((3-carboxypiperidin-1-yl) carbonyl)ethenyl)

- 20 phenyl] sulfide;

(Benzodioxan-6-yl)[2-chloro-4-(*E*-((3-(tetrazol-5-yl)piperidin-1-yl) carbonyl)ethenyl) phenyl] sulfide;

(Benzodioxan-6-yl)[2-chloro-4-(*E*-((4-*tert*-butoxycarbonyl)piperazin-1-yl) carbonyl)ethenyl] phenyl] sulfide;

(Benzodioxan-6-yl)[2-chloro-4-(*E*-((2-carboxypiperidin-1-yl) carbonyl)ethenyl] phenyl] sulfide;

- 5 (Benzodioxan-6-yl)[2-chloro-4-(*E*-((3-(tetrazol-5-yl)morpholin-1-yl) carbonyl)ethenyl] phenyl] sulfide;

(Benzodioxan-6-yl)[2-chloro-4-(*E*-((4-(methylaminocarbonyl)piperazin-1-yl) carbonyl)ethenyl] phenyl] sulfide;

(2-Methoxyphenyl)-[2,3-dichloro-4(*E*-[(4-carboxypiperidin-1-yl)carbonyl]ethenyl)

- 10 phenyl] sulfide;

(Benzodioxan-6-yl)[2-chloro-4-(*E*-((4-(tetrazol-5-yl)piperidin-1-yl) carbonyl)ethenyl] phenyl] sulfide;

(2-Methoxyphenyl)-[3-chloro-4(*E*-[(morpholin-1-yl)carbonyl]ethenyl)phenyl]sulfide;

(2-Isopropylphenyl)[2-nitro-4-(*E*-((4-oxopiperidin-1-yl)carbonyl)ethenyl) phenyl]

- 15 sulfide;

(Benzodioxan-6-yl)[2-trifluoromethyl-4-(*E*-((3-*R*-carboethoxypiperidin-1-yl)carbonyl)ethenyl] phenyl] sulfide;

(Benzodioxan-6-yl)[2-trifluoromethyl-4-(*E*-((3-*R*-carboxypiperidin-1-yl)carbonyl)ethenyl] phenyl] sulfide;

- 20 (Benzodioxan-6-yl)[2,3-dichloro-4-(*E*-((3-(pyrrolidin-2-on-1-yl)prop-1-ylamino)carbonyl) ethenyl)phenyl]sulfide;

(Benzodioxan-6-yl)[2,3-dichloro-4-(*E*-((4-acetylpiperazin-1-yl) carbonyl)ethenyl) phenyl] sulfide;

(Benzodioxan-6-yl)[2,3-dichloro-4-(*E*-((3-carboethoxypiperidin-1-yl) carbonyl)ethenyl) phenyl] sulfide;

- 5 (Benzodioxan-6-yl)[2,3-dichloro-4-(*E*-((4-carboethoxypiperidin-1-yl) carbonyl)ethenyl) phenyl] sulfide;

(Benzodioxan-6-yl)[2,3-dichloro-4-(*E*-((3-carboxypiperidin-1-yl) carbonyl)ethenyl) phenyl] sulfide;

- 10 (Benzodioxan-6-yl)[2,3-dichloro-4-(*E*-((4-carboxypiperidin-1-yl) carbonyl)ethenyl) phenyl] sulfide;

(2-Isopropylphenyl)[2,3-dichloro-4-(*E*-((3-(1-pyrrolidin-2-onyl)propylamino) carbonyl)ethenyl) phenyl] sulfide;

(2-Isopropylphenyl)[2,3-dichloro-4-(*E*-((4-acetylpiperazin-1-yl) carbonyl)ethenyl) phenyl] sulfide;

- 15 (2-Isopropylphenyl)[2,3-dichloro-4-(*E*-((3-carboethoxypiperidin-1-yl) carbonyl)ethenyl) phenyl] sulfide;

(2-Isopropylphenyl)[2,3-dichloro-4-(*E*-((4-carboethoxypiperidin-1-yl) carbonyl)ethenyl) phenyl] sulfide;

(2-Isopropylphenyl)[2,3-dichloro-4-(*E*-((3-carboxypiperidin-1-yl) carbonyl)ethenyl)

- 20 phenyl] sulfide;

(2-Isopropylphenyl)[2,3-dichloro-4-(*E*-((4-carboxypiperidin-1-yl) carbonyl)ethenyl) phenyl] sulfide;

(1-Methylindol-5-yl)[2,3-dichloro-4-(*E*-((3-carboethoxypiperidin-1-yl) carbonyl)ethenyl) phenyl] sulfide;

(1-Methylindol-5-yl)[2,3-dichloro-4-(*E*-((3-carboxypiperidin-1-yl) carbonyl)ethenyl) phenyl] sulfide;

5 (1-Methylindol-5-yl)[2,3-dichloro-4-(*E*-((4-carboethoxypiperidin-1-yl) carbonyl)ethenyl) phenyl] sulfide;

(1-Methylindol-5-yl)[2,3-dichloro-4-(*E*-((4-carboxypiperidin-1-yl) carbonyl)ethenyl) phenyl] sulfide;

(2-Ethoxyphenyl)-[2,3-dichloro-4(*E*-[(4-carboxypiperidin-1-yl)carbonyl]ethenyl) phenyl] sulfide;

10 (2-Ethoxyphenyl)-[2,3-dichloro-4(*E*-[(morpholin-1-yl)carbonyl]ethenyl)phenyl] sulfide;

(2-Ethoxyphenyl)-[2,3-dichloro-4(*E*-[(3-carboxypiperidin-1-yl)carbonyl]ethenyl) phenyl] sulfide;

15 (2-Isopropylphenyl)[2-nitro-4-(*E*-((3-carboethoxypyrrolidin-1-yl)carbonyl)ethenyl) phenyl] sulfide;

(2-Isopropylphenyl)[2-nitro-4-(*E*-((3-carboxypyrrolidin-1-yl)carbonyl)ethenyl) phenyl] sulfide;

(2-Isopropylphenyl)[2,3-difluoro-4-(*E*-((3-carboethoxypiperidin-1-yl)carbonyl)ethenyl) phenyl] sulfide;

20 (2-Isopropylphenyl)[2,3-difluoro-4-(*E*-((3-carboxypiperidin-1-yl)carbonyl)ethenyl) phenyl] sulfide;

(2-Isopropylphenyl)[2,3-difluoro-4-(*E*-((4-carboxypiperidin-1-yl)carbonyl)ethenyl)phenyl] sulfide;

(Benzodioxan-6-yl)[2-trifluoromethyl-4-(*E*-((3-ethoxycarbonylpyrrolidin-1-yl)carbonyl)ethenyl) phenyl] sulfide

- 5 (Benzodioxan-6-yl)[2-trifluoromethyl-4-(*E*-((3-carboxypyrrolidin-1-yl)carbonyl)ethenyl) phenyl] sulfide;

(2-Methoxyphenyl)[2-chloro-3-trifluoromethyl-4-(*E*-((4-carboethoxypiperidin-1-yl)carbonyl)ethenyl) phenyl] sulfide;

- (2-Methoxyphenyl)[2-chloro-3-trifluoromethyl-4-(*E*-((4-carboethoxypiperidin-1-yl)carbonyl)ethenyl) phenyl] sulfide;

(2-Methoxyphenyl)[2-chloro-3-trifluoromethyl-4-(*E*-((morpholin-1-yl)carbonyl)ethenyl) phenyl] sulfide;

(Benzodioxan-6-yl) [4-(*E*-((4-carboxypiperidin-1-yl) carbonyl)ethenyl)naphthyl] sulfide;

- 15 (2-Methoxyphenyl) [2,3-dichloro-4-(*E*-((4-(spiro-hydantoin-5-yl)-piperidin-1-yl)carbonyl)ethenyl)phenyl] sulfide;

(2-Methoxyphenyl) [2,3-dichloro-4-(*E*-(4-(2-(2-hydroxyethoxy)ethyl)piperazin-1-yl)carbonyl)ethenyl)phenyl] sulfide;

(2-Methoxyphenyl)[2,3-dichloro-4-(*E*-((4-ethylpiperazin-1-

- 20 yl)carbonyl)ethenyl)phenyl] sulfide;

(2-Isopropylphenyl)[2,3-dichloro-4-(*E*-((4-(2-(2-hydroxyethoxy)ethyl)piperazin-1-yl)carbonyl)ethenyl)phenyl] sulfide;

- (Benzodioxan-6-yl)[2,3-bis(trifluoromethyl)-4-(*E*-((4-carboxypiperidin-1-yl)carbonyl)ethenyl)phenyl]sulfide;
- (2-Methoxyphenyl) [2,3-dichloro-4-(*E*-((4-(carboxymethylamino)carbonyl)-piperidin-1-yl)carbonyl)ethenyl)phenyl] sulfide;
- 5 (2-Methoxyphenyl) [2,3-bis(trifluoromethyl)-4-(*E*-((4-carboxymethylpiperazin-1-yl)carbonyl)ethenyl)phenyl]sulfide;
- (2-Methoxyphenyl) [2,3-bis(trifluoromethyl)-4-(*E*-((4-*N*-(2-hydroxyethyl)piperazin-1-yl)carbonyl)ethenyl)phenyl]sulfide;
- (1-Methylindol-5-yl) [2,3-dichloro-4-(*E*-((4-(carbo-2,3-
- 10 dihydroxypropylamino)piperidin-1-yl)carbonyl)ethenyl)phenyl] sulfide;
- (2-Methoxyphenyl) [2,3-dichloro-4-(*E*-4-(2,3-dihydroxypropionyl)piperazin-1-yl)carbonyl)ethenyl)phenyl] sulfide;
- (2-Methoxyphenyl) [2,3-dichloro-4-(*E*-4-(2,3-dihydroxy-3-carboxypropionyl)piperazin-1-yl)carbonyl)ethenyl)phenyl] sulfide;
- 15 (1-Methylindol-5-yl) [2,3-dichloro-4-(*E*-((4-(carboxymethylamino)carbonyl)-piperidin-1-yl)carbonyl)ethenyl)phenyl] sulfide;
- (1-Methylindol-5-yl) [2,3-dichloro-4-(*E*-((4-sulfopiperidin-1-yl)carbonyl)ethenyl)phenyl] sulfide;
- (1-Methylindol-5-yl) [2,3-dichloro-4-(*E*-4-methylhomopiperazin-1-
- 20 ylcarbonyl)ethenyl)phenyl] sulfide;
- (1-Methylindol-5-yl) [2,3-dichloro-4-(*E*-4-tetrahydrofuroylpiperazin-1-yl)carbonyl)ethenyl)phenyl] sulfide;

- (2-Methoxyphenyl) [2,3-dichloro-4-(*E*-((4-amino-4-carboxypiperidin-1-yl)carbonyl)ethenyl)phenyl] sulfide;
- (2-Methoxyphenyl)[2,3-dichloro-4-((4-furoylpiperazin-1-yl)carbonyl)ethenyl)phenyl] sulfide;
- 5 (1-Methylindol-5-yl) [2,3-dichloro-4-(*E*-(4-(carbo-3-sulfopropylamino)piperadin-1-yl)carbonyl)ethenyl)phenyl] sulfide;
- (2-Methoxyphenyl)[2,3-dichloro-4-(*E*-(4-acetylamino-4-carboxypiperidin-1-yl)carbonyl)ethenyl)phenyl] sulfide;
- (2-Methoxyphenyl) [2,3-bis(trifluoromethyl)-4-(*E*-((4-carboxypiperidin-1-yl)carbonyl)ethenyl)phenyl]sulfide;
- 10 (2-Methoxyphenyl) 5-[8-(*E*-((4-(aminocarbonyl)piperidin-1-yl)carbonyl)ethenyl)quinoliny]sulfide;
- (2-Methoxyphenyl) [2-trifluoromethyl-4-(*E*-((4-carboxypiperidin-1-yl)carbonyl)ethenyl)phenyl]sulfide;
- 15 (1-Methylindol-5-yl) [2,3-dichloro-4-(*E/Z*-((1*S*,4*S*)-2,5-diazabicyclo(2,2,1)heptan-2-ylcarbonyl)ethenyl)-2,3-dichlorophenyl] sulfide;
- (1-Methylindol-5-yl) [2,3-dichloro-4-(*E*-(4-hydroxy-3-carboxypiperadin-1-yl)carbonyl)ethenyl)phenyl] sulfide;;
- (1-Methylindol-5-yl) [2,3-dichloro-4-(*E*-(*S*-oxothiomorpholin-1-ylcarbonyl)ethenyl)phenyl] sulfide;
- 20 (2-Methoxyphenyl) [2,3-dichloro-4-(*E*-((4-sulfophenylamino)carbonyl)ethenyl)phenyl] sulfide;

- (2-Methoxyphenyl) [2,3-dichloro-4-(*E*-((4-carboxyphenylamino)carbonyl)ethenyl)phenyl] sulfide;
- [3-(4-Morpholino)phenyl] [2,3-dichloro-4-(*E*-[(4-carboxypiperidin-1-yl)carbonyl]ethenyl)phenyl] sulfide;
- 5 [3-(4-Morpholino)phenyl] [2,3-dichloro-4-(*E*-((4-carboxypiperidin-1-yl)carbonyl)ethenyl)phenyl] sulfide;
- (2-Methoxyphenyl)[2,3-bis(trifluoromethyl)-4-(*E*-((4-phenylcarboxypiperidin-1-yl)carbonyl)ethenyl)phenyl]sulfide;
- (2-Methoxyphenyl) [2,3-dichloro-4-(*E*-(((4-hydroxylaminocarbonyl)piperidin-1-yl)carbonyl)ethenyl)phenyl] sulfide;
- 10 (2-Methoxyphenyl) [2,3-dichloro-4-(*E*-((N-carboxymethyl-N-phenylamino)carbonyl)ethenyl)phenyl] sulfide;

0541735.013100
DATE 05-27-99

- (2-Methoxyphenyl) [3-chloro-6-hydroxy-4-(E-((3-carboxypiperidin-1-yl)carbonyl)ethenyl)phenyl]sulfide;
- (2-Methoxyphenyl) [2,3-dichloro-4-(E-(4-((1-(2-phenyl-1-carboxyethyl)amino)carbonyl)piperidin-1-yl)carbonyl)ethenyl)phenyl]sulfide;
- 5 (2-Methoxyphenyl) [2,3-dichloro-4-(E-(4-((1-(2-hydroxy-1-carboxyethyl)amino)carbonyl)piperidin-1-yl)carbonyl)ethenyl)phenyl]sulfide;
- (3-(1-(3-Carboxypiperidinyl)phenyl)[2,3-dichloro-4-(E-((1,2,5,6-tetrahydropyridin-1-yl)carbonyl)ethenyl)phenyl]sulfide;
- (3-(4-Pyrrolidin-1-yl)piperidin-1-yl)phenyl [2,3-dichloro-4-(E-(((3-(2-pyrrolidinon-10 1-yl)propylamino)carbonyl)ethenyl)phenyl]sulfide;
- [3-(4-(Spiro-2,2-dioxolanyl)piperidin-1-yl)phenyl] [2,3-dichloro-4-(E-((4-morpholinyl)carbonyl)ethenyl)phenyl]sulfide;
- [3-(3-Carboxypiperidin-1-yl)phenyl] [2,3-dichloro-4-(E-[(4-carboxypiperidin-1-yl)carbonyl]ethenyl)phenyl] sulfide;
- 15 (2-(2-Carboxy)ethenyl)phenyl [2,3-dichloro-4-(E-((4-morpholinyl)carbonyl)ethenyl)phenyl]sulfide;
- [3-(4-Carboxylpiperidin-1-yl)phenyl] [2,3-dichloro-4-(E-[(1,2,3,6-tetrahydropyridine)-1-yl)carbonyl]ethenyl)phenyl] sulfide;
- [3-(4-Carboxylpiperidinyl)phenyl] [2,3-dichloro-4-(E-[(4-morpholinyl)carbonyl]ethenyl)phenyl] sulfide;
- 20 [2-(4-Acetyl)piperazin-1-yl)phenyl] [2,3-dichloro-4-(E-[(4-carboxypiperidin-1-yl)carbonyl]ethenyl)phenyl] sulfide;

0954795-033100

- 3-(3-Carboxypiperidin-1-yl)phenyl] [2,3-dichloro-4-(*E*-[(4-morpholinyl)carbonyl]ethenyl)phenyl] sulfide;
- [3-(4-Carboxypiperidin-1-yl)phenyl] [2,3-dichloro-4-(*E*-[(4-(dimethylaminosulfamoyl)piperazin-1-yl)carbonyl]ethenyl)phenyl] sulfide;
- 5 (2-Methoxyphenyl)[2,3-bis(trifluoromethyl)-4-(*E*-((3-carboxypiperidin-1-yl)carbonyl)ethenyl)phenyl]sulfide;
- (2-Methoxyphenyl) [2,3-bis(trifluoromethyl)-4-(*E*-((2-carboxypyrrolidin-1-yl)carbonyl)ethenyl)phenyl]sulfide;
- [3-(4-Carboxypiperidin-1-yl)phenyl] [2,3-dichloro-4-(*E*-((4-((trifluoromethylsulfonyl)piperazin-1-yl)carbonyl)ethenyl)phenyl] sulfide;
- 10 (2-Methoxyphenyl)[2,3-dichloro-4-(*E*-((piperidin-1-ylcarbonyl)ethenyl)phenyl] sulfide;
- (2-Hydroxyphenyl) [2,3-dichloro-4-(*E*-((4-morpholino)carbonyl)ethenyl)phenyl]sulfide;
- 15 (2-Methoxyphenyl) [2,3-dichloro-4-(*E*-(((4-carboxyphenyl)methyl)amino)carbonyl)ethenyl)phenyl]sulfide;
- (2-Methoxyphenyl) [2,3-dichloro-4-(*E*-(((4-pyrrolidin-1-yl)piperidin-1-yl)carbonyl)ethenyl)phenyl]sulfide;
- (2-Hydroxyphenyl)[2,3-dichloro-4-(*E*-((4-carboxypiperidin-1-yl)carbonyl)ethenyl)phenyl]sulfide;
- 20 [3-(4-Carboxypiperidin-1-yl)phenyl] [2,3-dichloro-4-(*E*-((4-(methylsulfonyl)piperazin-1-yl)carbonyl)ethenyl)phenyl] sulfide;

- (2-Aminophenyl) [2,3-dichloro-4-(*E*-((4-morpholinyl)carbonyl)ethenyl)phenyl] sulfide;
- (3-(4-carboxypiperidin-1-yl)phenyl)[2,3-dichloro-4-(*E*-((S-oxothiomorpholin-1-yl)carbonyl)ethenyl)phenyl] sulfide;
- 5 [3-(4-Carboxypiperidin-1-yl)phenyl] [2,3-dichloro-4-(*E*-((4-hydroxypiperidin-1-yl)carbonyl)ethenyl)phenyl] sulfide;
- (2-Glycoxyphenyl) [2,3-dichloro-4-(*E*-((4-morpholino)carbonyl)ethenyl)phenyl] sulfide;
- (2-(4-Butyroxyl)phenyl)[2,3-dichloro-4-(*E*-((4-morpholino)carbonyl)ethenyl)phenyl] sulfide;
- 10 [3-(4-Carboxypiperidin-1-yl)phenyl] [2,3-dichloro-4-(*E*-((4-hydroxyethylpiperazin-1-yl)carbonyl)ethenyl)phenyl] sulfide;
- [3-(4-Carboxypiperidin-1-yl)phenyl] [2,3-dichloro-4-(*E*-((4-furoyl)piperazin-1-yl)carbonyl)ethenyl)phenyl] sulfide;
- 15 [3-(4-Carboxypiperidin-1-yl)phenyl] [2,3-dichloro-4-(*E*-((pyrrolidin-1-yl)carbonyl)ethenyl)phenyl] sulfide;
- [3-(4-Carboxypiperidin-1-yl)phenyl] [2,3-dichloro-4-(*E*-((diethylaminocarbonyl)ethenyl)phenyl] sulfide;
- [3-(4-Carboxypiperidin-1-yl)phenyl] [2,3-dichloro-4-(*E*-((4-ethylpiperazin-1-yl)carbonyl)ethenyl)phenyl] sulfide;
- 20 [3-(4-Carboxypiperidin-1-yl)phenyl] [2,3-dichloro-4-(*E*-((4-(aminocarbonyl)piperidin-1-yl)carbonyl)ethenyl)phenyl] sulfide;

09544795.03160

- [3-(4-Carboxypiperidin-1-yl)phenyl] [2,3-dichloro-4-(*E*-((4-(2-ethoxyethyl)piperidin-1-yl)carbonyl)ethenyl)phenyl] sulfide;
- [3-((4-Carboxymethyl)piperazin-1-yl)phenyl] [(2,3-dichloro-4-(*E*-(4-morpholinyl)carbonyl)ethenyl)phenyl] sulfide;
- 5 [3-(4-Carboxypiperidin-1-yl)phenyl] [2,3-bis(trifluoromethyl)-4-(*E*-((4-morpholino)carbonyl)ethenyl)phenyl]sulfide;
- (3-Hydroxyphenyl) [2,3-dichloro-4-(*E*-((4-morpholino)carbonyl)ethenyl)phenyl]sulfide;
- [3-(4-Butyroxyl)phenyl] [2,3-dichloro-4-(*E*-((4-morpholino)carbonyl)ethenyl)phenyl]sulfide;
- 10 (2-Hydroxyphenyl) [2,3-bis(trifluoromethyl)-4-(*E*-((4-morpholino)carbonyl)ethenyl)phenyl]sulfide;
- (3-Hydroxyphenyl)[2,3-bis(trifluoromethyl)-4-(*E*-((4-morpholino)carbonyl)ethenyl)phenyl]sulfide;
- 15 [3-(4-Carboxypiperidin-1-yl)phenyl] [2,3-ditrifluoromethyl-4-(*E*-((4-hydroxypiperidin-1-yl)carbonyl)ethenyl)phenyl] sulfide;
- [3-(4-Carboxypiperidin-1-yl)phenyl] [2,3-ditrifluoromethyl-4-(*E*-((1,2,5,6-tetrahydropyridin-1-yl)carbonyl)ethenyl)phenyl] sulfide;

- [2-((4-Carboxy)butyloxy)phenyl] [2,3-dichloro-4-(E-((4-morpholino)carbonyl)ethenyl)phenyl]sulfide;
- (2-Glycoxyphenyl) [2,3-bis(trifluoromethyl)-4-(E-((4-morpholino)carbonyl)ethenyl)phenyl]sulfide;
- 5 (2-(4-Butyroxyl)phenyl)[2,3-bis(trifluoromethyl)-4-(E-((4-morpholino)carbonyl)ethenyl)phenyl]sulfide;
- [3-(4-Carboxypiperidin-1-yl)phenyl] [2,3-ditrifluoromethyl-4-(E-((bis-(2-ethoxyethyl)amino)carbonyl)ethenyl)phenyl] sulfide;
- [3-(4-Carboxypiperidin-1-yl)phenyl] [2,3-bis-(trifluoromethyl)-4-(E-((bis-(2-hydroxypropyl)amino)carbonyl)ethenyl)phenyl] sulfide;
- 10 [3-(4-Carboxypiperidin-1-yl)phenyl] [2,3-bis-(trifluoromethyl)-4-(E-((piperazin-1-yl)carbonyl)ethenyl)phenyl] sulfide;
- (3-(4-Butyroxyl)phenyl)[2,3-bis(trifluoromethyl)-4-(E-((4-morpholino)carbonyl)ethenyl)phenyl]sulfide;
- 15 [2-(3-Carboxypiperidin-1-yl)phenyl] [2,3-dichloro-4-(E-[(3-(2-pyrrolidinon-1-yl)propylaminocarbonyl)ethenyl)phenyl] sulfide;
- [2-(3-Carboxypiperidin-1-yl)phenyl] [2,3-bis(trifluoromethyl)-4-(E-[(3-(2-pyrrolidinon-1-yl)propylaminocarbonyl)ethenyl)phenyl] sulfide;
- [2-(3-Carboxypiperidin-1-yl)phenyl] [2,3-dichloro-4-(E-((3-(2-hydroxyethyl)piperazin-1-yl)carbonyl)ethenyl)phenyl] sulfide;
- 20 [2-(3-Carboxypiperidin-1-yl)phenyl] [2,3-bis(trifluoromethyl)-4-(E-((1,2,5,6-tetrahydropyridin-1-yl)carbonyl)ethenyl)phenyl] sulfide;

00541795-033100
00460-5674560

[2-(3-Carboxypiperidin-1-yl)phenyl] [2,3-bis(trifluoromethyl)-4-(*E*-((4-(2-hydroxyethyl)piperazin-1-yl)carbonyl)ethenyl)phenyl] sulfide;

[2-(3-Carboxypiperidin-1-yl)phenyl] [2,3-bis(trifluoromethyl)-4-(*E*-((4-(2-(hydroxyethoxy)ethyl)piperazin-1-yl)carbonyl)ethenyl)phenyl] sulfide; and

- 5 (3-(3-Propioxy)phenyl) [2,3-dichloro-4-(*E*-((4-morpholino)carbonyl)ethenyl)phenyl]sulfide.

001220-56274560

Pharmaceutical Compositions and Methods of Treatment

The present invention also provides pharmaceutical compositions which comprise compounds of the present invention formulated together with one or more pharmaceutically-acceptable carriers. The pharmaceutical compositions may be specially formulated for oral administration in solid or liquid form, for parenteral injection, or for rectal administration.

The pharmaceutical compositions of this invention can be administered to humans and other animals orally, rectally, parenterally, intracisternally, intravaginally, intraperitoneally, topically (as by powders, ointments, or drops), buccally, or as an oral or nasal spray. The term "parenteral" administration as used herein refers to modes of administration which include intravenous, intramuscular, intraperitoneal, intrasternal, subcutaneous and intraarticular injection and infusion.

Pharmaceutical compositions of this invention for parenteral injection comprise pharmaceutically-acceptable sterile aqueous or nonaqueous solutions, dispersions, suspensions or emulsions as well as sterile powders for reconstitution into sterile injectable solutions or dispersions just prior to use. Examples of suitable aqueous and nonaqueous carriers, diluents, solvents or vehicles include water, ethanol, polyols (such as glycerol, propylene glycol, polyethylene glycol, and the like), and suitable mixtures thereof, vegetable oils (such as olive oil), and injectable organic esters such as ethyl oleate. Proper fluidity can be maintained, for example, by the use of coating materials such as lecithin, by the maintenance of the required particle size in the case of dispersions, and by the use of surfactants.

These compositions may also contain adjuvants such as preservative, wetting agents, emulsifying agents, and dispersing agents. Prevention of the action of microorganisms may be ensured by the inclusion of various antibacterial and antifungal agents, for example, paraben, chlorobutanol, phenol sorbic acid, and the like. It may also be desirable to include isotonic agents such as sugars, sodium chloride, and the like, Prolonged absorption of the injectable pharmaceutical form may be brought about by the inclusion of agents which delay absorption such as aluminum monostearate and gelatin.

In some cases, in order to prolong the effect of the drug, it is desirable to slow the absorption of the drug from subcutaneous or intramuscular injection. This may be accomplished by the use of a liquid suspension of crystalline or amorphous material with poor water solubility. The rate of absorption of the drug then depends upon its rate of dissolution which, in turn, may depend upon crystal size and crystalline form. Alternatively, delayed absorption of a parenterally administered drug form is accomplished by dissolving or suspending the drug in an oil vehicle.

Injectable depot forms are made by forming microencapsule matrices of the drug in biodegradable polymers such as polylactide-polyglycolide. Depending upon the ratio of drug to polymer and the nature of the particular polymer employed, the rate of drug release can be controlled. Examples of other biodegradable polymers include poly(orthoesters) and poly(anhydrides). Depot injectable formulations are also prepared by entrapping the drug in liposomes or microemulsions which are compatible with body tissues.

The injectable formulations can be sterilized, for example, by filtration through a bacterial-retaining filter, or by incorporating sterilizing agents in the form of sterile solid compositions which can be dissolved or dispersed in sterile water or other sterile injectable medium just prior to use.

- 5 Solid dosage forms for oral administration include capsules, tablets, pills, powders, and granules. In such solid dosage forms, the active compound is mixed with at least one inert, pharmaceutically-acceptable excipient or carrier such as sodium citrate or dicalcium phosphate and/or (a) fillers or extenders such as starches, lactose, sucrose, glucose, mannitol, and silicic acid, (b) binders such as, for example,
- 10 carboxymethylcellulose, alginates, gelatin, polyvinylpyrrolidone, sucrose, and acacia, (c) humectants such as glycerol, (d) disintegrating agents such as agar-agar, calcium carbonate, potato or tapioca starch, alginic acid, certain silicates, and sodium carbonate, (e) solution retarding agents such as paraffin, (f) absorption accelerators such as quaternary ammonium compounds, (g) wetting agents such as, for example,
- 15 cetyl alcohol and glycerol monostearate, (h) absorbents such as kaolin and bentonite clay, and (i) lubricants such as talc, calcium stearate, magnesium stearate, solid polyethylene glycols, sodium lauryl sulfate, and mixtures thereof. In the case of capsules, tablets and pills, the dosage form may also comprise buffering agents.

- Solid compositions of a similar type may also be employed as fillers in soft
- 20 and hard-filled gelatin capsules using such excipients as lactose or milk sugar as well as high molecular weight polyethylene glycols and the like.

09544735-033400

The solid dosage forms of tablets, dragees, capsules, pills, and granules can be prepared with coatings and shells such as enteric coatings and other coatings well known in the pharmaceutical formulating art. They may optionally contain opacifying agents and can also be of a composition that they release the active ingredient(s) only, or preferentially, in a certain part of the intestinal tract, optionally, in a delayed manner. Examples of embedding compositions which can be used include polymeric substances and waxes.

The active compounds can also be in micro-encapsulated form, if appropriate, with one or more of the above-mentioned excipients.

Liquid dosage forms for oral administration include pharmaceutically-acceptable emulsions, solutions, suspensions, syrups and elixirs. In addition to the active compounds, the liquid dosage forms may contain inert diluents commonly used in the art such as, for example, water or other solvents, solubilizing agents and emulsifiers such as ethyl alcohol, isopropyl alcohol, ethyl carbonate, ethyl acetate, benzyl alcohol, benzyl benzoate, propylene glycol, 1,3-butylene glycol, dimethyl formamide, oils (in particular, cottonseed, groundnut, corn, germ, olive, castor, and sesame oils), glycerol, tetrahydrofurfuryl alcohol, polyethylene glycols and fatty acid esters of sorbitan, and mixtures thereof.

Besides inert diluents, the oral compositions can also include adjuvants such as wetting agents, emulsifying and suspending agents, sweetening, flavoring, and perfuming agents.

Suspensions, in addition to the active compounds, may contain suspending agents as, for example, ethoxylated isostearyl alcohols, polyoxyethylene sorbitol and sorbitan esters, microcrystalline cellulose, aluminum metahydroxide, bentonite, agar-agar, and tragacanth, and mixtures thereof.

5 Compositions for rectal or vaginal administration are preferably suppositories which can be prepared by mixing the compounds of this invention with suitable non-irritating excipients or carriers such as cocoa butter, polyethylene glycol or a suppository wax which are solid at room temperature but liquid at body temperature and therefore melt in the rectum or vaginal cavity and release the active compound.

10 Compounds of the present invention can also be administered in the form of liposomes. As is known in the art, liposomes are generally derived from phospholipids or other lipid substances. Liposomes are formed by mono- or multi-lamellar hydrated liquid crystals that are dispersed in an aqueous medium. Any non-toxic, physiologically-acceptable and metabolizable lipid capable of forming
15 liposomes can be used. The present compositions in liposome form can contain, in addition to a compound of the present invention, stabilizers, preservatives, excipients, and the like. The preferred lipids are the phospholipids and the phosphatidyl choline (lecithins), both natural and synthetic.

 Methods to form liposomes are known in the art. See, for example, Prescott,
20 Ed., Methods in Cell Biology, Volume XIV, Academic Press, New York, N.Y. (1976), p. 33 et seq.

001100 56214560

The compounds of the present invention may be used in the form of pharmaceutically-acceptable salts derived from inorganic or organic acids. By "pharmaceutically-acceptable salt" is meant those salts which are, within the scope of sound medical judgment, suitable for use in contact with the tissues of humans and lower animals without undue toxicity, irritation, allergic response and the like, and are commensurate with a reasonable benefit/risk ratio. Pharmaceutically-acceptable salts are well-known in the art. For example, S. M. Berge, *et al.* Describe pharmaceutically-acceptable salts in detail in J. Pharmaceutical Sciences, 1977, 66: 1 et seq. The salts may be prepared *in situ* during the final isolation and purification of the compounds of the invention or separately by reacting a free base function with a suitable acid. Representative acid addition salts include acetate, adipate, alginate, citrate, aspartate, benzoate, benzenesulfonate, bisulfate, butyrate, camphorate, camphorsulfonate, digluconate, glycerophosphate, hemisulfate, heptanoate, hexanoate, fumarate, hydrochloride, hydrobromide, hydroiodide, 2-hydroxyethanesulfonate (isethionate), lactate, maleate, methanesulfonate, nicotinate, 2-naphthalenesulfonate, oxalate, pamoate, pectinate, persulfate, 3-phenylpropionate, picrate, pivalate, propionate, succinate, tartrate, thiocyanate, phosphate, glutamate, bicarbonate, p-toluenesulfonate and undecanoate. Also, the basic nitrogen-containing groups can be quaternized with such agents as lower alkyl halides such as methyl, ethyl, propyl, and butyl chlorides, bromides and iodides; dialkyl sulfates like dimethyl, diethyl, dibutyl and diamyl sulfates; long chain halides such as decyl, lauryl, myristyl and stearyl chlorides, bromides and iodides; arylalkyl halides like benzyl and phenethyl bromides

09544795.033400

and others. Water or oil-soluble or dispersible products are thereby obtained.

Examples of acids which may be employed to form pharmaceutically acceptable acid addition salts include such inorganic acids as hydrochloric acid, hydrobromic acid, sulphuric acid and phosphoric acid and such organic acids as oxalic acid, maleic acid, succinic acid and citric acid.

Basic addition salts can be prepared *in situ* during the final isolation and purification of compounds of this invention by reacting a carboxylic acid-containing moiety with a suitable base such as the hydroxide, carbonate or bicarbonate of a pharmaceutically acceptable metal cation or with ammonia or an organic primary, secondary or tertiary amine. Pharmaceutically-acceptable basic addition salts include cations based on alkali metals or alkaline earth metals such as lithium, sodium, potassium, calcium, magnesium and aluminum salts and the like and nontoxic quaternary ammonia and amine cations including ammonium, tetramethylammonium, tetraethylammonium, methylamine, dimethylamine, trimethylamine, triethylamine, diethylamine, ethylamine and the like. Other representative organic amines useful for the formation of base addition salts include ethylenediamine, ethanolamine, diethanolamine, piperidine, piperazine and the like.

Dosage forms for topical administration of a compound of this invention include powders, sprays, ointments and inhalants. The active compound is mixed under sterile conditions with a pharmaceutically-acceptable carrier and any needed preservatives, buffers, or propellants which may be required. Ophthalmic

09544795-03310
001300-5627150

formulations, eye ointments, powders and solutions are also contemplated as being within the scope of this invention.

Actual dosage levels of active ingredients in the pharmaceutical compositions of this invention may be varied so as to obtain an amount of the active compound(s) that is effective to achieve the desired therapeutic response for a particular patient, compositions, and mode of administration. The selected dosage level will depend upon the activity of the particular compound, the route of administration, the severity of the condition being treated, and the condition and prior medical history of the patient being treated. However, it is within the skill of the art to start doses of the compound at levels lower than required for to achieve the desired therapeutic effect and to gradually increase the dosage until the desired effect is achieved.

Generally dosage levels of about 0.1 to about 50 mg, more preferably of about 5 to about 20 mg of active compound per kilogram of body weight per day are administered orally or intravenously to a mammalian patient. If desired, the effective daily dose may be divided into multiple doses for purposes of administration, e.g. two to four separate doses per day.

00541795.033100
001220.5624560

Preparation of Compounds of the Invention

The compounds and processes of the present invention may be better understood in connection with the following synthetic Schemes which illustrate the methods by which the compounds of the invention can be prepared.

Scheme 1




10

Scheme 1 describes the synthesis of a typical cinnamide-substituted diaryl sulfide **4** through an aldehyde intermediate **2**. Aldehyde **2** is prepared by reaction of a thiophenol (for example 2,4-dichlorothiophenol, 2-bromothiophenol, or the like) with halo-substituted benzaldehyde derivative **1** (e.g. 2-chlorobenzaldehyde, 3-chloro, 4-fluorobenzaldehyde, or the like) in the presence of base (e.g. sodium carbonate, triethylamine, or the like) and a polar solvent (e.g. dimethylformamide, dimethylsulfoxide, or the like). The aldehyde group is homologated to the corresponding cinnamic acid **3**, using an acetate equivalent (for example, malonic acid, triethoxyphosphonoacetate, or the like) in the presence of an appropriate base and solvent. In some cases, it may be necessary to hydrolyze an intermediate ester (for example using sodium hydroxide in alcohol). The acid group is activated (for

20


example using thionyl chloride, or dicyclohexylcarbodiimide and N-hydroxysuccinimide, or the like) and reacted with a primary or secondary amine (for example, 6-aminohexanol, pyrrolidone-3-propylamine, or the like) to provide the desired analog **4**. In one variant, a halo-acetophenone can replace benzaldehyde **2**; the resultant cinnamides **4** are substituted with a methyl group at the 3-position.

Scheme 2




Alternatively, the order of these coupling steps may be reversed (Scheme 2). A substituted halocinnamic acid **5** (e.g. 3-chloro-2-nitrocinnamic acid or the like) may be coupled with a primary or secondary amine (e.g. N-acetyl piperazine or the like) as described above to give the corresponding amide **6**. The halo-group can then be displaced with a substituted thiophenol in the presence of base to provide the product **7**.

Scheme 3




A number of the compounds described herein may be prepared from intermediate benzylic alcohols like **8** (Scheme 3). Activation of the alcohol moiety (for example, using phosphorus tribromide or methanesulfonyl chloride and lithium halide in dimethylformamide) and displacement with a primary or secondary amine (e.g. morpholine, N-formylpiperazine or the like) provides analogs with structures related to **9**. Alternatively the alcohol may be oxidized (for example using TPAP or PCC or the like) to give aldehyde **10**.

Scheme 4




Cinnamides like **13** may be prepared from halo-substituted derivatives **11** by palladium-mediated coupling [e.g. using tetrakis (o-tolyl phosphine) palladium (0), $\text{Pd}_2(\text{dba})_3$, or the like] with acrylamide derivatives **12** (Scheme 4). In similar manner, anilino-cinnamides like **16** can be prepared by palladium-mediated coupling of amines **15** with halo-cinnamides **14**.

Scheme 5



- 5 In some cases, functional groups on the aromatic rings can be modified to produce new analogs (Scheme 5). For example, a nitro group in compounds like 17 may be reduced (for example, with tin(II) chloride, or by catalytic hydrogenation, or the like) to the corresponding amine 18. This amine may then itself be converted to a halogen, for example by diazotization using nitrous acid or t-butyl nitrite in the
- 10 presence of a metal halide salt like cupric bromide, providing analog 19.

Scheme 6




15

It is also possible to assemble cinnamide-substituted diaryl sulfides in a “reverse” sense (Scheme 6). Thus, for example, compound 20, prepared as described in Scheme 1, may be deprotected by treatment with base (e.g. potassium t-butoxide or the like) to provide thiolate anion 21, which may be reacted with an activated

haloarene (e.g. 2,3-dichlorobenzaldehyde, 3-chloro,4-fluorobenzaldehyde or the like) to provide the corresponding product **22**. Alternatively, this same thiolate anion may be coupled with unactivated aryl halides (e.g. aryl bromide or Aryl iodides) using a metal-catalyzed Ullman coupling procedure (for example, using a palladium or nickel catalyst) to give product **23**.


A further method for producing diarylsulfide cinnamides is shown in Scheme 7, wherein the diaryl sulfide is formed through coupling of a suitably protected aryl thiol **28** to an activated cinnamate ester **27**. Substituted phenol **24** may be brominated to give bromophenol **25**. Heck-type coupling of bromide **25** with an appropriate olefinic substrate, for example methyl acrylate, is effected with palladium catalysis, leading to the cinnamate ester **26**. The phenol is then activated towards further reaction, for example by conversion to the corresponding triflate **27** under standard conditions. The required protected thiol **28** may be prepared by the method of XXX (*Tetrahedron Lett.* **1994**, 35, 3221-3224), by coupling an aryl halide or triflate with triisopropylsilyl thiol under palladium catalysis. The two partners **27** and **28** are then reacted in the presence of a fluoride source, for example cesium fluoride, to provide the diarylsulfide cinnamate **29**. Hydrolysis is accomplished by basic media, such as lithium or sodium hydroxide in water-THF, and the resulting acid **30** is coupled to amines under standard amide-bond forming conditions (for example, EDC/HOBt) to produce the amides **31**.

Scheme 7



A method for preparing cinnamides bearing two arylthio groups is outlined in Scheme 8. Commercially available difluoro cinnamic acid 32 was coupled with an amine, using standard conditions, and this derived amide 33 was reacted with excess aryl thiol to provide the bis-sulfide 34.


Scheme 8



Compounds which contain trifluoromethyl groups on the cinnamide-portion of inhibitors were made by the method shown in Scheme 9. According to the method of

XXX (Ref), Diels-Alder reaction between 1,1,1,4,4,4-hexafluoro-2-butyne and 2-methylfuran led to bicyclic ether **35**, which was rearranged with Lewis acid (for example, boron trifluoride etherate) to the phenol **36**. The methyl group is then converted to the corresponding aldehyde **37** by bromination followed by reaction with dimethylsulfoxide. Using the analogous procedures described for Scheme 1 above, the phenol was activated and condensed with thiols under basic conditions to afford diarylsulfide aldehydes **38**, and further converted to cinnamides **39** by the previously described procedures.

Scheme 9




10

15


Cinnamides bearing more complex substituted piperidine amides can be produced by the methods outlined in Scheme 10 and 11. Cinnamic acids **40** are coupled to spiro-hydantoin piperidine **41**, and the derived amide **42** is first reacted with an activating reagent (for example di-tert-butyl dicarbonate), and then hydrolyzed to the amino acid **43**. The derived amino group may then be reacted further, for example with acid anhydrides or acid chlorides, to produce amides **44**.

Scheme 10



Further derivatives of piperidine amides can be obtained by coupling of piperidinone **45** with cinnamic acids **40**, as shown in Scheme 11. Standard coupling conditions lead to amide **46**, which is first reduced to the corresponding alcohol, then hydrolyzed to afford hydroxy acid **47**.


Scheme 11



Also included in this invention are compounds derived from coupling of amines, or amino acid derivatives (such as α -amino esters) to the carboxylic acid group of cinnamides **48**, using standard coupling and hydrolysis methods, as outlined in Scheme 12. Thus, amides **49** are produced directly from amine coupling reactions.

- 5 Amino acid esters are coupled to **48**, and the derived esters are hydrolyzed to the corresponding acids **50**.


Scheme 12



Inhibitors bearing substituted piperazine (or homopiperazine) cinnamides may


- 10 be produced by the methods described in Scheme 13. The methods described may be utilized to produce piperazine amide **51**. Secondary amine **51** then serves as educt for preparing amides **52**, through standard coupling reactions. Alternatively, **51** may be converted to tertiary amines **53**, through standard reductive alkylation methods (for example, condensation with an aldehyde in the presence of a reducing agent such as
- 15 sodium triacetoxyborohydride).

Scheme 13



A process for preparing analogs with amino substitutions of the aryl portion of the sulfides is illustrated in Scheme 14. The intermediate triflate **27** is reacted with halo-substituted thiophenols **54** ($X = Br, Cl, OTf, OTs$) under basic catalysis, to provide the sulfide derivative **55**. The halogen or activated hydroxyl is then substituted with an amine, using the method of Buchwald (Old, D. W.; Wolfe, J. P.; Buchwald, S. L. *J. Am. Chem. Soc.* **1998**, *120*, 9722-9723). Similar transition-metal catalyzed reactions may be applied, for example, the method of Hartwig {Hamann, B. C.; Hartwig, J. F. *J. Am. Chem. Soc.* **1998**, *120*, 7369-7370). The NR_3R_4 group may constitute a cyclic or acyclic group, optionally substituted with additional functionalities that may enhance the activities of the compounds, and that further synthetic transformations familiar to those skilled in the art may be applied. For instance, ester groups may be hydrolyzed to the corresponding carboxylic acids or amides. The derived anilino sulfides may then be processed as described above to produce the cinnamides **56**.

Scheme 14



Scheme 15 presents a synthesis of a particular class of substituted aniline derivatives bearing a carboxylic acid. A cyclic amino acid **58** may be converted into the corresponding t-butyl ester **61**, through the intermediacy of carbamate **59** and ester **60**, using standard synthetic methods. The amino ester **61** was then reacted with 2-fluoronitrobenzene with mild basic catalysis (for example, cesium fluoride, potassium bicarbonate), to provide the aniline derivative **62**. The nitro group may then be transformed into an iodo-substituted derivative **64**, by first conversion to the aniline **63**, followed by standard diazotization and reaction of the diazonium salt with potassium iodide (among other similar methods for this Sandmeyer reaction). Using the method outlined in Scheme 7, the iodide **64** may be converted to the TIPS-protected arylthiol **65**. In a sequence analogous with that described in Scheme 7, silyl thioether **65** may be reacted with cinnamide triflate **27** in the presence of a fluoride source (for example, cesium fluoride), and thus converted to the diarylsulfide **66**. Standard synthetic transformations (ester hydrolysis, amide coupling, and tert-butyl ester cleavage) provides the desired acid **68**, through intermediate ester **67**.


Compounds bearing elaborated ether groups on the arylsulfide ring were made according to Scheme 16. Methyl ether cinnamate esters such as **69** were hydrolyzed to the corresponding acids, and then the methyl ether was cleaved with boron tribromide (or alternatively using similar ether cleaving agents, such as trimethylsilyl iodide) to provide the hydroxy acids **70**. Standard coupling methods provided the amides **71**, which were then alkylated on the phenolic group using an appropriate alkyl halide **72** (where L is a linking group consisting of an acyclic or cyclic alkyl, or

heterocyclic group) or lactone ($m=1,2$) in the presence of a base (such as potassium *tert*-butoxide, sodium hydride, or cesium carbonate). Alternatively, the phenolic group was alkylated with an ester-bearing alcohol **73**, using Mitsunobu conditions. The resulting ester-bearing ethers **74** were then hydrolyzed to the corresponding acids


5 **75** using standard hydrolysis conditions. Alternatively, the ester of **74** may be *tert*-butyl, in which case acidic deprotection to acid **75** would be employed (for example, using trifluoroacetic acid in dichloromethane, or hydrochloric acid in dioxane).

001320.56274560

Scheme 15



Scheme 16



Related compounds bearing elaborated functionalized amino substituents were

- 5 made according to Scheme 17. Triflate **27** was reacted with an amino thiophenol to produce the diarylsulfide cinnamide **76** in a similar manner to that described in Schemes 1, 2 and 7. The cinnamate ester was hydrolyzed to give acid **77**, which was coupled under standard conditions to provide amide **78**. The amino group of **78** then underwent reductive alkylation with an ester-bearing alkyl aldehyde, using standard
- 10 conditions (or alternatively using sodium triacetoxyborohydride) to provide the secondary amine **79**. The ester functionality was hydrolyzed to the corresponding acid salt **80**.


An alternative strategy for producing intermediate **78** is shown in Scheme 18.

Nitro-substituted *tert*-butyl ester derivative **81** (prepared according to Scheme 14,


using the tert-butyl analog of cinnamate **27**) was cleaved to the carboxylic acid, converted to the cinnamide using standard conditions, and then the nitro group was reduced using iron powder in aqueous ammonium chloride solution.

5

Scheme 17




Scheme 18



A modified method for the preparation of analogs bearing 2,3-bis-(trifluoromethyl)cinnamides is illustrated in Scheme 19. Commercially available acrylic acid **82** was esterified with ethyl iodide, and the ester **83** was condensed with 1,1,1,4,4,4-hexafluoro-2-butyne at 110 °C to give the bicyclic adduct **84**. The bicyclic ether was then converted to the corresponding phenol **85** using a Lewis acid (for example boron trifluoride-etherate). Phenol **85** was then utilized as illustrated in Scheme 7 or Scheme 14 to prepare the desired inhibitors.

Scheme 19




Scheme 20 illustrates an alternative method for preparing substituted anilinosulfides **57**. Cinnamate ester **55** was converted to the corresponding tert-butyl ester **87**, via reaction of acid **86** with tert-butyl trichloroacetimidate under Lewis acid catalysis. The bromide **87** was then coupled with an appropriately functionalized amine (illustrated in Scheme 20 with ethyl pyrrolidinecarboxylates) using palladium

5 catalysis (for example, using the conditions of Buchwald or Hartwig noted for Scheme 14). The resultant substituted anilines **88** were then first cleaved to acids **89** using acidic conditions (TFA, HCl, or similar known deprotections for tert-butyl esters), then the acids **89** were coupled to amines HNR_3R_4 using standard conditions to provide amides **90**. The ethyl ester group of **90** was then hydrolyzed using lithium or sodium hydroxide in aqueous media to produce acids **91**.


10 Compounds with a 2,6-disubstitution pattern on the cinnamide ring system were made according to the method of Scheme 21. Commercially available 4,6-dichlorosalicylaldehyde was condensed with arylthiols under basic conditions to provide the diarylsulfide **92**. The phenolic group was protected with allyl bromide, providing the O-allyl derivative **93**. The method outlined in Scheme 1 was used to prepare the corresponding cinnamic acid **94**, then the allyl group was removed using palladium(0)-catalyzed transfer to morpholine, thus producing hydroxy cinnamic acid **95**. The acid group was coupled to a cyclic amino ester ($n=0, 1, 2$; $\text{R} = \text{Me}, \text{Et}$) under standard conditions to yield the amide **96**. Basic hydrolysis conditions reveal the acid **97**.

Scheme 20



90

Scheme 21



EXAMPLES

The compounds and processes of the present invention may be better understood in connection with the following Examples, which are intended as an illustration of and not a limitation upon the scope of the invention.

Example 1

(2,4-Dichlorophenyl)[2-(*E*-((6-hydroxyhexylamino)carbonyl)ethenyl)phenyl] sulfide

Example 1A2-[(2,4-Dichlorophenyl)thio]benzaldehyde

To a stirred solution of 2,4-dichlorothiophenol (2.0 g, 11.2 mmol) in 25 mL of anhydrous DMF was added potassium carbonate (3.09 g, 22.4 mmol), followed by 2-chlorobenzaldehyde (1.26 mL, 11.3 mmol). The mixture was then heated under nitrogen atmosphere at 70 °C for 5 hours. The reaction mixture was then allowed to cool to room temperature and partitioned between ether and water. The aqueous layer was extracted with ether once and the combined organic layer was washed with water and brine, dried over sodium sulfate and condensed in vacuo. The crude product was purified via silica gel flash chromatography, eluting with 5-10 % ether/hexanes, to give 2.62 g (9.25 mmol, 83%) of the desired aldehyde as a colorless oil, which solidified slowly upon standing at room temperature.

Example 1B*trans*-2-[(2,4-Dichlorophenyl)thio]cinnamic acid

A mixture of the aldehyde (1.50 g, 5.3 mmol) from Example 1A, malonic acid (1.21 g, 11.6 mmol), piperidine (78.6 μ L, 0.80 mmol) in 8.0 mL of anhydrous pyridine was heated at 110 °C for 2 hours. Gas evolution ceased during this period. Pyridine was then removed under vacuum. Water and 3N aq. HCl were then added with stirring. The desired cinnamic acid was then collected through filtration, washed with cold water and dried in a vacuum oven overnight to give 1.56 g (4.8 mmol, 91 %) of white solid.

10

Example 1C(2,4-Dichlorophenyl)[2-(*E*-((6-hydroxyhexylamino)carbonyl)ethenyl)phenyl] sulfide

A suspension of the acid (284 mg, 0.87 mmol) from Example 1B in 5 mL of methylene chloride was stirred with (COCl)₂ (84 μ L, 0.97 mmol), and one drop of DMF under nitrogen atmosphere for 90 minutes. The solvent was then removed under vacuum. The residue (COCl)₂ was removed with benzene (2x) in vacuo. To a separate flask, previously filled with 6-amino-1-hexanol (12 mg, 0.10 mmol), Hunig's base (22.8 μ L, 0.13 mmol) and DMAP (1.1 mg, 0.008 mmol) in 2.0 mL of CH₂Cl₂, the acid chloride (30 mg, 0.087 mmol) in 1.0 mL of CH₂Cl₂ was then dropped in slowly. After 30 minutes, the reaction mixture was poured into 3N HCl and extracted with ethyl acetate (EtOAc). The organic layer was washed with brine, dried with Na₂SO₄, condensed under reduced pressure. The crude product was purified by

20

preparative TLC to give 21.0 mg (90 %) of the title compound as a colorless oil. ¹H NMR (CDCl₃, 300 MHz) δ 1.31-1.48 (m, 4H), 1.48-1.70 (m, 4H), 3.37 (q, *J* = 6.7 Hz, 2H), 3.65 (t, *J* = 6.3 Hz, 2H), 5.63 (br s, 1H), 6.36 (d, *J* = 15.9 Hz, 1H), 6.71 (d, *J* = 9.3 Hz, 1H), 7.05 (dd, *J* = 2.4, 8.7 Hz, 1H), 7.31-7.49 (m, 4H), 7.65 (dd, *J* = 2.1, 7.5 Hz, 1H), 7.99 (d, *J* = 15.9 Hz, 1H). MS (DCI/NH₃) (M+NH₄)⁺ at *m/z* 441, 443, 445.

Example 2

(2,4-Dichlorophenyl)[2-(*E*-((3-(1-imidazolyl)propylamino)carbonyl)ethenyl)phenyl] sulfide

The title compound was prepared by the procedures described in Example 1C substituting 6-amino-1-hexanol with 1-(3-aminopropyl)imidazole. White powder; ¹H NMR (d⁶-DMSO, 300 MHz) δ 1.88 (p, *J* = 7.7 Hz, 2H), 3.11 (q, *J* = 7.7 Hz, 2H), 3.97 (t, *J* = 7.7 Hz, 2H), 6.63 (d, *J* = 15.9 Hz, 1H), 6.70 (d, *J* = 8.7 Hz, 1H), 6.89 (d, *J* = 0.9 Hz, 1H), 7.17 (d, *J* = 0.9 Hz, 1H), 7.33 (dd, *J* = 2.7, 8.7 Hz, 1H), 7.46-7.65 (m, 4H), 7.72 (d, *J* = 2.7 Hz, 1H), 7.78 (d, *J* = 15.9 Hz, 1H), 7.80 (d, *J* = 8.7 Hz, 1H), 8.24 (t, *J* = 5.9 Hz, 1H). MS (DCI/NH₃) (M+H)⁺ at *m/z* 448, 450, 452. Analysis calculated for C₂₁H₁₉N₃O₁Cl₂S₁·0.87 H₂O: C, 56.30; H, 4.67; N, 9.38. Found: C, 56.30; H, 4.56; N, 9.27.

Example 3

(2,4-Dichlorophenyl)[2-chloro-4-(*E*-((2-hydroxyethylamino)carbonyl)ethenyl)phenyl] sulfide

The title compound was prepared by the procedures described in Example 1 substituting 2-chlorobenzaldehyde with 3-chloro-4-fluoro-benzaldehyde, and 6-amino-1-hexanol with ethanolamine. Colorless oil; ^1H NMR (CDCl_3 , 300 MHz) δ 3.57 (q, J = 7.65 Hz, 2H), 3.71 (q, J = 7.65 Hz, 2H), 6.06 (br s, 1H), 6.40 (d, J = 15.3 Hz, 1H), 6.96 (d, J = 8.7 Hz, 1H), 7.22-7.30 (m, 4H), 7.49-7.60 (m, 1H), 7.55 (d, J = 15.3 Hz, 1H). MS (APCI) ($\text{M}+\text{H}$) $^+$ at m/z 402, 404, 406, 408. Analysis calculated for $\text{C}_{17}\text{H}_{14}\text{N}_1\text{O}_2\text{Cl}_3\text{S}_1\cdot 0.25\text{H}_2\text{O}$: C, 50.14; H, 3.59; N, 3.44. Found: C, 50.16; H, 3.62; N, 3.29.

Example 4

(2,4-Dichlorophenyl)[2-chloro-4-(*E*-((6-hydroxyhexylamino)carbonyl)ethenyl)phenyl] sulfide

The title compound was prepared by the procedures described in Example 1 substituting 2-chlorobenzaldehyde with 3-chloro-4-fluoro-benzaldehyde. Colorless oil; ^1H NMR (CDCl_3 , 300 MHz) δ 1.42 (m, 4H), 1.58 (m, 4H), 3.40 (q, J = 6.7 Hz, 2H), 3.65 (br m, 2H), 5.60 (br t, 1H), 6.35 (d, J = 15.3 Hz, 1H), 6.98 (d, J = 8.7 Hz, 1H), 7.22-7.30 (m, 4H), 7.49-7.60 (m, 1H), 7.55 (d, J = 15.3 Hz, 1H). MS (APCI) ($\text{M}+\text{H}$) $^+$ at m/z 458, 460, 462, 464. Analysis calculated for $\text{C}_{21}\text{H}_{22}\text{N}_1\text{O}_2\text{Cl}_3\text{S}_1\cdot 0.27\text{H}_2\text{O}$: C, 54.39; H, 4.90; N, 3.02. Found: C, 54.40; H, 4.85; N, 2.71.

Example 5

(2,4-Dichlorophenyl)[2-chloro-4-(*E*-((bis-(2-hydroxyethyl)amino)carbonyl)ethenyl) phenyl] sulfide

The title compound was prepared by the procedures described in Example 1 substituting 2-chlorobenzaldehyde with 3-chloro-4-fluoro-benzaldehyde, and 6-amino-1-hexanol with diethanolamine. Colorless oil; ¹H NMR (CDCl₃, 300 MHz) δ 2.99 (br s, 2H), 3.67 (br m, 4H), 3.88 (t, *J* = 5.1 Hz, 2H), 3.94 (t, *J* = 5.1 Hz, 2H), 6.94 (d, *J* = 15.3 Hz, 1H), 6.97 (d, *J* = 8.7 Hz, 1H), 7.21-7.32 (m, 3H), 7.50-7.54 (m, 1H), 7.58 (d, *J* = 2.4 Hz, 1H), 7.58 (d, *J* = 15.3 Hz, 1H). MS (APCI) (M+H)⁺ at *m/z* 446, 448, 450, 452. Analysis calculated for C₁₉H₁₈N₁O₃Cl₃S₁·1.09H₂O: C, 48.93; H, 4.36; N, 3.00. Found: C, 48.88; H, 4.00; N, 3.01.

Example 6

(2,4-Dichlorophenyl)[2-chloro-4-(*E*-(3-(1-pyrrolidin-2-yl)propylamino)carbonyl)ethenyl]phenyl sulfide

The title compound was prepared by the procedures described in Example 1 substituting 2-chlorobenzaldehyde with 3-chloro-4-fluoro-benzaldehyde, and 6-amino-1-hexanol with 1-(3-aminopropyl)-2-pyrrolidinone. Colorless oil; ¹H NMR (CDCl₃, 300 MHz) δ 1.74 (qu, *J* = 6.0 Hz, 2H), 2.09 (qu, *J* = 7.5 Hz, 2H), 2.45 (t, *J* = 8.25 Hz, 2H), 3.33 (q, *J* = 6.0 Hz, 2H), 3.42 (q, *J* = 8.25 Hz, 4H), 6.46 (d, *J* = 15.6 Hz, 1H), 7.02 (d, *J* = 8.7 Hz, 1H), 7.14-7.23 (m, 2H), 7.30 (dd, *J* = 2.4, 8.7 Hz, 1H), 7.51 (d, *J* = 2.4 Hz, 1H), 7.51 (d, *J* = 15.6 Hz, 1H), 7.60 (d, *J* = 2.1 Hz, 1H). MS (DCI/NH₃) (M+H)⁺ at *m/z* 483, 485, 487, 489. Analysis calculated for C₂₂H₂₁N₂O₂Cl₃S₁·0.57H₂O: C, 53.48; H, 4.52; N, 5.67. Found: C, 53.49; H, 4.60; N, 5.65.

Example 7

(2,4-Dichlorophenyl)[2-chloro-4-(*E*-((1-morpholinyl)carbonyl)ethenyl)phenyl]

sulfide

The title compound was prepared by the procedures described in Example 1 substituting 2-chlorobenzaldehyde with 3-chloro-4-fluoro-benzaldehyde, and 6-amino-1-hexanol with morpholine. White solid; ¹H NMR (CDCl₃, 300 MHz) δ 3.59-3.80 (m, 8H), 6.83 (d, *J* = 15.6 Hz, 1H), 6.97 (d, *J* = 8.7 Hz, 1H), 7.16-7.32 (m, 3H), 7.49-7.53 (m, 1H), 7.59 (d, *J* = 2.4 Hz, 1H), 7.59 (d, *J* = 15.6 Hz, 1H). MS (DCI/NH₃) (M+H)⁺ at *m/z* 428, 430, 432, 434. Analysis calculated for C₁₉H₁₆N₁O₂Cl₃S₁·0.46H₂O: C, 52.22; H, 3.90; N, 3.20. Found: C, 52.20; H, 3.76; N, 3.12.

Example 8

(2,4-Dichlorophenyl)[2-chloro-4-(*E*-((4-methylpiperazin-1-yl)carbonyl)

ethenyl)phenyl] sulfide

The title compound was prepared by the procedures described in Example 1 substituting 2-chlorobenzaldehyde with 3-chloro-4-fluoro-benzaldehyde, and 6-amino-1-hexanol with 1-methylpiperazine. Colorless oil; ¹H NMR (CDCl₃, 300 MHz) δ 2.37 (s, 3H), 2.51 (br m, 4H), 3.63-3.87 (br m, 4H), 6.85 (d, *J* = 15.6 Hz, 1H), 6.98 (d, *J* = 8.7 Hz, 1H), 7.19-7.25 (m, 2H), 7.27 (dd, *J* = 2.1, 8.7 Hz, 1H), 7.52 (t, *J* = 0.9 Hz, 1H), 7.57 (d, *J* = 15.6 Hz, 1H), 7.60 (d, *J* = 2.1 Hz, 1H). MS (DCI/NH₃) (M+H)⁺ at *m/z* 441, 443, 445, 447. Analysis calculated for C₂₀H₁₉N₂O₁Cl₃S₁·0.45H₂O: C, 53.39; H, 4.46; N, 6.23. Found: C, 53.37; H, 4.46; N, 6.07.

00541795.03100

Example 9

(2,4-Dichlorophenyl)[2-chloro-4-(*E*-((4-acetyl)piperazin-1-yl)carbonyl)
ethenyl]phenyl] sulfide

5 The title compound was prepared by the procedures described in Example 1 substituting 2-chlorobenzaldehyde with 3-chloro-4-fluoro-benzaldehyde, and 6-amino-1-hexanol with 1-acetyl-piperazine. White solid; ¹H NMR (CDCl₃, 300 MHz) δ 2.15 (s, 3H), 3.50-3.58 (m, 2H), 3.58-3.85 (m, 6H), 6.85 (d, *J* = 15.3 Hz, 1H), 6.96 (d, *J* = 8.7 Hz, 1H), 7.24-7.36 (m, 3H), 7.54 (d, *J* = 2.4 Hz, 1H), 7.61 (d, *J* = 15.3 Hz, 1H),
10 7.61 (d, *J* = 2.1 Hz, 1H). MS (DCI/NH₃) (M+H)⁺ at *m/z* 486, 488, 490, 492. Analysis calculated for C₂₁H₁₉N₂O₂Cl₃S₁·0.85H₂O: C, 51.99; H, 4.30; N, 5.77. Found: C, 52.03; H, 4.27; N, 5.67.

Example 10

15 (2,4-Dichlorophenyl)[2-chloro-4-(*E*-((4-(2-pyridyl)piperazin-1-yl)carbonyl)
ethenyl]phenyl] sulfide

The title compound was prepared by the procedures described in Example 1 substituting 2-chlorobenzaldehyde with 3-chloro-4-fluoro-benzaldehyde, and 6-amino-1-hexanol with 1-(2-pyridyl)piperazine. White solid; ¹H NMR (CDCl₃, 300 MHz) δ
20 3.59 (br m, 2H), 3.69 (br m, 2H), 3.78 (br m, 2H), 3.86 (br m, 2H), 6.64-6.72 (m, 2H), 6.90 (d, *J* = 15.6 Hz, 1H), 6.99 (d, *J* = 8.7 Hz, 1H), 7.22-7.25 (m, 2H), 7.31(dd, *J* = 2.4, 8.7 Hz, 1H), 7.49-7.57 (m, 2H), 7.61 (d, *J* = 15.6 Hz, 1H), 7.62 (d, *J* = 2.4 Hz,

1H), 8.19-8.24 (m, 1H). MS (DCI/NH₃) (M+H)⁺ at *m/z* 504, 506, 508, 510. Analysis calculated for C₂₄H₂₀N₃O₁Cl₃S₁: C, 57.10; H, 3.99; N, 8.32. Found: C, 57.12; H, 4.06; N, 8.29.

5

Example 11

(2-(Hydroxymethyl)phenyl)[2-chloro-4-(*E*-((1-morpholinyl)carbonyl)ethenyl)phenyl] sulfide

The title compound was prepared by the procedures described in Example 1 substituting 2,4-dichlorothiophenol with 2-mercaptobenzyl alcohol, 2-

- 10 chlorobenzaldehyde with 3-chloro-4-fluoro-benzaldehyde, and 6-amino-1-hexanol with morpholine. White solid; ¹H NMR (CDCl₃, 300 MHz) δ 3.50-3.62 (br m, 6H), 3.65-3.74 (br m, 2H), 4.54 (d, *J* = 5.7 Hz, 2H), 5.33 (t, *J* = 5.7 Hz, 1H), 6.62 (d, *J* = 8.7 Hz, 1H), 7.28 (d, *J* = 15.0 Hz, 1H), 7.36 (d, *J* = 7.8 Hz, 1H), 7.42 (d, *J* = 15.0 Hz, 1H), 7.43 (dd, *J* = 1.8, 8.7 Hz, 1H), 7.50 (dd, *J* = 2.1, 8.7 Hz, 1H), 7.55 (dd, *J* = 2.1, 7.8 Hz, 1H), 7.68 (dd, *J* = 1.5, 8.1 Hz, 1H), 8.02 (d, *J* = 2.1 Hz, 1H). MS (DCI/NH₃) (M+H)⁺ at *m/z* 390, 392. Analysis calculated for C₂₀H₂₀N₁O₃Cl₁S₁ · 0.09H₂O: C, 61.35; H, 5.20; N, 3.58. Found: C, 61.37; H, 5.48; N, 3.81.
- 15

Example 12

- 20 (2-Bromophenyl)[2-chloro-4-(*E*-((1-morpholinyl)carbonyl)ethenyl)phenyl] sulfide

The title compound was prepared by the procedures described in Example 1 substituting 2,4-dichlorothiophenol with 2-bromothiophenol, 2-chlorobenzaldehyde

with 3-chloro-4-fluoro-benzaldehyde, and 6-amino-1-hexanol with morpholine. White solid; ^1H NMR (d_6 -DMSO, 300 MHz) δ 3.50-3.66 (br m, 6H), 3.66-3.79 (br m, 2H), 7.05 (d, $J = 8.7$ Hz, 1H), 7.26 (dd, $J = 2.1, 8.1$ Hz, 1H), 7.33 (dd, $J = 2.1, 8.1$ Hz, 1H), 7.36 (d, $J = 15.6$ Hz, 1H), 7.39 (dd, $J = 1.8, 12.0$ Hz, 1H), 7.45 (dd, $J = 1.8, 6.3$ Hz, 1H), 7.48 (d, $J = 15.6$ Hz, 1H), 7.64 (dd, $J = 2.1, 8.7$ Hz, 1H), 7.80 (dd, $J = 2.8, 8.7$ Hz, 1H), 8.09 (d, $J = 2.1$ Hz, 1H). MS (DCI/ NH_3) ($\text{M}+\text{H}$) $^+$ at m/z 438, 440, 442.

Example 13

(2,4-Dichlorophenyl)[2-chloro-4-(*E*-((4-(2-hydroxyethyl)piperazin-1-yl)carbonyl)ethenyl)phenyl] sulfide

10

The title compound was prepared by the procedures described in Example 1 substituting 2-chlorobenzaldehyde with 3-chloro-4-fluoro-benzaldehyde, and 6-amino-1-hexanol with 1-hydroxyethylpiperazine. Colorless oil; ^1H NMR (CDCl_3 , 300 MHz) δ 2.85-3.20 (br m, 6H), 3.84-4.19 (m, 6H), 6.80 (d, $J = 15.3$ Hz, 1H), 6.94 (d, $J = 8.7$ Hz, 1H), 7.22-7.38 (m, 3H), 7.50-7.56 (m, 1H), 7.56-7.62 (m, 1H), 7.60 (d, $J = 15.3$ Hz, 1H). MS (DCI/ NH_3) ($\text{M}+\text{H}$) $^+$ at m/z 471, 473, 475, 477.

15

Example 14

(2,4-Dichlorophenyl)[2-chloro-4-(*E*-((4-(2-hydroxyethoxyethyl)piperazin-1-yl)carbonyl)

20

ethenyl)phenyl] sulfide

The title compound was prepared by the procedures described in Example 1 substituting 2-chlorobenzaldehyde with 3-chloro-4-fluoro-benzaldehyde, and 6-amino-1-hexanol with 1-[2-(2-hydroxyethoxy)ethyl]piperazine. Colorless oil; ¹H NMR (CDCl₃, 300 MHz) δ 2.73 (br m, 6H), 3.58-3.68 (m, 2H), 3.68-4.00 (m, 8H), 6.84 (d, *J* = 15.3 Hz, 1H), 6.97 (d, *J* = 8.7 Hz, 1H), 7.20-7.34 (m, 3H), 7.54 (d, *J* = 7.5 Hz, 1H), 7.58 (d, *J* = 15.3 Hz, 1H), 7.58-7.65 (overlapping d, 1H). MS (DCI/NH₃) (M+H)⁺ at *m/z* 515, 517, 519, 521.

Example 15

10 (2-Bromophenyl)[2-chloro-4-(*E*-((3-(hydroxymethyl)piperidin-1-yl)carbonyl)ethenyl]phenyl] sulfide

The title compound was prepared by the procedures described in Example 1 substituting 2,4-dichlorothiophenol with 2-bromothiophenol, 2-chlorobenzaldehyde with 3-chloro-4-fluoro-benzaldehyde, and 6-amino-1-hexanol with 3-hydroxymethylpiperidine. ¹H NMR (DMSO-d₆, 300MHz) δ 8.07 (d, *J* = 17.7 Hz, 1H), 7.80 (d, *J* = 7.7 Hz, 1H), 7.63 (br d, *J* = 7.7 Hz, 1H), 7.44 (d, *J* = 7.0 Hz, 1H), 7.40 (br s, 2H), 7.35 (m, 1H), 7.25 (dd 7.7, 1.5, 1H), 7.06 (dd, *J* = 8.1, 2.9, 1H), 4.57 (m, 1H), 4.45 (m, 1H), 4.16 (br m, 2H), 1.2 – 1.8 (m, 8H). HRMS calculated for C₂₁H₂₁N₁O₂S₁Br₁Cl₁: 466.0243. Observed: 466.0247.

20

Example 16

(2-Bromophenyl)[2-chloro-4-(*E*-((2-(hydroxymethyl)piperidin-1-yl)carbonyl)

ethenyl)phenyl] sulfide

The title compound was prepared by the procedures described in Example 1 substituting 2,4-dichlorothiophenol with 2-bromothiophenol, 2-chlorobenzaldehyde with 3-chloro-4-fluoro-benzaldehyde, and 6-amino-1-hexanol with 2-

- 5 hydroxymethylpiperidine. ^1H NMR (DMSO- d_6 , 300MHz) δ 8.03 (m, 1H), 7.79 (d, J = 7.8 Hz, 1H), 7.61 (m, 1H), 7.30 – 7.45 (m, 4H), 7.23 (m, 1H), 7.07 (m, 1H), 4.79 (m, 2H), 4.61 (m, 2H), 4.10 (m, 1H), 1.50 (m, 6H). HRMS calculated for $\text{C}_{21}\text{H}_{21}\text{N}_1\text{O}_2\text{S}_1\text{Br}_1\text{Cl}_1$: 466.0243. Observed: 466.0247.

10

Example 17(2-Bromophenyl)[2-chloro-4-(*E*-(3-acetamidopyrrolidin-1-yl)carbonyl)ethenyl)phenyl] sulfide

The title compound was prepared by the procedures described in Example 1 substituting 2,4-dichlorothiophenol with 2-bromothiophenol, 2-chlorobenzaldehyde

- 15 with 3-chloro-4-fluoro-benzaldehyde, and 6-amino-1-hexanol with 3-acetamidopyrrolidine. ^1H NMR (DMSO- d_6 , 300MHz) δ 8.14 (m, 1H), 8.07 (dd, J = 9.8, 1.7 Hz, 1H), 7.80 (d, J = 7.8 Hz, 1H), 7.64 (dd, J = 8.1, 1.7 Hz, 1H), 7.25 – 7.47 (m, 4H), 7.10 (t, J = 7.8 Hz, 1H), 7.03 (dd, J = 8.1, 1.7 Hz, 1H), 3.45 – 4.34 (m, 6H), 2.02 (m, 2H), 1.81 (ap d, J = 1.4 Hz, 1H). HRMS calculated for $\text{C}_{21}\text{H}_{20}\text{N}_2\text{O}_2\text{S}_1\text{Br}_1\text{Cl}_1$: 479.0196. Observed: 479.0183.
- 20

Example 18

(2-Bromophenyl)[2-chloro-4-(*E*-((4-hydroxypiperidin-1-yl)carbonyl)
ethenyl)phenyl] sulfide

The title compound was prepared by the procedures described in Example 1 substituting 2,4-dichlorothiophenol with 2-bromothiophenol, 2-chlorobenzaldehyde with 3-chloro-4-fluoro-benzaldehyde, and 6-amino-1-hexanol with 4-hydroxypiperidine. ¹H NMR (DMSO-d₆, 300MHz) δ 8.08 (d, *J* = 1.7 Hz, 1H), 7.80 (dd, *J* = 8.0, 1.5 Hz, 1H), 7.63 (dd, *J* = 8.3, 1.9 Hz, 1H), 7.44 (ap dd, *J* = 7.5, 1.4 Hz, 2H), 7.40 (ap d, *J* = 3.7 Hz, 2H), 7.34 (dt, *J* = 7.6, 1.8 Hz, 1H), 7.25 (dd, *J* = 7.5, 1.7 Hz, 1H), 7.05 (d, *J* = 8.1 Hz, 1H), 4.76 (br s, 1H), 4.01 (m, 2H), 3.72 (m, 1H), 3.12 (m, 1H), 1.75 (m, 2H), 1.32 (m, 2H). HRMS calculated for C₂₀H₁₉N₁O₂S₁Br₁Cl₁: 452.0087. Observed: 452.0076.

Example 19

(2-Bromophenyl)[2-chloro-4-(*E*-((piperidin-1-yl)carbonyl) ethenyl)phenyl] sulfide

The title compound was prepared by the procedures described in Example 1 substituting 2,4-dichlorothiophenol with 2-bromothiophenol, 2-chlorobenzaldehyde with 3-chloro-4-fluoro-benzaldehyde, and 6-amino-1-hexanol with piperidine. ¹H NMR (DMSO-d₆, 300MHz) δ 8.08 (d, *J* = 1.7 Hz, 1H), 7.80 (dd, *J* = 8.1, 1.4 Hz, 1H), 7.63 (dd, *J* = 8.1, 1.7 Hz, 1H), 7.44 (ap dd, *J* = 7.6, 1.5 Hz, 1H), 7.39 (ap d, *J* = 4.8 Hz, 2H), 7.34 (dt, *J* = 7.5, 1.6, 1H), 7.24 (dd, *J* = 7.5, 1.7, 1H), 7.05 (d, *J* = 8.1 Hz, 1H), 3.65 (br m, 2H), 3.53 (br m, 2H), 1.62 (br m, 2H), 1.50 (br m, 4H). HRMS calculated for C₂₀H₁₉N₁O₁S₁Br₁Cl₁: 436.0130. Observed: 436.0122.

00544795.033100

Example 20

(2,4-Dichlorophenyl)[2-chloro-4-(*E*-((3-carboxypiperidin-1-yl)carbonyl)
ethenyl)phenyl] sulfide

5

The title compound was prepared by the procedures described in Example 1 substituting 2-chlorobenzaldehyde with 3-chloro-4-fluoro-benzaldehyde, and 6-amino-1-hexanol with nipecotic acid. Colorless oil; ¹H NMR (CDCl₃, 300 MHz) δ 1.44-1.68 (br m, 1H), 1.68-2.00 (br m, 2H), 2.51-2.67 (br m, 1H), 3.13-3.37 (br m, 1H), 3.80-10 4.12 (br m, 1H), 4.30-5.00 (br m, 3H), 6.86 (d, *J* = 15.3 Hz, 1H), 6.99 (d, *J* = 8.7 Hz, 1H), 7.16-7.24 (m, 2H), 7.29 (d, *J* = 8.7 Hz, 1H), 7.47-7.55 (m, 1H), 7.55 (d, *J* = 15.3 Hz, 1H), 7.60 (br d, 1H). MS (APCI) (M+H)⁺ at *m/z* 470, 472, 474, 476.

Example 21

15 (2,4-Dichlorophenyl)[2-chloro-4-(*E*-((4-carboxypiperidin-1-yl)carbonyl)
ethenyl)phenyl] sulfide

The title compound was prepared by the procedures described in Example 1 substituting 2-chlorobenzaldehyde with 3-chloro-4-fluoro-benzaldehyde, and 6-amino-1-hexanol with isonipecotic acid. Colorless oil; ¹H NMR (CDCl₃, 300 MHz) δ 1.68-20 1.85 (m, 2H), 1.98-2.09 (m, 2H), 2.60-2.72 (m, 1H), 2.90-3.13 (br m, 1H), 3.17-3.38 (br m, 1H), 3.93-4.12 (br m, 1H), 4.38-4.59 (br m, 1H), 6.86 (d, *J* = 15.3 Hz, 1H), 6.99 (dd, *J* = 8.7 Hz, 1H), 7.20-7.25 (m, 2H), 7.28 (dd, *J* = 1.8, 8.7 Hz, 1H), 7.49-7.53

(m, 1H), 7.56 (d, $J = 15.3$ Hz, 1H), 7.60 (d, $J = 1.8$ Hz, 1H). MS (APCI) $(M+H)^+$ at m/z 470, 472, 474, 476.

Example 22

5 (2-Bromophenyl)[2-chloro-4-(*E*-((4-acetylhomopiperazin-1-yl)carbonyl)ethenyl)phenyl] sulfide

The title compound was prepared by the procedures described in Example 1 substituting 2,4-dichlorothiophenol with 2-bromothiophenol, 2-chlorobenzaldehyde with 3-chloro-4-fluoro-benzaldehyde, and 6-amino-1-hexanol with 4-

10 acetylhomopiperazine. ^1H NMR (DMSO- d_6 , 300MHz) δ 8.10 (m, 1H), 7.81 (d, $J = 7.7$ Hz, 1H), 7.64 (m, 1H), 7.24 – 7.51 (m, 5H), 7.05 (m, 1H), 3.39 – 3.77 (m, 8H), 1.97 (m, 3H), 1.68 (m, 2H). HRMS calculated for $C_{22}H_{22}N_2O_2S_1Br_1Cl_1$: 493.0352. Observed: 493.0352.

Example 23

15 (2-Bromophenyl)[2-chloro-4-(*E*-((thiomorpholin-1-yl)carbonyl)ethenyl)phenyl] sulfide

The title compound was prepared by the procedures described in Example 1 substituting 2,4-dichlorothiophenol with 2-bromothiophenol, 2-chlorobenzaldehyde with 3-chloro-4-fluoro-benzaldehyde, and 6-amino-1-hexanol with thiomorpholine. ^1H NMR (DMSO- d_6 , 300MHz) δ 8.10 (d, $J = 1.5$ Hz, 1H), 7.80 (d, $J = 8.5$ Hz, 1H), 7.64 (dd, $J = 8.1, 1.5$ Hz, 1H), 7.31 – 7.48 (m, 4H), 7.36 (m, 1H), 7.26 (dd, $J = 8.1, 1.8$ Hz,

1H), 7.05 (d $J = 8.1$ Hz, 1H), 3.96 (m, 2H), 3.82 (m, 2H), 2.62 (m, 4H). HRMS calculated for $C_{19}H_{17}N_1O_1S_2Br_1Cl_1$: 455.9681. Observed: 455.9676.

Example 24

5 (2-Bromophenyl)[2-chloro-4-(*E*-((4-(1-benzimidazol-2-onyl)piperidin-1-yl)carbonyl)ethenyl)phenyl] sulfide

The title compound was prepared by the procedures described in Example 1 substituting 2,4-dichlorothiophenol with 2-bromothiophenol, 2-chlorobenzaldehyde with 3-chloro-4-fluoro-benzaldehyde, and 6-amino-1-hexanol with 4-(1-benzimidazol-2-onyl)piperidine. 1H NMR (DMSO- d_6 , 300MHz) δ 8.14 (d, $J = 1.5$ Hz, 1H), 7.80 (dd, $J = 7.9, 1.3$ Hz, 1H), 7.67 (dd, $J = 8.1, 1.8$ Hz, 1H), 7.48 (ap s, 2H), 7.44 (dt, $J = 7.5, 1.2$, 1H), 7.34 (dt, $J = 7.6, 1.6$, 1H), 7.26 (dd, $J = 7.7, 1.8$ Hz, 1H), 7.22 (m, 1H), 7.06 (d, $J = 8.1$, 1H), 6.97 (ap d, $J = 2.6$, 3H), 4.64 (m, 1H), 4.48 (m, 2H), 2.79 (m, 2H), 2.29 (m, 2H), 1.78 (m, 2H). HRMS calculated for $C_{27}H_{23}N_3O_2S_1Br_1Cl_1$: 568.0461. Observed: 568.0477.

Example 25

(2-Bromophenyl)[2-chloro-4-(*E*-((2-tetrahydroisoquinolinyl)carbonyl)ethenyl)phenyl] sulfide

20 The title compound was prepared by the procedures described in Example 1 substituting 2,4-dichlorothiophenol with 2-bromothiophenol, 2-chlorobenzaldehyde with 3-chloro-4-fluoro-benzaldehyde, and 6-amino-1-hexanol with

tetrahydroisoquinoline. ^1H NMR ($\text{DMSO}-d_6$, 300MHz) δ 8.12 (d, $J = 7.4$ Hz, 1H), 7.81 (dd, $J = 7.7$, 1.1 Hz, 1H), 7.67 (dd, $J = 8.3$, 1.3 Hz, 1H), 7.47 (m, 2H), 7.43 (dd, $J = 7.5$, 1.3 Hz, 2H), 7.34 (dt, $J = 7.6$, 1.7 Hz, 1H), 7.27 (d 7.7 Hz, 1H), 7.19 (m, 4H), 7.05 (d, $J = 8.1$ Hz, 1H), 4.92 (s, 1H), 4.72 (s, 1H), 3.95 (t, $J = 5.9$ Hz, 1H), 3.78 (t, $J = 5.7$ Hz, 1H), 2.89 (t, $J = 5.3$ Hz, 1H), 2.83 (t, $J = 3.7$, 1H). HRMS calculated for $\text{C}_{24}\text{H}_{19}\text{N}_4\text{O}_2\text{S}_1\text{Br}_1\text{Cl}_1$: 484.0138. Observed: 484.0128.

Example 26

(2-Methylphenyl)[2-trifluoromethyl-4-(*E*-((4-acetyl)piperazin-1-yl)carbonyl)

10 ethenyl)phenyl] sulfide

The title compound was prepared by the procedures described in Example 1 substituting 2,4-dichlorothiophenol with 2-methylthiophenol, 2-chlorobenzaldehyde with 4-fluoro-3-trifluoromethylbenzaldehyde, and 6-amino-1-hexanol with 1-acetyl piperazine. ^1H NMR (CDCl_3 , 300MHz) δ 7.79 (s, 1H); 7.63 (d, $J = 15.4$ Hz, 1H); 7.51 (d, $J = 6.8$ Hz, 1H); 7.41-7.33 (m, 3H); 7.28 (m, 1H); 6.83 (d, $J = 15.4$ Hz, 1H); 6.79 (d, $J = 6.8$ Hz, 1H); 3.80-3.60 (m, 6H); 3.57-3.50 (m, 2H); 2.34 (s, 3H); 2.14 (s, 3H). MS (ESI) m/z 919 ($2\text{M}+\text{Na}$) $^+$, 897 ($2\text{M}+\text{H}$) $^+$, 471 ($\text{M}+\text{Na}$) $^+$, 449 ($\text{M}+\text{H}$) $^+$.

Example 27

20 (2-Methylphenyl)[2-trifluoromethyl-4-(*E*-((1-morpholinyl)carbonyl)

ethenyl)phenyl] sulfide

The title compound was prepared by the procedures described in Example 1 substituting 2,4-dichlorothiophenol with 2-methylthiophenol, 2-chlorobenzaldehyde with 4-fluoro-3-trifluoromethylbenzaldehyde, and 6-amino-1-hexanol with morpholine. ¹H NMR (CDCl₃, 300MHz) δ 7.79 (s, 1H); 7.63 (d, J = 14.0 Hz, 1H); 7.52 (d, J = 7.6 Hz, 1H); 7.40-7.30 (m, 3H); 7.28 (m, 1H); 6.87 (d, J = 14.0 Hz, 1H); 6.84 (d, J = 7.6 Hz, 1H); 3.73 (br s, 8H); 2.34 (s, 3H). MS (ESI) *m/z* 837 (2M+Na)⁺, 815 (2M+H)⁺, 408 (M+H)⁺.

Example 28

10 (2-Methylphenyl)[2-trifluoromethyl-4-(E-((2-(1-morpholinyl)ethylamino)carbonyl)ethenyl)phenyl] sulfide

The title compound was prepared by the procedures described in Example 1 substituting 2,4-dichlorothiophenol with 2-methylthiophenol, 2-chlorobenzaldehyde with 4-fluoro-3-trifluoromethylbenzaldehyde, and 6-amino-1-hexanol with 2-(1-

15 morpholinyl)ethylamine. ¹H NMR (CDCl₃, 300MHz) δ 7.80 (s, 1H); 7.56 (d, J = 15.8 Hz, 1H); 7.50 (d, J = 8.1 Hz, 1H); 7.40-7.32 (m, 3H); 7.28 (m, 1H); 6.79 (d, J = 15.8 Hz, 1H); 6.40 (d, J = 8.1 Hz, 1H); 3.75 (t, J = 4.6 Hz, 4H); 3.51 (q, J = 5.5 Hz, 2H), 2.57 (t, J = 5.8 Hz, 2H); 2.55-2.48 (m, 4H); 2.34 (s, 3H). MS (ESI) *m/z* 923 (2M+Na)⁺, 473 (M+Na)⁺, 451 (M+H)⁺.

Example 29

(2-Methylphenyl)[2-trifluoromethyl-4-(E-((4-phenylpiperazin-1-yl)carbonyl)

ethenyl)phenyl] sulfide

The title compound was prepared by the procedures described in Example 1 substituting 2,4-dichlorothiophenol with 2-methylthiophenol, 2-chlorobenzaldehyde with 4-fluoro-3-trifluoromethylbenzaldehyde, and 6-amino-1-hexanol with 4-

- 5 phenylpiperazine. ¹H NMR (CDCl₃, 300MHz) δ 7.81 (s, 1H); 7.64 (d, J = 16.0 Hz, 1H); 7.51 (d, J = 8.2 Hz, 1H); 7.40-7.27 (m, 6H); 6.98-6.90 (m, 4H); 6.80 (d, J = 8.2 Hz, 1H); 3.88 (br s, 4H); 2.23 (br s, 4H); 2.34 (s, 3H). MS (ESI) *m/z* 987 (2M+Na)⁺, 965 (2M+H)⁺, 505 (M+Na)⁺, 483 (M+H)⁺, 451.

10

Example 30(2-Methylphenyl)[2-trifluoromethyl-4-(E-((3-(1-pyrrolidin-2-only)propylamino)carbonyl)ethenyl)phenyl] sulfide

The title compound was prepared by the procedures described in Example 1 substituting 2,4-dichlorothiophenol with 2-methylthiophenol, 2-chlorobenzaldehyde with 4-fluoro-3-trifluoromethylbenzaldehyde, and 6-amino-1-hexanol with 1-

- 15 pyrrolidin-2-only)propylamine. ¹H NMR (CDCl₃, 300MHz) δ 7.78 (s, 1H); 7.53 (d, J = 15.6 Hz, 1H); 7.49 (d, J = 7.2 Hz, 1H); 7.40-7.33 (m, 3H); 7.14 (m, 1H); 6.80 (d, J = 8.2 Hz, 1H); 6.43 (d, J = 15.6 Hz, 1H); 3.41 (m, 4H); 3.32 (q, J = 6.1 Hz, 2H); 2.43 (t, 20 J = 6.6 Hz, 2H); 2.34 (s, 3H), 2.08 (m, 2H), 1.75 (m, 2H). MS (ESI) *m/z* 947 (2M+Na)⁺, 925 (2M+H)⁺, 485 (M+Na)⁺, 463 (M+H)⁺.

Example 31(2-Methylphenyl)[2-trifluoromethyl-4-(E-((cyclopropylamino)carbonyl)ethenyl)phenyl] sulfide

The title compound was prepared by the procedures described in Example 1

5 substituting 2,4-dichlorothiophenol with 2-methylthiophenol, 2-chlorobenzaldehyde with 4-fluoro-3-trifluoromethylbenzaldehyde, and 6-amino-1-hexanol with cyclopropylamine. ¹H NMR (CDCl₃, 300MHz) δ 7.76 (s, 1H); 7.56 (d, J = 15.4 Hz, 1H); 7.50 (d, J = 8.4 Hz, 1H); 7.40-7.30 (m, 3H); 7.28 (m, 1H); 6.88 (d, J = 8.4 Hz, 1H); 6.30 (d, J = 15.4 Hz, 1H); 5.70 (br s, 1H), 2.95 (m, 1H); 2.34 (s, 3H); 0.85 (m, 10 2H); 0.57 (m, 2H). MS (ESI) *m/z* 777 (2M+Na)⁺, 755 (2M+H)⁺, 400 (M+Na)⁺, 378 (M+H)⁺.

Example 32(2,4-Dichlorophenyl)[2-nitro-4-(E-((4-acetyl)piperazin-1-yl)carbonyl)ethenyl)phenyl] sulfide

15

Example 32A1-Chloro-2-nitro-4-(E-((4-acetyl)piperazin-1-yl)carbonyl)ethenyl) benzene

To a stirred solution of *trans*-4-chloro-3-nitrocinnamic acid (1.50 g, 6.59 mmol)

20 and 1-acetylpiperazine (0.89 g, 6.94 mmol) in 20 mL of DMF at room temperature was added EDAC (1.4 g, 7.30 mmol). The mixture was then stirred at room temperature for 2 hours. TLC indicated the complete consumption of the acid. Water

0011795-03310
0120-567450

was then added to quench the reaction and to precipitate out the product. Cinnamide was then collected through filtration and washed with cold water. The light yellow product was dried in vacuum oven overnight at 40 °C to give 2.04 g (6.03 mmol, 91.6 %) of the title compound.

5

Example 32B

(2,4-Dichlorophenyl)[2-nitro-4-(*E*-((4-acetylpiperazin-1-yl)carbonyl) ethenyl)phenyl] sulfide

To a stirred solution of 4-chloro-3-nitro-cinnamide (275 mg, 0.814 mmol) from
10 Example 32A in 1.0 mL of DMF was added potassium carbonate (169 mg, 1.22
mmol), followed by the dropwise addition of 2,4-dichlorothiophenol (146 mg, 0.815
mmol). The mixture was then stirred at room temperature for 60 minutes.
Completion of the reaction was indicated by the TLC. Water was then added to
precipitate the product. Filtration, washing with cold water, and drying in a vacuum
15 oven afforded 350 mg (0.728 mmol, 89%) of the titled compound as light yellow
solid. ¹H NMR (d⁶-DMSO, 300 MHz) δ 2.05 (s, 3H), 3.42-3.50 (br m, 4H), 3.50-3.64
(br m, 2H), 3.64-3.79 (br m, 2H), 6.83 (d, *J* = 8.7 Hz, 1H), 7.44 (d, *J* = 15.3 Hz, 1H),
7.55 (d, *J* = 15.3 Hz, 1H), 7.63 (dd, *J* = 2.7, 8.7 Hz, 1H), 7.83 (d, *J* = 8.7 Hz, 1H), 7.93
(d, *J* = 8.7 Hz, 1H), 7.96 (d, *J* = 2.7 Hz, 1H), 8.69 (d, *J* = 1.8 Hz, 1H). MS (DCI/NH₃)
20 (M+H)⁺ at *m/z* 497, 499, 501. Analysis calculated for C₂₁H₁₉N₃O₄ Cl₂ S₁•0.82H₂O: C,
50.94; H, 4.20; N, 8.49. Found: C, 50.91; H, 4.21; N, 8.69.

00720-56Z4560

Example 33

(2,4-Dichlorophenyl)[2-nitro-4-(E-((3-(1-pyrrolidin-2-onyl)propylamino)carbonyl)

ethenyl)phenyl] sulfide

The title compound was prepared by the procedures described in Example 32 substituting 1-acetylpiperazine with 1-(3-aminopropyl)-2-pyrrolidinone. Light-yellow powder; ¹H NMR (d⁶-DMSO, 300 MHz) δ 1.64 (p, *J* = 7.1 Hz, 2H), 1.91 (p, *J* = 7.5 Hz, 2H), 2.21 (t, *J* = 8.3 Hz, 2H), 3.15 (q, *J* = 6.3 Hz, 2H), 3.21 (dd, *J* = 9.9, 17.7 Hz, 2H), 3.32 (overlapping t, *J* = 8.4 Hz, 2H), 6.72 (d, *J* = 15.6 Hz, 1H), 6.86 (d, *J* = 8.7 Hz, 1H), 7.46 (d, *J* = 15.6 Hz, 1H), 7.63 (dd, *J* = 2.4, 8.1 Hz, 1H), 7.79 (dd, *J* = 2.4, 8.7 Hz, 1H), 7.84 (d, *J* = 8.7 Hz, 1H), 7.96 (d, *J* = 2.4 Hz, 1H), 8.18 (t, *J* = 6.0 Hz, 1H), 8.46 (d, *J* = 2.1 Hz, 1H). MS (DCI/NH₃) (M+H)⁺ at *m/z* 494, 496.

Example 34

(2,3-Dichlorophenyl)[2-nitro-4-(E-((4-acetylpiperazin-1-yl)carbonyl)

ethenyl)phenyl] sulfide

The title compound was prepared by the procedures described in Example 32B substituting 2,4-dichlorothiophenol with 2,3-dichlorothiophenol. Light-yellow powder; ¹H NMR (d⁶-DMSO, 300 MHz) δ 2.04 (s, 3H), 3.42-3.50 (br m, 4H), 3.50-3.64 (br m, 2H), 3.64-3.79 (br m, 2H), 6.88 (d, *J* = 8.7 Hz, 1H), 7.45 (d, *J* = 15.6 Hz, 1H), 7.55 (t, *J* = 7.65 Hz, 1H), 7.57 (d, *J* = 15.6 Hz, 1H), 7.78 (dd, *J* = 1.8, 8.1 Hz, 1H), 7.87 (dd, *J* = 1.8, 8.1 Hz, 1H), 7.95 (dd, *J* = 2.7, 9.0 Hz, 1H), 8.69 (d, *J* = 1.8 Hz, 1H). MS (DCI/NH₃) (M+H)⁺ at *m/z* 497, 499, 501.

Example 35

(4-Bromophenyl)[2-nitro-4-(*E*-(4-acetyl)piperazin-1-yl)carbonyl]ethenyl]phenyl]

sulfide

5 The title compound was prepared by the procedures described in Example 32 substituting 2,4-dichlorothiophenol with 4-bromothiophenol. Light-yellow powder; ¹H NMR (d⁶-DMSO, 300 MHz) δ 2.04 (s, 3H), 3.47 (br m, 4H), 3.52 (br m, 1H), 3.60 (br m, 1H), 3.68 (br m, 1H), 3.74 (br m, 1H), 6.90 (d, *J* = 8.7 Hz, 1H), 7.43 (d, *J* = 15.0 Hz, 1H), 7.54 (d, *J* = 15.0 Hz, 1H), 7.58 (d, *J* = 9.0 Hz, 2H), 7.78 (d, *J* = 9.0 Hz, 2H),
10 7.92 (dd, *J* = 2.1, 9.0 Hz, 1H), 8.65 (d, *J* = 2.1 Hz, 1H). MS (DCI/NH₃) (M+H)⁺ at *m/z* 507, 509.

Example 36

(4-Methylphenyl)[2-nitro-4-(*E*-(4-acetyl)piperazin-1-yl)carbonyl]ethenyl]phenyl]

sulfide

15 The title compound was prepared by the procedures described in Example 32 substituting 2,4-dichlorothiophenol with *p*-thiocresol. Light-yellow powder; ¹H NMR (d⁶-DMSO, 300 MHz) δ 2.04 (s, 3H), 2.39 (s, 3H), 3.47 (br m, 4H), 3.52 (br m, 1H), 3.60 (br m, 1H), 3.68 (br m, 1H), 6.89 (d, *J* = 8.7 Hz, 1H), 7.20 (d, *J* = 8.1 Hz, 1H),
20 7.39 (d, *J* = 8.4 Hz, 2H), 7.40 (d, *J* = 15.0 Hz, 1H), 7.53 (d, *J* = 15.0 Hz, 1H), 7.54 (d, *J* = 8.4 Hz, 2H), 7.89 (dd, *J* = 2.1, 8.7 Hz, 1H), 8.64 (d, *J* = 2.1 Hz, 1H). MS (DCI/NH₃) (M+NH₄)⁺ at *m/z* 443.

Example 37

(2,4-Dichlorophenyl)[2-nitro-4-(E-((4-(*tert*-butoxycarbonyl)piperazin-1-yl)carbonyl)ethenyl)phenyl] sulfide

5 The title compound was prepared by the procedures described in Example 32 substituting 1-acetylpiperazine with *tert*-butyl piperazine carboxylate. Light-yellow powder; ¹H NMR (d⁶-DMSO, 300 MHz) δ 1.42 (s, 9H), 3.36 (overlapping m, 4H), 3.55 (br m, 2H), 3.70 (br m, 2H), 6.83 (d, *J* = 8.7 Hz, 1H), 7.42 (d, *J* = 15.6 Hz, 1H), 7.54 (d, *J* = 15.6 Hz, 1H), 7.63 (dd, *J* = 2.4, 8.4 Hz, 1H), 7.83 (d, *J* = 8.7 Hz, 1H), 7.92 (dd, *J* = 2.4, 8.7 Hz, 1H), 7.96 (d, *J* = 2.7 Hz, 1H), 8.68 (d, *J* = 2.4 Hz, 1H). MS (APCI) (M+H)⁺ at *m/z* 538, 540, 542.

Example 38

(2,4-Dichlorophenyl)[2-nitro-4-(E-((4-(2-furoylcarbonyl)piperazin-1-yl)carbonyl)ethenyl)phenyl] sulfide

15

Example 38A

(2,4-Dichlorophenyl)[2-nitro-4-(E-((piperazin-1-yl)carbonyl) ethenyl)phenyl] sulfide
Trifluoroacetic Acid Salt

20 The compound (100 mg, 0.186 mmol) from Example 37 was dissolved in 0.5 mL of neat trifluoroacetic acid (TFA). The mixture was stirred at room temperature

for 1 hour. The TFA was then removed under vacuum to give the title compound (105 mg) as a yellow solid.

Example 38B

5 (2,4-Dichlorophenyl)[2-nitro-4-(*E*-((4-(2-furoylcarbonyl)piperazin-1-yl)carbonyl)ethenyl)phenyl] sulfide

To a stirred solution of piperazine TFA salt (35 mg, 0.067 mmol) from Example 38A in 2.0 mL of CH₂Cl₂ was added Et₃N (23 μL, 0.17 mmol), 4-dimethylaminopyridine (DMAP) (1.0 mg, 0.0082 mmol), and furyl chloride (8.0 μL, 0.080 mmol). The mixture was then stirred at room temperature for 30 minutes before the solvent was removed. The crude product was purified with Gilson HPLC system, YMC C-18 column, 75x30 mm I.D., S-5 μM, 120 Å, and a flow rate of 25 mL/min, λ=214, 245 nm; mobile phase A, 0.05 M NH₄OAc, and B, CH₃CN; linear gradient 20-100% of B in 20 minutes to give the title compound (24 mg, 67%) as light-yellow powder; ¹H NMR (d⁶-DMSO, 300 MHz) δ 3.62-3.87 (br m, 8H), 6.66 (q, *J* = 2.1 Hz, 1H), 6.84 (d, *J* = 8.7 Hz, 1H), 7.04 (d, *J* = 3.3 Hz, 1H), 7.44 (d, *J* = 15.3 Hz, 1H), 7.56 (d, *J* = 15.3 Hz, 1H), 7.63 (dd, *J* = 2.4, 8.1 Hz, 1H), 7.83 (d, *J* = 8.4 Hz, 1H), 7.87 (d, *J* = 2.1 Hz, 1H), 7.92 (dd, *J* = 2.1, 12.0 Hz, 1H), 7.96 (d, *J* = 2.1 Hz, 1H), 8.70 (d, *J* = 2.1 Hz, 1H). MS (APCI) (M+H)⁺ at *m/z* 532, 534, 536.

20

Example 39

(2,4-Dichlorophenyl)[2-nitro-4-(*E*-((4-(methanesulfonyl)piperazin-1-yl)carbonyl)

ethenyl)phenyl] sulfide

The title compound was prepared by the procedures described in Example 38B substituting furoyl chloride with methanesulfonyl chloride. Light-yellow powder; ¹H NMR (d⁶-DMSO, 300 MHz) δ 2.90 (s, 3H), 3.25 (br m, 4H), 3.68 (br m, 2H), 3.83 (br m, 2H), 6.84 (d, *J* = 9.0 Hz, 1H), 7.45 (d, *J* = 15.6 Hz, 1H), 7.56 (d, *J* = 15.6 Hz, 1H), 7.63 (dd, *J* = 2.4, 8.7 Hz, 1H), 7.83 (d, *J* = 9.0 Hz, 1H), 7.93 (dd, *J* = 2.1, 9.0 Hz, 1H), 7.95 (d, *J* = 2.7 Hz, 1H), 8.70 (d, *J* = 2.1 Hz, 1H). MS (ESI) (M+H)⁺ at *m/z* 516, 518, 520.

Example 40(2,4-Dichlorophenyl)[2-nitro-4-(*E*-(4-(diethylaminocarbonylmethyl)piperazin-1-yl)carbonyl)ethenyl)phenyl] sulfide

The title compound was prepared by the procedures described in Example 38B substituting furoyl chloride with 2-chloro-*N,N*-diethylacetamide. Light-yellow powder; ¹H NMR (d⁶-DMSO, 300 MHz) δ 1.01 (t, *J* = 7.2 Hz, 3H), 1.13 (t, *J* = 7.2 Hz, 3H), 2.46 (br m, 4H), 3.16 (s, 2H), 3.24 (q, *J* = 7.2 Hz, 2H), 3.37 (q, *J* = 7.2 Hz, 2H), 3.56 (br m, 2H), 3.69 (br m, 2H), 6.83 (d, *J* = 9.0 Hz, 1H), 7.46 (d, *J* = 15.3 Hz, 1H), 7.52 (d, *J* = 15.3 Hz, 1H), 7.62 (dd, *J* = 2.4, 8.7 Hz, 1H), 7.82 (d, *J* = 9.0 Hz, 1H), 7.92 (dd, *J* = 2.1, 9.0 Hz, 1H), 7.95 (d, *J* = 2.7 Hz, 1H), 8.67 (d, *J* = 2.1 Hz, 1H). MS (ESI) (M+NH₄)⁺ at *m/z* 573, 575, 577.

Example 41

(2,4-Dichlorophenyl)[2-nitro-4-(E-((4-(diethylaminocarbonyl)piperazin-1-yl)carbonyl)ethenyl)phenyl] sulfide

The title compound was prepared by the procedures described in Example 38B substituting furoyl chloride with *N,N*-diethylcarbonyl chloride. Light-yellow powder;

- 5 ¹H NMR (d⁶-DMSO, 300 MHz) δ 1.06 (t, *J* = 6.9 Hz, 6H), 3.12 (br m, 4H), 3.15 (q, *J* = 6.9 Hz, 4H), 3.58 (br m, 2H), 3.72 (br m, 2H), 6.83 (d, *J* = 8.7 Hz, 1H), 7.42 (d, *J* = 15.6 Hz, 1H), 7.53 (d, *J* = 15.6 Hz, 1H), 7.63 (dd, *J* = 2.7, 9.0 Hz, 1H), 7.82 (d, *J* = 8.7 Hz, 1H), 7.92 (dd, *J* = 2.4, 8.7 Hz, 1H), 7.95 (d, *J* = 2.7 Hz, 1H), 8.68 (d, *J* = 2.1 Hz, 1H). MS (APCI) (M+H)⁺ at *m/z* 537, 539, 541.

10

Example 42

(2,4-Dichlorophenyl)[2-nitro-4-(E-((4-(*tert*-butoxycarbonylmethyl)piperazin-1-yl)carbonyl)ethenyl)phenyl] sulfide

The title compound was prepared by the procedures described in Example 38B

- 15 substituting CH₂Cl₂ with CH₃CN as solvent, and furoyl chloride with *tert*-butyl bromoacetate. Light-yellow powder; ¹H NMR (CDCl₃, 300 MHz) δ 1.47 (s, 9H), 2.70 (br m, 4H), 3.21 (s, 2H), 3.74 (br m, 2H), 3.82 (br m, 2H), 6.73 (d, *J* = 8.7 Hz, 1H), 6.92 (d, *J* = 15.0 Hz, 1H), 7.39 (dd, *J* = 2.4, 8.7 Hz, 1H), 7.47 (d, *J* = 8.7 Hz, 1H), 7.61 (d, *J* = 15.0 Hz, 1H), 7.62 (d, *J* = 2.4 Hz, 1H), 7.66 (d, *J* = 8.7 Hz, 1H), 8.43 (br d,
- 20 1H). MS (APCI) (M+H)⁺ at *m/z* 552, 554, 556.

Example 43

(2,4-Dichlorophenyl)[2-nitro-4-(*E*-((4-(carboxycarbonyl)piperazin-1-yl)carbonyl)
ethenyl)phenyl] sulfide

Example 43A

5 (2,4-Dichlorophenyl)[2-nitro-4-(*E*-((4-(carboxycarbonyl)piperazin-1-yl)carbonyl)
ethenyl)phenyl] sulfide

The title compound was prepared by the procedures described in Example 38B substituting furoyl chloride with ethyl oxalyl chloride.

10 Example 43B

(2,4-Dichlorophenyl)[2-nitro-4-(*E*-((4-(carboxycarbonyl)piperazin-1-yl)carbonyl)
ethenyl)phenyl] sulfide

To a stirred solution of the ethyl ester (40 mg, 0.074 mmol) from Example 43A in 2 mL of ethanol was added saturated LiOH (0.25 mL). The mixture was then stirred at room temperature for 2 hours. Water (2 mL) was then added to the reaction mixture, which was then acidified to pH = 2 with concentrated HCl. The precipitates were collected through filtration, washed with cold water, dried under vacuum to give the titled compound (30 mg, 79%) as light yellow solid. ¹H NMR (d⁶-DMSO, 300 MHz) δ 3.52 (br m, 4H), 3.62 (br m, 2H), 3.76 (br m, 2H), 6.84 (d, *J* = 9.0 Hz, 1H), 7.46 (d, *J* = 15.3 Hz, 1H), 7.56 (d, *J* = 15.3 Hz, 1H), 7.63 (dd, *J* = 2.7, 8.7 Hz, 1H), 7.83 (d, *J* = 9.0 Hz, 1H), 7.93 (d, *J* = 9.0 Hz, 1H), 7.96 (d, *J* = 2.7 Hz, 1H), 8.70 (br d, 1H). MS (APCI) (M-COO)⁺ at *m/z* 466, 468, 470.

Example 44

(2,4-Dichlorophenyl)[2-nitro-4-(E-((4-(carboxymethyl)piperazin-1-yl)carbonyl)ethenyl)phenyl] sulfide

5

The title compound was prepared by the procedures described in Example 38A substituting compound from Example 37 with compound from Example 42. Light-yellow powder; ¹H NMR (d⁶-DMSO, 300 MHz) δ 3.14 (s, 2H), 3.40 (overlapping br m, 4H), 3.44 (br m, 1H), 3.51 (br m, 1H), 3.57 (br m, 1H), 3.71 (br m, 1H), 6.82 (d, *J* = 8.7 Hz, 1H), 7.42 (d, *J* = 15.6 Hz, 1H), 7.52 (d, *J* = 15.6 Hz, 1H), 7.63 (dd, *J* = 2.4, 8.7 Hz, 1H), 7.83 (d, *J* = 8.7 Hz, 1H), 7.92 (dd, *J* = 2.4, 8.7 Hz, 1H), 7.96 (d, *J* = 2.4 Hz, 1H), 8.68 (d, *J* = 2.4 Hz, 1H). MS (APCI) (M+H)⁺ at *m/z* 496, 498, 500.

10

Example 45

(2-Methylphenyl)[2-nitro-4-(E-((4-acetyl)piperazin-1-yl)carbonyl)ethenyl]phenyl] sulfide

15

The title compound was prepared by the procedures described in Example 32 substituting 2,4-dichlorothiophenol with *o*-thiocresol. Light-yellow powder; ¹H NMR (d⁶-DMSO, 300 MHz) δ 2.03 (s, 3H), 2.29 (s, 3H), 3.47 (br m, 4H), 3.53 (br m, 1H), 3.60 (br m, 1H), 3.67 (br m, 1H), 3.83 (br m, 1H), 6.64 (d, *J* = 8.7 Hz, 1H), 7.40 (d, *J* = 15.0 Hz, 1H), 7.36-7.42 (m, 1H), 7.46-7.57 (m, 3H), 7.63 (d, *J* = 6.9 Hz, 1H), 7.89 (dd, *J* = 2.4, 9.0 Hz, 1H), 8.66 (d, *J* = 2.4 Hz, 1H). MS (APCI) (M+H)⁺ at *m/z* 426.

20

Example 46

(2-Chlorophenyl)[2-nitro-4-(E-((4-acetyl)piperazin-1-yl)carbonyl)ethenyl]phenyl]

sulfide

5 The title compound was prepared by the procedures described in Example 32 substituting 2,4-dichlorothiophenol with 2-chlorothiophenol. Light-yellow powder; ¹H NMR (d⁶-DMSO, 300 MHz) δ 2.04 (s, 3H), 3.47 (br m, 4H), 3.52 (br m, 1H), 3.60 (br m, 1H), 3.68 (br m, 1H), 3.73 (br m, 1H), 6.75 (d, *J* = 9.0 Hz, 1H), 7.43 (d, *J* = 15.3 Hz, 1H), 7.54 (d, *J* = 15.3 Hz, 1H), 7.55 (dd, *J* = 1.8, 8.1 Hz, 1H), 7.64 (t, *J* = 1.8, 8.1 Hz, 1H), 7.76 (d, *J* = 1.8, 8.1 Hz, 1H), 7.82 (d, *J* = 1.8, 8.1 Hz, 1H), 7.93 (dd, *J* = 2.4, 9.0 Hz, 1H), 8.68 (d, *J* = 2.4 Hz, 1H). MS (APCI) (M+H)⁺ at *m/z* 446, 448, 450.

Example 47

(2-Aminophenyl)[2-nitro-4-(E-((4-acetyl)piperazin-1-yl)carbonyl)ethenyl]phenyl]

sulfide

15 The title compound was prepared by the procedures described in Example 32 substituting 2,4-dichlorothiophenol with 2-aminothiophenol. Light-yellow powder; ¹H NMR (d⁶-DMSO, 300 MHz) δ 2.04 (s, 3H), 3.47 (br m, 4H), 3.52 (br m, 1H), 3.60 (br m, 1H), 3.68 (br m, 1H), 3.74 (br m, 1H), 5.58 (s, 2H), 6.65 (td, *J* = 1.5, 15.0 Hz, 1H), 6.72 (dd, *J* = 1.5, 8.7 Hz, 1H), 7.00 (dd, *J* = 1.8, 8.7 Hz, 1H), 7.27 (t, *J* = 1.5, 8.6 Hz, 1H), 7.36 (dd, *J* = 1.5, 8.7 Hz, 1H), 7.39 (d, *J* = 15.3 Hz, 1H), 7.53 (d, *J* = 15.3 Hz,

5 (2-Hydroxymethyl)phenyl][2-nitro-4-(*E*-((4-acetyl)piperazin-1-yl)carbonyl)
ethenyl)phenyl] sulfide

15

(2-Ethylphenyl)[2-nitro-4-(*E*-((4-acetyl)piperazin-1-yl)carbonyl)ethenyl]phenyl
sulfide

The title compound was prepared by the procedures described in Example 32 substituting 2,4-dichlorothiophenol with 2-ethylthiophenol. Light-yellow powder; ¹H NMR (d⁶-DMSO, 300 MHz) δ 1.01 (t, *J* = 7.65 Hz, 3H), 2.04 (s, 3H), 2.69 (q, *J* = 7.65 Hz, 2H), 3.47 (br m, 4H), 3.52 (br m, 1H), 3.59 (br m, 1H), 3.67 (br m, 1H), 3.73

(br m, 1H), 6.64 (d, $J = 8.7$ Hz, 1H), 7.38 (dd, $J = 2.4, 7.5$ Hz, 1H), 7.40 (d, $J = 15.6$ Hz, 1H), 7.50-7.61 (m, 3H), 7.53 (d, $J = 15.6$ Hz, 1H), 7.89 (dd, $J = 2.4, 8.7$ Hz, 1H), 8.64 (d, $J = 2.4$ Hz, 1H). MS (APCI) (M+Cl)⁻ at m/z 474, 476.

5

Example 50

(2-iso-Propylphenyl)[2-nitro-4-(E-((4-acetyl)piperazin-1-yl)carbonyl)
ethenyl]phenyl] sulfide

The title compound was prepared by the procedures described in Example 32 substituting 2,4-dichlorothiophenol with 2-isopropylthiophenol. Light-yellow powder;

- 10 ¹H NMR (d⁶-DMSO, 300 MHz) δ 1.05 (d, $J = 6.9$ Hz, 6H), 2.04 (s, 3H), 3.47 (br m, 4H), 3.52 (br m, 1H), 3.60 (br m, 1H), 3.67 (br m, 1H), 3.72 (br m, 1H), 6.64 (d, $J = 8.4$ Hz, 1H), 7.34-7.41 (m, 2H), 7.39 (d, $J = 15.3$ Hz, 1H), 7.52 (d, $J = 15.3$ Hz, 1H), 7.56-7.73 (m, 2H), 7.90 (dd, $J = 2.1, 8.7$ Hz, 1H), 8.64 (d, $J = 2.1$ Hz, 1H). MS (APCI) (M+NH₄)⁺ at m/z 471. Analysis calculated for C₂₄H₂₇N₃O₄ S₁·0.21H₂O: C, 63.03; H, 5.96; N, 9.13. Found: C, 63.03; H, 6.04; N, 9.19.
- 15

Example 51

(2-tert-Butylphenyl)[2-nitro-4-(E-((4-acetyl)piperazin-1-yl)carbonyl)
ethenyl]phenyl] sulfide

- 20 The title compound was prepared by the procedures described in Example 32 substituting 2,4-dichlorothiophenol with 2-tert-butylthiophenol. Light-yellow powder;
- ¹H NMR (d⁶-DMSO, 300 MHz) δ 1.46 (s, 9H), 2.04 (s, 3H), 3.47 (br m, 4H), 3.52 (br

m, 1H), 3.60 (br m, 1H), 3.67 (br m, 1H), 3.73 (br m, 1H), 6.68 (d, $J = 8.7$ Hz, 1H), 7.35 (t, $J = 7.5$ Hz, 1H), 7.39 (d, $J = 15.3$ Hz, 1H), 7.45-7.57 (m, 2H), 7.50 (d, $J = 15.3$ Hz, 1H), 7.65 (d, $J = 8.1$ Hz, 1H), 7.88 (dd, $J = 2.4, 8.7$ Hz, 1H), 8.64 (d, $J = 2.4$ Hz, 1H). MS (APCI) $(M+NH_4)^+$ at m/z 485.

5

Example 52

(2-Chlorophenyl)[2-chloro-4-(*E*-(4-acetyl)piperazin-1-yl)carbonyl)]

2-propenyl)phenyl] sulfide

10

Example 52A

3'-Chloro-4'-[(2-chlorophenyl)thio]acetophenone

The title compound was prepared by the procedures described in Example 1A substituting 2,4-dichlorothiophenol with 2-chlorothiophenol, and 2-chlorobenzaldehyde with 4'-fluoro-3'-chloroacetophenone.

15

Example 52B

(2-Chlorophenyl)[2-chloro-4-(*E*-(1-ethoxycarbonyl) 2-propenyl)phenyl] sulfide

To a stirred suspension of NaH (60% in mineral oil, 121 mg, 3.03 mmol) in 20 mL of anhydrous THF under nitrogen atmosphere was added triethyl phosphonoacetate dropwise. After 20 minutes, the acetophenone (600 mg, 2.02 mmol) from Example 52A in THF (5 mL) was added in one portion. The resulting clear solution was then stirred at room temperature for 7 hours. Reaction was then stopped, most of the solvent was evaporated, and the residue was partitioned between

20

001320-56214560

EtOAc (2x20 mL) and water. The combined organic layer was washed with water and brine, dried over Na_2SO_4 , concentrated in vacuo. The crude product was purified using silica gel flash column chromatography eluting with 5-10% Et_2O in hexanes to give the (*E*)-isomer of the cinnamate (500 mg, 68%) as a white solid.

5

Example 52C

(2-Chlorophenyl)[2-chloro-4-(*E*-(1-carboxy) 2-propenyl)phenyl] sulfide

A mixture of the cinnamate (500 mg, 1.37 mmol) from Example 52B in 5 mL of EtOH/THF (4:1) was stirred with sat. LiOH solution (0.50 mL) at 50 °C for 2
10 hours. The mixture was then acidified with 3N HCl and extracted with CH_2Cl_2 (3x10 mL). The combined organic layer was dried over MgSO_4 , concentrated under reduced pressure to give the titled compound (450mg, 97%) as a white solid.

Example 52D

(2-Chlorophenyl)[2-chloro-4-(*E*-((4-acetylpiperazin-1-yl)carbonyl))

2-propenyl)phenyl] sulfide

The title compound was prepared using the cinnamic acid from Example 52C by the procedures described in Example 1C substituting 6-amino-1-hexanol with 1-acetylpiperazine. White solid; ^1H NMR (CDCl_3 , 300 MHz) δ 2.10-2.20 (m, 3H), 2.25
20 (s, 3H), 3.40-3.80 (m, 8H), 6.28 (s, 1H), 7.00 (d, $J = 8.7$ Hz, 1H), 7.19-7.36 (m, 4H), 7.46-7.56 (m, 2H). MS (APCI) ($\text{M}+\text{NH}_4$) $^+$ at m/z 466, 468, 470.

Example 53

(2-(1-Morpholinylmethyl)phenyl)[2-chloro-4-(E-((1-morpholinyl)carbonyl)
ethenyl) phenyl] sulfide

5

Example 53A

(2-(1-Bromomethyl)phenyl)[2-chloro-4-(E-((1-morpholinyl)carbonyl)
ethenyl) phenyl] sulfide

00544795.033400
00100.562450

To a stirred solution of benzyl alcohol (195 mg, 0.32 mmol) from Example 11
in 2.0 mL of anhydrous DMF was added LiBr (48 mg, 0.35 mmol). The mixture was
10 then cooled in an ice-water bath, and PBr₃ (60 μ L, 0.40 mmol) was dropped in slowly.
The ice bath was then removed and the mixture was stirred at room temperature for 1
hour. Water was then added, the mixture was then partitioned between EtOAc and
aqueous NaHCO₃. The aqueous layer was extracted with EtOAc once. The combined
organic layer was washed with water and brine, dried over Na₂SO₄, concentrated on a
15 rotavap. The crude bromide (230mg) was used directly for the alkylation without
purification.

Example 53B

(2-(1-Morpholinylmethyl)phenyl)[2-chloro-4-(E-((1-morpholinyl)carbonyl)
ethenyl) phenyl] sulfide

20

To a stirred solution of morpholine (10 μ L, 0.11 mmol) in 0.5 mL of CH₃CN
was added Hunig's base (23.7 μ L, 0.14 mmol), followed by the bromide (40 mg,

0.091 mmol). The mixture was then stirred at room temperature for 2 hours. Solvent was then removed and the crude product was purified with Gilson Preparative HPLC as described in Example 38B to give the titled compound as a white solid. ¹H NMR (d⁶-DMSO, 300 MHz) δ 2.33 (br t, 4H), 3.45 (br t, 4H), 3.50-3.65 (m, 6H), 3.56 (s, 2H), 3.65-3.80 (br m, 2H), 6.74 (d, *J* = 8.7 Hz, 1H), 7.30 (d, *J* = 15.3 Hz, 1H), 7.35-7.41 (m, 2H), 7.43 (d, *J* = 15.3 Hz, 1H), 7.46 (td, *J* = 2.4, 8.1 Hz, 1H), 7.52 (dd, *J* = 2.1, 8.7 Hz, 1H), 7.56 (d, *J* = 8.1 Hz, 1H), 8.02 (d, *J* = 2.1 Hz, 1H). MS (DCI/NH₃) (M+H)⁺ at *m/z* 459, 461.

10

Example 54

(2-(4-(1,3-Benzodioxolyl-5-methyl)piperazin-1-ylmethyl)phenyl)[2-chloro-4-(*E*-(1-morpholinyl)carbonyl)ethenyl]phenyl] sulfide

The title compound was prepared by the procedures described in Example 53B substituting morpholine with 1-piperonylpiperazine. White solid; ¹H NMR (d⁶-DMSO, 300 MHz) δ 2.13-2.40 (br m, 8H), 3.28 (s, 2H), 3.49-3.64 (br m, 6H), 3.54 (s, 2H), 3.70 (br m, 2H), 5.97 (s, 2H), 6.69 (dd, *J* = 1.8, 8.1 Hz, 1H), 6.74 (d, *J* = 8.7 Hz, 1H), 6.79 (d, *J* = 1.8 Hz, 1H), 6.81 (d, *J* = 8.1 Hz, 1H), 7.39 (d, *J* = 15.3 Hz, 1H), 7.33-7.38 (m, 2H), 7.38-7.50 (m, 2H), 7.43 (d, *J* = 15.3 Hz, 1H), 7.53 (d, *J* = 8.4 Hz, 1H), 8.00 (d, *J* = 2.1 Hz, 1H). MS (DCI/NH₃) (M+H)⁺ at *m/z* 592, 594.

20

Example 55

(2-(4-(*iso*-Propylaminocarbonylmethyl)piperazin-1-ylmethyl)phenyl)[2-chloro-4-(*E*-
((1-
morpholinyl)carbonyl) ethenyl)phenyl] sulfide

The title compound was prepared by the procedures described in Example 53B

- 5 substituting morpholine with *N*-isopropyl-1-piperazineacetamide. White solid; ¹H
 NMR (d⁶-DMSO, 300 MHz) δ 1.04 (d, *J* = 6.3 Hz, 6H), 2.20-2.42 (br m, 8H), 2.78 (s,
 2H), 3.47-3.64 (br m, 6H), 3.56 (s, 2H), 3.64-3.76 (br m, 2H), 3.85 (qd, *J* = 6.3, 8.1
 Hz, 1H), 6.73 (d, *J* = 8.7 Hz, 1H), 7.29 (d, *J* = 15.6 Hz, 1H), 7.31-7.39 (m, 2H), 7.43
 (d, *J* = 15.6 Hz, 1H), 7.45 (td, *J* = 2.7, 6.3 Hz, 1H), 7.50 (dd, *J* = 2.1, 8.7 Hz, 1H),
 10 7.55 (d, *J* = 7.8 Hz, 1H), 8.00 (d, *J* = 2.1 Hz, 1H). MS (DCI/NH₃) (M+H)⁺ at *m/z* 557,
 559.

Example 56

(2-((*N*-Ethoxycarbonylmethyl)-*N*-methyl)aminomethyl)phenyl)[2-chloro-4-(*E*-((1-
morpholinyl)carbonyl) ethenyl)phenyl] sulfide

- 15 The title compound was prepared by the procedures described in Example 53B
 substituting morpholine with ethyl sarcosinate hydrochloride. White solid; ¹H NMR
 (d⁶-DMSO, 300 MHz) δ 1.16 (t, *J* = 7.2 Hz, 3H), 2.27 (s, 2H), 3.30 (s, 2H), 3.51-3.66
 (br m, 6H), 3.66-3.75 (br m, 2H), 3.78 (s, 2H), 4.05 (q, *J* = 7.2 Hz, 2H), 6.75 (d, *J* =
 20 8.7 Hz, 1H), 7.30 (d, *J* = 15.3 Hz, 1H), 7.33-7.38 (m, 2H), 7.42-7.50 (m, 2H), 7.43 (d,
J = 15.3 Hz, 1H), 7.53 (dd, *J* = 2.1, 8.7 Hz, 1H), 7.60 (d, *J* = 7.8 Hz, 1H), 8.02 (d, *J* =
 2.1 Hz, 1H). MS (DCI/NH₃) (M+H)⁺ at *m/z* 489, 491.

001100.56214560

Example 57(2-Formylphenyl)[2-chloro-4-(*E*-((1-morpholinyl)carbonyl)ethenyl)phenyl] sulfide

To a stirred solution of the alcohol (368 mg, 0.94 mmol) from Example 11 in 5 mL of anhydrous acetonitrile was added activated 4Å molecular sieves, TPAP (3.3 mg, 0.0094 mmol), and NMO (110 mg, 1.03 mmol). The mixture was then stirred at room temperature for 3 hours. The reaction mixture was then quenched with dimethyl sulfide (100 µL). The crude product was filtered through celite, washed with acetonitrile, condensed in vacuo. The titled compound was purified by silica gel column chromatography to give a white solid (216 mg, 59 %). ¹H NMR (d⁶-DMSO, 300 MHz) δ 3.60 (br m, 6H), 3.73 (br m, 2H), 7.00 (d, *J* = 8.4 Hz, 1H), 7.40 (d, *J* = 15.3 Hz, 1H), 7.42 (d, *J* = 8.4 Hz, 1H), 7.51 (d, *J* = 15.3 Hz, 1H), 7.52 (td, *J* = 1.8, 8.1 Hz, 1H), 7.61 (td, *J* = 1.8, 8.1 Hz, 1H), 7.71 (dd, *J* = 2.1, 8.4 Hz, 1H), 8.02 (dd, *J* = 2.1, 8.4 Hz, 1H), 8.14 (d, *J* = 2.1 Hz, 1H). MS (DCI/NH₃) (M+H)⁺ at *m/z* 388, 390.

Example 58(2-(4-Formylpiperazin-1-ylmethyl)phenyl)[2-chloro-4-(*E*-((1-morpholinyl)carbonyl)ethenyl)phenyl] sulfide

The title compound was prepared by the procedures described in Example 53B substituting morpholine with 1-formyl piperazine. White solid; ¹H NMR (d⁶-DMSO, 300 MHz) δ 2.20-2.32 (m, 6H), 2.74 (br m, 2H), 3.48 (s, 2H), 3.59 (m, 6H), 3.70 (br m, 2H), 6.74 (d, *J* = 8.7 Hz, 1H), 7.29 (d, *J* = 15.6 Hz, 1H), 7.35-7.41 (m, 2H), 7.42

(d, $J = 15.6$ Hz, 1H), 7.45-7.52 (m, 3H), 7.98 (d, $J = 2.1$, 1H). MS (DCI/NH₃) (M+H)⁺ at m/z 486, 488.

Example 59

5 (2-(*E*-((1-Morpholinyl)carbonyl)ethenyl)phenyl)[2-chloro-4-(*E*-((1-morpholinyl)carbonyl)ethenyl)phenyl] sulfide

A mixture of bromide (80 mg, 0.18 mmol) from Example 12, acryloylmorpholine (33 mg, 0.23 mmol), Pd(OAc)₂ (2.0 mg, 0.009 mmol), P(*o*-tolyl)₃ (17 mg, 0.056 mmol), Et₃N (39 μ L, 0.27 mmol), and anhydrous DMF (1.0 mL) in a pressure tube was flushed with nitrogen for 5 minutes before it capped and heated at 110 °C over night. TLC indicated almost complete consumption of the starting bromide. The reaction mixture was then allowed to cool down to room temperature, partitioned between EtOAc and water. The aqueous layer was extracted once with EtOAc. The combined organic layer was washed with water and brine, dried over Na₂SO₄, condensed under reduced pressure. The crude product was purified with Gilson Preparative HPLC as described in Example 38B to give the titled compound as a light-brown solid (35 mg, 39%). ¹H NMR (d⁶-DMSO, 300 MHz) δ 3.43-3.88 (m, 16H), 6.58 (d, $J = 8.7$ Hz, 1H), 7.30 (d, $J = 15.3$ Hz, 2H), 7.43 (d, $J = 15.3$ Hz, 1H), 7.47-7.64 (m, 4H), 7.86 (d, $J = 15.3$ Hz, 1H), 8.06 (d, $J = 2.1$ Hz, 1H), 8.14 (d, $J = 7.5$ Hz, 1H). MS (DCI/NH₃) (M+NH₄)⁺ at m/z 516, 518. Analysis calculated for

$C_{26}H_{27}N_2O_4ClS_1 \cdot 0.46H_2O$: C, 61.56; H, 5.55; N, 5.21. Found: C, 61.56; H, 5.50; N, 5.43.

Example 60

5 (2-Formylphenyl)[2-nitro-4-(E-((4-acetyl)piperazin-1-yl)carbonyl)ethenyl)phenyl]
sulfide

The title compound was prepared by the procedures described in Example 57 substituting compound from Example 11 with compound from Example 48. Yellow solid; ¹H NMR (d⁶-DMSO, 300 MHz) δ 2.04 (s, 3H), 3.47 (br m, 4H), 3.52 (br m, 1H), 3.60 (br m, 1H), 3.68 (br m, 1H), 3.74 (br m, 1H), 6.85 (d, *J* = 8.4 Hz, 1H), 7.44 (d, *J* = 15.6 Hz, 1H), 7.55 (d, *J* = 15.6 Hz, 1H), 7.61 (d, *J* = 7.5 Hz, 1H), 7.73 (t, *J* = 7.5 Hz, 1H), 7.80 (td, *J* = 2.4, 7.5 Hz, 1H), 7.92 (dd, *J* = 2.1, 9.0 Hz, 1H), 8.04 (dd, *J* = 2.4, 7.5 Hz, 1H), 8.66 (d, *J* = 2.1 Hz, 1H), 10.29 (s, 1H). MS (APCI) (M+Cl)⁺ at *m/z* 474, 476.

Example 61

(2-Formylphenyl)[2-chloro-4-(*E*-((1-morpholinyl)carbonyl)ethenyl)phenyl] sulfide.

N,N-dimethyl hydrazone

A mixture of the aldehyde (20 mg, 0.052 mmol) from Example 57, 1,1-dimethyl hydrazine (3.9 μ L, 0.052 mmol) in 0.5 mL of EtOH with a tiny amount of AcOH was stirred at room temperature over night. The solvent was then removed and the product was purified by preparative TLC to give the titled compound (20 mg,

90%) as a white solid. ^1H NMR (CDCl_3 , 300 MHz) δ 2.91 (s, 6H), 3.55-3.82 (br m, 8H), 6.64 (d, J = 8.7 Hz, 1H), 6.76 (d, J = 15.3 Hz, 1H), 7.05 (dd, J = 1.8, 8.7 Hz, 1H), 7.26 (td, J = 1.8, 7.8 Hz, 1H), 7.43 (t, J = 7.8 Hz, 1H), 7.47-7.57 (m, 2H), 7.54 (m, 2H), 8.04 (dd, J = 1.8, 8.7 Hz, 1H). MS (DCI/NH_3) ($\text{M}+\text{H}$) $^+$ at m/z 430, 432, 434,

5 436.

Example 62

(2-((3-(1-Morpholinyl)propyl)-1-amino)phenyl)[2-chloro-4-(*E*-(1-morpholinyl)carbonyl) ethenyl]phenyl sulfide

10 A mixture of bromide (60 mg, 0.14 mmol) from Example 12, aminopropylmorpholine (24 μL , 0.17 mmol), $\text{Pd}_2(\text{dba})_3$ (1.2 mg, 0.0013 mmol), BINAP (2.5 mg, 0.004 mmol), NaOt-Bu (19 mg, 0.20 mmol), 18-crown-6 (50 mg, 0.20 mmol), and anhydrous toluene (1 mL) in a pressure tube was flushed with nitrogen for 3 minutes before it was capped and heated at 80 $^\circ\text{C}$ over night. The

15 reaction was then stopped, and allowed to cool down to room temperature. The reaction mixture was partitioned between EtOAc and water, and the aqueous layer was extracted once with EtOAc. The combined organic layer was then washed with water and brine, dried over Na_2SO_4 , condensed under reduced pressure. The crude product was purified with Gilson Preparative HPLC as described in Example 38B to

20 give the titled compound as a light-brown oil (30 mg, 44%). ^1H NMR (d^6 -DMSO, 300 MHz) δ 1.62 (quintet, J = 6.5 Hz, 2H), 2.15-2.26 (m, 8H), 3.17 (q, J = 6.5 Hz, 2H), 3.22-3.76 (m, 12 H), 3.50 (t, J = 6.5 Hz, 2H), 5.72 (t, J = 5.7 Hz, 1H), 6.47 (d, J

= 8.7 Hz, 1H), 6.68 (t, J = 7.2 Hz, 1H), 6.81 (d, J = 8.4 Hz, 1H), 7.26 (d, J = 15.6 Hz, 1H), 7.35-7.42 (m, 2H), 7.43 (d, J = 15.6 Hz, 1H), 7.44 (d, J = 8.4 Hz, 1H), 7.49 (d, J = 8.4 Hz, 1H), 8.00 (d, J = 2.1 Hz, 1H). MS (APCI) (M+H)⁺ at m/z 502, 504.

5

Example 63

(2,4-Dichlorophenyl)[2-bromo-4-(*E*-(3-(1-pyrrolidin-2-yl)propylamino)carbonyl)
ethenyl)phenyl] sulfide

Example 63A

10 (2,4-Dichlorophenyl)[2-amino-4-(*E*-(3-(1-pyrrolidin-2-yl)propylamino)carbonyl)
ethenyl)phenyl] sulfide

A mixture of nitro compound (780 mg, 1.58 mmol) from Example 33, SnCl₂ (1.50 g, 7.91 mmol) in 25 mL of anhydrous EtOH was refluxed under nitrogen atmosphere for 90 minutes. The reaction was then allowed to cool down to room
15 temperature, quenched with sat. NaHCO₃, extracted with EtOAc (2x50 mL). The combined organic layer was washed with water and brine, dried over Na₂SO₄, condensed in vacuo to give the crude aniline as yellowish brown solid, which was converted to the bromide without purification.

20

Example 63B

(2,4-Dichlorophenyl)[2-bromo-4-(*E*-(3-(1-pyrrolidin-2-yl)propylamino)carbonyl)
ethenyl)phenyl] sulfide

To a stirred solution of *t*-butyl nitrite (57 μ L, 0.48 mmol), CuBr₂ (87 mg, 0.39 mmol) in 2.0 mL of CH₃CN at room temperature was added a solution of aniline from Example 63A (150 mg, 0.323 mmol) in 1.0 mL of CH₃CN. The dark green solution was then heated at 65 °C under nitrogen atmosphere for 90 minutes. The reaction mixture was then allowed to cool down to room temperature, partitioned between EtOAc and 3N HCl. The organic layer was then washed with brine, dried over Na₂SO₄, condensed in vacuo. The crude product was then purified with Gilson Preparative HPLC as described in Example 38B to give the titled compound as a light-brown solid (50 mg, 29%). Colorless oil; ¹H NMR (d⁶-DMSO, 300 MHz) δ 1.63 (quintet, *J* = 7.2 Hz, 2H), 1.91 (quintet, *J* = 8.4 Hz, 2H), 2.22 (t, *J* = 8.4 Hz, 2H), 3.09-3.47 (m, 6H), 6.67 (d, *J* = 15.3 Hz, 1H), 7.07 (d, *J* = 8.4 Hz, 1H), 7.32 (d, *J* = 8.7 Hz, 1H), 7.38 (d, *J* = 15.3 Hz, 1H), 7.50 (dd, *J* = 2.4, 8.7 Hz, 1H), 7.57 (dd, *J* = 2.1, 8.4 Hz, 1H), 7.86 (d, *J* = 2.4 Hz, 1H), 7.96 (d, *J* = 2.1 Hz, 1H), 8.13 (t, *J* = 6.0 Hz, 1H). MS (ESI) (M+H)⁺ at *m/z* 527, 529, 531, 533.

Example 64

(2,4-Dichlorophenyl)[2-formyl-4-(*E*-((1-morpholinyl)carbonyl)ethenyl)phenyl]
sulfide

Example 64A

[1-Fluoro-2-formyl-4-(*E*-((1-morpholinyl)carbonyl)ethenyl) benzene

The title compound was prepared by the procedures described in Example 59 substituting the bromide from Example 12 with 2-fluoro-5-bromobenzaldehyde.

Example 64B

5 (2,4-Dichlorophenyl)[2-formyl-4-(*E*-((1-morpholinyl)carbonyl)ethenyl)phenyl]
sulfide

The title compound was prepared by the procedures described in Example 32 substituting 4-chloro-3-nitro-cinnamide with the compound from Example 64A.

White solid; ¹H NMR (d⁶-DMSO, 300 MHz) δ 3.60 (br m, 6H), 3.71 (br m, 2H), 6.82
 10 (d, *J* = 8.7 Hz, 1H), 7.35 (d, *J* = 15.6 Hz, 1H), 7.54 (d, *J* = 15.6 Hz, 1H), 7.55 (dd, *J* =
 2.4, 8.7 Hz, 1H), 7.61 (d, *J* = 8.7 Hz, 1H), 7.86 (dd, *J* = 2.4, 8.4 Hz, 1H), 7.91 (d, *J* =
 2.4 Hz, 1H), 8.41 (d, *J* = 2.1 Hz, 1H), 10.19 (s, 1H). MS (DCl/NH₃) (M+H)⁺ at *m/z*
 422. 424. 426. 428.

15 Example 65

(2-Chloro-6-formylphenyl)[2-chloro-4-(*E*-((4-acetyl)piperazin-1-yl)carbonyl
ethenyl) phenyl] sulfide

Example 65A

20 (2-Carbomethoxyethyl)[2-chloro-4-(E-((4-acetyl)piperazin-1-yl)carbonyl)
ethenyl] phenyl] sulfide

The title compound was prepared by the procedures described in Example 1 substituting 2,4-dichlorothiophenol with methyl 3-mercaptopropionate, and 6-amino-1-hexanol with 1-acetyl piperazine.

Example 65B

5 (2-Chloro-6-formylphenyl)[2-chloro-4-(*E*-((4-acetyl)piperazin-1-yl)carbonyl)ethenyl] phenyl] sulfide

To a stirred solution of the compound (105 mg, 0.26 mmol) from Example 65A in 2 mL of THF under nitrogen atmosphere at 0 °C was added *t*-BuOK solution (1.0M, 281 µL, 0.29 mmol). Light orange precipitates appeared immediately. After
10 completion of the addition, the reaction mixture was stirred at room temperature for 1 hour before the solvent was removed on a rotavap under reduced pressure.

The yellow thiolate thus obtained was dissolved in 0.5 mL of DMF, and 2,3-dichlorobenzaldehyde was then added. The mixture was then heated at 80 °C under nitrogen for 2 hours. Reaction was then stopped and the solvent was removed under
15 vacuum. The crude product was purified with Gilson Preparative HPLC as described in Example 38B to give the titled compound as a white solid (25 mg, 21%). ¹H NMR (CDCl₃, 300 MHz) δ 2.05 (s, 3H), 3.48-3.58 (m, 2H), 3.58-3.84 (m, 6H), 6.53 (d, *J* = 8.7 Hz, 1H), 6.80 (d, *J* = 15.3 Hz, 1H), 7.19 (dd, *J* = 1.8, 8.7 Hz, 1H), 7.51-7.62 (m, 2H), 7.60 (d, *J* = 15.3 Hz, 1H), 7.84 (dd, *J* = 1.8, 8.4 Hz, 1H), 7.99 (dd, *J* = 1.8, 8.4
20 Hz, 1H). MS (APCI) (M+NH₄)⁺ at *m/z* 480, 482, 484.

Example 66

(2-Cyanophenyl)[2-chloro-4-(*E*-((4-acetyl)piperazin-1-yl)carbonyl)
ethenyl) phenyl] sulfide

The title compound was prepared by the procedures described in Example 65B substituting 2,3-dichlorobenzaldehyde with 2-fluorobenzonitrile, giving a white solid.

- 5 ¹H NMR (CDCl₃, 300 MHz) δ 2.15 (s, 3H), 3.48-3.57 (m, 2H), 3.59-3.84 (m, 6H), 6.86 (d, J = 15.6 Hz, 1H), 7.12 (d, J = 8.4 Hz, 1H), 7.32 (d, J = 8.4 Hz, 1H), 7.41 (d, J = 6.6 Hz, 1H), 7.46 (dd, J = 1.8, 8.4 Hz, 1H), 7.55 (dd, J = 1.8, 8.1 Hz, 1H), 7.61 (d, J = 15.6 Hz, 1H), 7.64 (d, J = 1.8 Hz, 1H), 7.75 (dd, J = 1.8, 8.4 Hz, 1H). MS (DCI/NH₃) (M+NH₄)⁺ at *m/z* 443.

10

Example 67

(2-Isopropylphenyl)[2-cyano-4-(*E*-((morpholin-1-yl)carbonyl)
ethenyl) phenyl] sulfide

15

Example 67A

(2-Isopropylphenyl)(4-bromo-2-cyanophenyl)sulfide

The title compound was prepared by the procedures described in Example 1A substituting 2,4-dichlorothiophenol with isopropylthiophenol, and 2-chlorobenzaldehyde with 2-fluorobenzonitrile.

20

Example 67B

(2-Isopropylphenyl)[2-cyano-4-(*E*-((morpholin-1-yl)carbonyl)

ethenyl) phenyl] sulfide

The title compound was prepared by the procedures described in Example 59 substituting the bromide from Example 12 with the bromide from Example 67A, giving a white solid. ¹H NMR (CDCl₃, 300 MHz) δ 1.19 (d, *J* = 6.9 Hz, 6H), 3.49 (septet, *J* = 6.9 Hz, 1H), 3.58-3.87 (m, 8H), 6.73 (d, *J* = 8.4 Hz, 1H), 6.83 (d, *J* = 15.6 Hz, 1H), 7.20-7.30 (m, 1H), 7.42 (dd, *J* = 2.4, 8.4 Hz, 1H), 7.46 (d, *J* = 3.0 Hz, 2H), 7.49 (dd, *J* = 1.8, 6.9 Hz, 1H), 7.57 (d, *J* = 15.6 Hz, 1H), 7.76 (d, *J* = 1.8 Hz, 1H). MS (APCI⁺) (M+H)⁺ at *m/z* 393.

10

Example 68(2-Bromophenyl)[2-nitro-4-(*E*-((4-acetyl)piperazin-1-yl)carbonyl)ethenyl) phenyl] sulfide

The title compound was prepared by the procedures described in Example 32B substituting 2,4-dichlorothiophenol with 2-bromothiophenol, providing a light-yellow solid; ¹H NMR (d⁶-DMSO, 300 MHz) δ 2.04 (s, 3H), 3.40-3.65 (m, 8H), 6.75 (d, *J* = 8.7 Hz, 1H), 7.42 (d, *J* = 15.6 Hz, 1H), 7.51 (dd, *J* = 2.1, 6.9 Hz, 1H), 7.54 (d, *J* = 15.6 Hz, 1H), 7.55 (t, *J* = 2.1 Hz, 1H), 7.59 (dd, *J* = 2.1, 6.9 Hz, 1H), 7.82 (dd, *J* = 2.4, 7.8 Hz, 1H), 7.92 (td, *J* = 2.4, 8.4 Hz, 1H), 8.67 (d, *J* = 2.4 Hz, 1H). MS (APCI⁺) (M+Cl)⁺ at *m/z* 524, 526, 528.

20

Example 69(2-(Pyrrolidin-1-yl)phenyl)[2-chloro-4-(*E*-((morpholin-1-yl)carbonyl)

ethenyl) phenyl] sulfide

- To a stirred solution of bromide (75 mg, 0.17 mmol) from Example 12 in toluene in a sealed tube was added sequentially pyrrolidine (18.4 mL, 0.22 mmol), $\text{Pd}_2(\text{dba})_3$ (3.0 mg, 0.0034mmol), BINAP (6.0 mg, 0.010mmol), followed by NaOr-Bu (26 mg, 0.27 mmol). The resulting mixture was then flushed with anhydrous N_2 for 2 min before it was capped and heated at 90 °C for 24 h. The reaction mixture was then allowed to cool down to room temperature and partitioned between ethyl acetate and brine. The organic layer was then dried with Na_2SO_4 , filtered, and concentrated in vacuo. The crude product was purified using Gilson Preparative HPLC as described in Example 38B to give the title compound (40 mg, 55% yield) as a white solid; ^1H NMR (CDCl_3 , 300 MHz) δ 1.83 (br s, 4H), 3.40 (br s, 4H), 3.56-3.80 (m, 8H), 6.57 (d, J = 8.4 Hz, 1H), 6.75 (d, J = 15.6 Hz, 1H), 6.81 (br t, J = 8.4 Hz, 1H), 6.90 (br s, 1H), 7.15 (dd, J = 2.1, 8.4 Hz, 1H), 7.18-7.27 (m, 1H), 7.32 (td, J = 1.8, 8.4 Hz, 1H), 7.42 (dd, J = 1.8, 7.8 Hz, 1H), 7.50 (d, J = 1.8 Hz, 1H), 7.55 (d, J = 15.6 Hz, 1H). MS (APCI $^+$) ($\text{M}+\text{H}$) $^+$ at m/z 429, 431.

Example 70(2-Methoxyphenyl)-[2-chloro-4(E-[(morpholin-1-yl)carbonyl]ethenyl)phenyl]sulfide

- The title compound was prepared according to the procedures of Example 1, giving a white solid, m.p. 162-164C. ^1H NMR (CDCl_3 , 300 MHz) δ 3.60-3.78 (m, 8H), 3.84 (s, 3H), 6.72 (d, J =9Hz, 1H), 6.78 (d, J =16Hz, 1H), 6.96-7.04 (m, 2H), 7.16 (dd, J =9Hz, 2Hz, 1H), 7.40-7.46 (, 2H), 7.55 (d, J =2H, 1H), 7.58 (d, J =16Hz, 1H).

Anal. Calcd. for $C_{20}H_{20}ClNO_3S$: C, 61.61; H, 5.17; N, 3.59. Found: C, 61.53, H, 5.22; N, 3.50.

Example 71

5 (2-Isopropylphenyl)[2-nitro-4-(*E*-((3-carbomethoxypiperazin-1-yl)carbonyl)ethenyl) phenyl] sulfide

Example 71A

1-*tert*-Butyloxycarbonyl-2-carbomethoxypiperazine

2-Carbomethoxypiperazine was treated with benzyl chloroformate (1.0 eq) in
10 aqueous $NaHCO_3$ to give 1-benzyloxycarbonyl-3-carbomethoxypiperazine. This
material was treated with di-*tert*-butyldicarbonate (1.1 eq) and triethylamine (1.0 eq)
in THF to produce 1-*tert*-butyloxycarbonyl-4-benzyloxycarbonyl-2-
carbomethoxypiperazine. Hydrogenation of this compound in methanol using 10%
Pd-C gives the title compound after filtration and solvent removal.

15 Example 71B

(2-Isopropylphenyl)[2-nitro-4-(*E*-((3-carbomethoxypiperazin-1-yl)carbonyl)ethenyl) phenyl] sulfide

A mixture of (2-isopropylphenyl)[2-nitro-4-*E*-(carboxyethenyl)phenyl]
sulfide (prepared according to the procedures of Example 32), the amine from
20 Example 71A (1.0 eq), 2-(1H-benzotriazol-1-yl)-1,1,3,3-tetramethyluronium
tetrafluoroborate (1.0 eq), and diisopropylethylamine (2.0 eq) in DMF was stirred at
ambient temperature for 4 hr. Ethyl acetate was added, and the mixture was washed

09541795-033100

sequentially with 1N HCl, bicarb, and brine. The resultant yellow solid was treated with 1:1 TFA/dichloromethane at ambient temperature to give the title compound as a yellow solid. ¹H NMR (DMSO-d₆, 300MHz) δ 1.15 (d, J = 6.6 Hz, 6H); 2.52-3.16 (br m, 4H); 3.25-3.47 (m, 1H); 3.60-3.65 (br d, 3H); 3.60, 3.66 (br s, br s, 3H); 6.61-6.67 (br m, 1H); 7.30-7.62 (m, 6H); 7.88-7.93 (br m, 1H); 8.58-8.65 (br m, 1H). MS (APCI) (M+H)⁺ at m/z 470. Anal calcd for C₂₄H₂₇N₃O₅: C, 61.39; H, 5.80; N, 8.95. Found: C, 61.51; H, 5.87; N, 8.68.

Example 72

10 (2-Methylphenyl)[2-nitro-4-(E-((3-carboxamido-4-carbobenzoxypiperazin-1-yl)carbonyl)ethenyl) phenyl] sulfide

Prepared according to the procedures of Example 71, giving a yellow solid. ¹H NMR (DMSO-d₆, 300MHz) δ 2.30 (s, 3H); 2.80-4.80 (br m, 7H); 5.05-5.15 (br m, 2H); 6.61-6.67 (br m, 1H); 7.02-7.64 (m, 13H); 7.80-7.90 (br m, 1H); 8.56-8.65 (br m, 1H). MS (APCI) (M+H)⁺ at m/z 561. Anal calcd for C₂₉H₂₈N₄S₁O₈·0.42CH₃COOCH₂CH₃: C, 61.66; H, 5.29; N, 9.38. Found: C, 61.41; H, 5.28; N, 9.53.

Example 73

20 (2-Isopropylphenyl)[2-nitro-4-(E-((2-carbomethoxy-4-tert-butoxycarbonyl)piperazin-1-yl)carbonyl)ethenyl) phenyl] sulfide

Prepared according to the procedures of Example 71, giving a yellow solid. ¹H NMR (DMSO-d₆, 300MHz) δ 1.13 (d, J = 6.6 Hz, 6H); 1.40, 1.41 (s, s, 9H); 2.72-3.08 (br m, 1H); 3.17-3.24 (m, 1H); 3.30-3.40 (m, 1H); 3.68 (br s, 3H); 3.79-4.51 (br m, 4H); 5.06, 5.36 (br s, br s, 1H); 6.61-6.67 (m, 1H); 7.30-7.62 (m, 6H); 7.85-7.93 (br m, 1H); 8.64-8.69 (br m, 1H). MS (APCI) (M+H)⁺ at m/z 570. Anal calcd for C₂₉H₃₃N₃S₁O₄·0.15C₆H₁₄: C, 61.66; H, 6.43; N, 7.21. Found: C, 61.69; H, 6.35; N, 7.02.

Example 74

10 (2-Isopropylphenyl)[2-nitro-4-(E-((2-carboxy-4-tert-butoxycarbonylpiperazin-1-yl)carbonyl)ethenyl) phenyl] sulfide

Prepared according to the procedures of Example 71, giving a yellow solid. ¹H NMR (CDCl₃, 300MHz) δ 1.14 (d, J = 6.6 Hz, 6H); 1.45 (s, 9H); 2.72-4.75 (br m, 6H); 3.38-3.49 (m, 1H); 5.78 (br s, 1H); 6.68, 6.72 (s, s, 1H); 6.88, 6.94 (br s, br s, 1H); 7.26-7.71 (m, 6H); 8.44 (br s, 1H). MS (APCI) (M-H)⁺ at m/z 554. Anal calcd for C₂₈H₃₃N₃S₁O₇: C, 60.53; H, 5.99; N, 7.56. Found: C, 60.42; H, 6.21; N, 7.31.

Example 75

20 (2-Isopropylphenyl)[2-trifluoromethyl-4-(E-((4-acetylpiperazin-1-yl)carbonyl)ethenyl) phenyl] sulfide

The title compound was prepared according to the procedures of Example 1. ¹H NMR (CDCl₃, 300 MHz) δ 7.78 (s, 1H), 7.62 (d, 1H, J = 15.5 Hz), 7.43-7.49 (m,

3H), 7.37 (d, 1H, $J = 8.1$ Hz), 7.23 (m, 1H), 6.85 (d, 1H, $J = 15.5$ Hz), 6.82 (d, 1H, $J = 8.5$ Hz), 3.63-3.77 (m, 6H), 3.45-3.55 (m, 3H), 2.14 (s, 3H), 1.17 (d, 6H, $J = 6.6$ Hz). MS (ESI) m/z 477, 499, 975, 953. Anal. Calcd for $C_{25}H_{27}F_3N_2O_2S \cdot 0.5$ EtOAc: C, 62.29; H, 6.00; N, 5.38. Found: C, 62.40; H, 6.21; N, 5.35.

5

Example 76

(2-Isopropylphenyl)[2-trifluoromethyl-4-(*E*-(morpholin-1-yl)carbonyl)ethenyl]phenyl] sulfide

The title compound was prepared according to the procedures of Example 1.

10 1H NMR ($CDCl_3$, 300 MHz) 7.78 (s, 1H), 7.62 (br, 1H), 7.33-7.48 (m, 3H), 7.22 (m, 1H), 6.85 (m, 1H), 6.80 (d, 1H, $J = 8.5$ Hz), 3.73 (br, 8H), 3.49 (dq, 1H, $J_1 = J_2 = 6.9$ Hz), 1.17 (d, 6H, $J = 7.1$ Hz). MS (ESI) m/z 436, 871, 893. Anal. Calcd for $C_{23}H_{24}F_3N_1O_2S$: C, 63.43; H, 5.55; N, 3.22. Found: C, 63.12; H, 5.81; N, 3.10.

15

Example 77

(2-Isopropylphenyl)[2-trifluoromethyl-4-(*E*-(3-(pyrrolidin-2-on-1-yl)prop-1-ylamino)carbonyl)ethenyl]phenyl]sulfide

The title compound was prepared according to the procedures of Example 1.

20 1H NMR ($CDCl_3$, 300 MHz) δ 7.77 (s, 1H), 7.52 (d, 1H, $J = 15.4$ Hz), 7.43-7.51 (m, 3H), 7.36 (d, 1H, $J = 8.8$ Hz), 7.22 (m, 1H), 7.10 (br, 1H), 6.80 (d, 1H, $J = 8.4$ Hz), 6.44 (d, 1H, $J = 15.4$ Hz), 3.49 (dq, 1H, $J_1 = J_2 = 6.9$ Hz), 3.40 (m, 4H), 3.31 (dd, 2H, $J_1 = 5.7$ Hz, $J_2 = 12.0$ Hz), 2.44 (t, 2H, $J = 8.1$ Hz), 2.08 (tt, 2H, $J_1 = J_2 = 7.5$ Hz), 1.74

5 (2-Isopropylphenyl)[2-trifluoromethyl-4-(*E*-((cyclobutylamino)carbonyl)
ethenyl) phenyl] sulfide

¹H NMR (CDCl₃, 300 MHz) δ 7.76 (s, 1H), 7.52 (d, 1H, J = 15.4 Hz), 7.43-7.49 (m, 3H), 7.33 (d, 1H, J = 7.7 Hz), 7.22 (m, 1H), 6.79 (d, 1H, J = 8.1 Hz), 6.33 (d, 1H, J = 15.4 Hz), 5.72 (br, 1H), 4.52 (m, 1H), 3.49 (dq, 1H, J₁ = J₂ = 6.9 Hz), 2.40 (m, 2H), 1.90 (m, 2H), 1.74 (m, 2H), 1.17 (d, 6H, J = 6.6 Hz). MS (ESI) *m/z* 420, 839, 861. Anal. Calcd for C₂₃H₂₄F₃N₁O₁S: C, 65.85; H, 5.77; N, 3.34. Found: C, 65.53; H, 5.83; N, 3.21.

15 Example 79
(2-Isopropylphenyl)[2-trifluoromethyl-4-(E-((cyclopentylamino)carbonyl)
ethenyl) phenyl] sulfide

¹H NMR (CDCl₃, 300 MHz) δ 7.77 (s, 1H), 7.52 (d, 1H, J = 15.5 Hz), 7.43-7.48 (m, 3H), 7.33 (d, 1H, J = 8.8 Hz), 7.22 (m, 1H), 6.79 (d, 1H, J = 8.1 Hz), 6.33 (d, 1H, J = 15.5 Hz), 5.54 (d, J = 7.7, 1H), 4.35 (m, 1H), 3.49 (dq, 1H, J₁ = J₂ = 6.9 Hz), 2.05 (m, 2H), 1.68 (m, 4H), 1.44(m, 2H), 1.17 (d, 6H, J = 7.0 Hz). MS (ESI) *m/z* 434, 867, 889.

Anal. Calcd for $C_{24}H_{26}F_3N_1O_1S$: C, 66.49; H, 6.04; N, 3.23. Found: C, 66.24; H, 6.14, N, 3.06.

Example 80

5 (2-Isopropylphenyl)[2-trifluoromethyl-4-(E-((5-hydroxypent-1-ylamino)carbonyl)ethenyl) phenyl] sulfide

The title compound was prepared according to the procedures of Example 1.

¹H NMR (CDCl₃, 300 MHz) δ 7.77 (s, 1H), 7.54 (d, 1H, J = 15.5 Hz), 7.43-7.49 (m, 3H), 7.33 (d, 1H, J = 8.0 Hz), 7.22 (m, 1H), 6.79 (d, 1H, J = 8.4 Hz), 6.35 (d, 1H, J = 15.6 Hz), 5.67 (br, 1H), 3.67 (t, 2H, J = 6.4 Hz), 3.49 (dq, 1H, J₁ = J₂ = 6.9 Hz), 3.40 (m, 2H), 2.40 (m, 2H), 1.45-1.62 (m, 6H), 1.17 (d, 6H, J = 7.0 Hz). MS (ESI) *m/z* 452, 474, 903, 925. Anal. Calcd for $C_{24}H_{28}F_3NO_2S \cdot 0.56$ EtOAc: C, 62.92; H, 6.54; N, 2.80. Found: C, 62.86; H, 6.53; N, 2.96.

Example 81

15 (2-Isopropylphenyl)[2-nitro-4-(E-((3-carbomethoxy-4-acetyl)piperazin-1-yl)carbonyl)ethenyl) phenyl] sulfide

Prepared according to the procedures of Example 71, giving a yellow solid.

¹H NMR (CDCl₃, 300MHz) δ 1.14 (d, J = 6.6 Hz, 6H); 2.20 (s, 3H); 2.75-3.80 (br m, 4H); 3.39-3.50 (m, 1H); 3.70, 3.77 (br s, br s, 3H); 4.49-4.75 (br m, 2H); 5.39 (br s, 1H); 6.71(m, 1H); 6.91-7.04 (br m, 1H); 7.25-7.64 (m, 6H); 8.42 (br m, 1H). MS

(APCI) (M+H)⁺ at *m/z* 512. Anal calcd for C₂₆H₂₉N₃S₁O₆: C, 61.04; H, 5.71; N, 8.21.

Found: C, 61.40; H, 6.05; N, 7.88.

Example 82

5 (2-Biphenyl)[2-chloro-4-(*E*-((morpholin-1-yl)carbonyl)ethenyl) phenyl] sulfide

To a stirred solution of bromide from Example 12 (60 mg, 0.14 mmol) in 1 mL of toluene was added 0.5 mL of sat. Na₂CO₃, Pd(PPh₃)₄ (8 mg, 0.007 mmol), phenylboronic acid (17 mg, 0.14 mmol). The mixture was flushed with nitrogen and
10 heated at 100 °C for 3 h. The reaction mixture was then allowed to cool down to room temperature and partitioned between ethyl acetate and brine. The organic layer was then dried with Na₂SO₄, filtered, and concentrated in vacuo. The crude product was purified using Gilson Preparative HPLC as described in Example 38B to give the title
15 compound as colorless oil (40 mg, 67% yield); ¹H NMR (CDCl₃, 300 MHz) δ 3.58-3.86 (m, 8H), 6.77 (d, *J* = 15.6 Hz, 1H), 6.86 (d, *J* = 8.4 Hz, 1H), 7.67 (dd, *J* = 2.1, 8.4 Hz, 1H), 7.29-7.40 (m, 3H), 7.40-7.48 (m, 6H), 7.56 (d, *J* = 15.6 Hz, 1H), 7.65 (d, *J* = 1.8 Hz, 1H). MS (APCI⁺) (M+H)⁺ at *m/z* 436, 438.

Example 83

20 (3,4-Dimethylphenyl)[2-nitro-4-(*E*-((4-acetylpiperazin-1-yl)carbonyl)ethenyl)phenyl]sulfide

To a solution of the compound of Example 32A (40 mg, 0.12 mmole) in 2.5 mL of dimethylformamide was added 3,4-dimethylthiophenol (17 mg, 0.12 mmole), followed by potassium carbonate powder (20 mg, 0.14 mmole). The mixture was heated at 100°C for 20 h. The solvent was removed using N₂ gas flow. Water (5 mL) was then added to the residue, the resulting precipitate was collected through filtration, washed with cold water, and air dried to give the title compound (42 mg, 81%) as light yellow solid. ¹H-NMR (CDCl₃, 400 MHz) δ 2.08 (s, 3H), 2.23 (s, 3H), 2.27 (s, 3H), 3.45 (br, m, 2H), 3.63 (br, m, 6H), 6.79 (s, 1H), 6.82 (d, J = 19 Hz, 1H), 7.18 (d, J = 19 Hz, 1H), 7.24 (dd, J = 4, 19 Hz, 1H), 7.27 (s, 1H), 7.34 (d, J = 21 Hz, 1H), 7.56 (d, J = 39 Hz, 1H), 8.32 (d, J = 4 Hz, 1H). MS (APCI) (M+H)⁺ at m/z 440. FAB High Resolution MS calculated m/z for C₂₃H₂₆N₃O₄S (M+H)⁺: 440.1644. Observed m/z: 440.1646.

Example 84

(2-Bromophenyl)[2-trifluoromethyl-4-(*E*-((4-acetyl)piperazin-1-yl)carbonyl)ethenyl] sulfide

The title compound was prepared by the procedures described in Example 9 substituting 2,4-dichlorothiophenol with 2-bromothiophenol, and 3,4-dichlorobenzaldehyde with 4-fluoro-3-trifluoromethylbenzaldehyde, to give a white solid. ¹H NMR (d⁶-DMSO, 300 MHz) δ 2.04 (s, 3H), 3.43-3.80 (m, 8H), 7.21 (dd, J = 2.1, 8.4 Hz, 1H), 7.24 (d, J = 8.4 Hz, 1H), 7.33 (td, J = 2.1, 7.65 Hz, 1H), 7.42 (td, J = 1.8, 7.65 Hz, 1H), 7.45 (d, J = 15.6 Hz, 1H), 7.58 (d, J = 15.6 Hz, 1H), 7.78 (dd, J =

1.8, 8.4 Hz, 1H), 7.96 (dd, $J = 1.8, 8.4$ Hz, 1H), 8.25 (d, $J = 1.8$ Hz, 1H). MS (APCI⁺) ($M + NH_4$)⁺ at m/z 530, 532, 534.

Example 85

5 (5-Indolyl)[2-chloro-4-(*E*-((4-acetylpiperazin-1-yl)carbonyl)
ethenyl) phenyl] sulfide

To a stirred solution of 5-iodoindole (255 mg, 1.05 mmol) in 5.0 mL of anhydrous DMF was added the potassium thiolate (457 mg, 1.26 mmol) from Example 65B, followed by K_2CO_3 (174 mg, 1.26 mmol), and cuprous iodide (20 mg, 0.11 mmol). The resulting mixture was then heated at 120 °C for overnight. The reaction mixture was then allowed to cool to ambient temperature and poured into water. The aqueous mixture was extracted twice with 25 mL of ethyl acetate. The combined organic layer was then washed with water and brine, dried over Na_2SO_4 , filtered, concentrated on a rotavap under reduced pressure. The crude product was purified using Gilson Preparative HPLC as described in Example 38B to give the title compound (115 mg, 25 % based on the iodide) as a light-brown solid. 1H NMR (d^6 -DMSO, 300 MHz) δ 2.03 (s, 3H), 3.40-3.78 (m, 8H), 6.51 (d, J = 8.4 Hz, 1H), 6.53 (s, 1H), 7.23 (dd, J = 2.1, 8.4 Hz, 1H), 7.27 (d, J = 15.6 Hz, 1H), 7.39 (d, J = 15.6 Hz, 1H), 7.41 (dd, J = 1.8, 8.4 Hz, 1H), 7.49 (t, J = 2.7 Hz, 1H), 7.56 (d, J = 8.4 Hz, 1H), 7.85 (d, J = 1.8 Hz, 1H), 7.99 (d, J = 1.8 Hz, 1H). MS (APCI $^{+}$) ($M+NH_4$) $^{+}$ at m/z 440, 442. Anal. Calcd for $C_{23}H_{22}ClN_3O_2S \cdot 0.53 CH_2Cl_2$: C, 58.28; H, 4.79; N, 8.66. Found: C, 58.31; H, 4.93; N, 8.65.

Example 86

(5-Benzodioxolyl)[2-chloro-4-(E-((4-acetyl)piperazin-1-yl)carbonyl)
ethenyl) phenyl] sulfide

- 5 The title compound was prepared by the procedures described in Example 85 substituting 5-iodoindole with 1-iodo-3,4-methylenedioxybenzene, providing a white solid. ¹H NMR (CDCl₃, 300 MHz) δ 2.14 (s, 3H), 3.48-3.60 (m, 2H), 3.60-3.84 (m, 6H), 6.05 (s, 2H), 6.75 (d, *J* = 8.4 Hz, 1H), 6.80 (d, *J* = 15.3 Hz, 1H), 6.88 (d, *J* = 8.4 Hz, 1H), 6.98 (d, *J* = 2.1 Hz, 1H), 7.08 (dd, *J* = 2.1, 8.4 Hz, 1H), 7.19 (d, *J* = 1.8, 8.4 Hz, 1H), 7.52 (d, *J* = 2.1 Hz, 1H), 7.58 (d, *J* = 15.6 Hz, 1H). MS (APCI⁺) (M+NH₄)⁺
- 10 at *m/z* 445, 447.

Example 87

(2-Isopropylphenyl)[2-nitro-4-(E-((2-carbomethoxypiperazin-1-yl)carbonyl)ethenyl)
phenyl] sulfide

- 15 Prepared according to the procedures of Example 71, giving a yellow solid. ¹H NMR (DMSO-*d*₆, 300MHz) δ 1.14 (d, *J* = 6.6 Hz, 6H); 2.52-2.91 (br m, 5H); 3.30-3.40 (m, 1H); 3.68, 3.69 (s, s, 3H); 4.10-4.25 (br m, 1H); 5.00-5.21 (br m, 1H); 6.60-6.65 (m, 1H); 7.29-7.62 (m, 6H); 7.85-7.95 (m, 1H); 8.64-8.68 (m, 1H). MS (APCI)
- 20 (M+H)⁺ at *m/z* 470.

Example 88

(2,3-Dimethoxyphenyl)-[2-chloro-4-(E-[(morpholin-1-yl)carbonyl]ethenyl)phenyl]
sulfide

The title compound was prepared according to the procedures of Example 1, giving a white solid, m.p. 148-150C. ¹H NMR (CDCl₃, 300 MHz) δ 3.60-3.78 (m, 8H), 3.85 (s, 3H), 3.91 (s, 3H), 6.78 (d, J=16Hz, 1H), 6.86-6.98 (m, 3H), 7.20 (dd, J=9Hz, 2Hz, 1H), 7.54 (d, J=2Hz, 1H), 7.58 (d, J=16Hz, 1H). Anal. Calcd. for C₂₁H₂₂ClNO₄S: C, 60.06; H, 5.28; N, 3.33. Found: C, 59.72; H, 5.34; N, 2.97.

Example 89

(2-Fluorophenyl)[2-nitro-4-(E-((4-acetyl)piperazin-1-yl)carbonyl]ethenyl)phenyl]sulfide

The title compound was prepared by the procedures described in Example 83 substituting 3,4-dimethylthiophenol with 2-fluorothiophenol. Yellow solid (40 mg, 78%); ¹H-NMR (CDCl₃, 400 MHz) δ 2.17 (s, 3H), 3.56 (br, m, 2H), 3.77 (br, m, 6H), 6.88 (dd, J=3, 21 Hz, 1H), 6.93 (d, J=39 Hz, 1H), 7.26 (dd, J=3, 21 Hz, 1H), 7.33 (dd, J=3, 19 Hz, 1H), 7.49 (br, d, J=20 Hz, 1H), 7.58 (m, 1H), 7.66 (m, 2H), 8.46 (d, J=4 Hz, 1H). MS (APCI) (M+H)⁺ at m/z 430. FAB High Resolution MS calculated m/z for C₂₁H₂₁N₃O₄FS (M+H)⁺: 430.1237. Observed m/z: 430.1246.

Example 90

(2-Bromophenyl)[2-trifluoromethyl-4-(E-((4-tert-butoxycarbonyl)piperazin-1-yl)carbonyl]ethenyl)phenyl] sulfide

The title compound was prepared by the procedures described in Example 1 substituting 2,4-dichlorothiophenol with 2-bromothiophenol, 2-chlorobenzaldehyde with 4-fluoro-3-trifluoromethylbenzaldehyde, and 6-amino-1-hexanol with *t*-butyl 1-piperazinecarboxylate, to give a white solid. ¹H NMR (CDCl₃, 300 MHz) δ 1.48 (s, 9H), 3.49 (br s, 4H), 3.56-3.78 (m, 4H), 6.89 (d, *J* = 15.6 Hz, 1H), 7.10 (d, *J* = 8.4 Hz, 1H), 7.18-7.35 (m, 3H), 7.49 (d, *J* = 8.4 Hz, 1H), 7.65 (d, *J* = 15.6 Hz, 1H), 7.68 (dd, *J* = 2.1, 8.4 Hz, 1H), 7.85 (br s, 1H). MS (APCI) (M+Cl)⁺ at *m/z* 605, 607, 609. Anal. Calcd for C₂₅H₂₆N₂O₃BrF₃S · 0.03 H₂O: C, 52.50; H, 4.59; N, 4.90. Found: C, 52.54; H, 4.71; N, 4.68.

Example 91

(2-(Pyrrolidin-1-yl)phenyl)[2-trifluoromethyl-4-(*E*-((4-*tert*-butoxycarbonyl)piperazin-1-yl)carbonyl)ethenyl] phenyl] sulfide

The title compound was prepared by the procedures described in Example 69 substituting the bromide from Example 12 with the bromide from Example 90, to give a white solid. ¹H NMR (CDCl₃, 300 MHz) δ 1.85 (s, 9H), 1.85 (br s, 4H), 3.32-3.55 (m, 8H), 3.55-3.78 (m, 4H), 6.76 (d, *J* = 8.4 Hz, 1H), 6.82 (d, *J* = 15.6 Hz, 1H), 7.23-7.45 (m, 5H), 7.61 (d, *J* = 15.6 Hz, 1H), 7.75 (br s, 1H). MS (APCI) (M+H)⁺ at *m/z* 562.

Example 92

(3-Carboxamidophenyl)[2-nitro-4-(*E*-((4-acetyl)piperazin-1-yl)carbonyl]

ethenyl phenyl sulfideExample 92A(3-Carboxyphenyl)[2-nitro-4-(*E*-((4-acetyl)piperazin-1-yl)carbonyl)5 ethenyl phenyl sulfide

The title compound was prepared by the procedures described in Example 32B substituting 2,4-dichlorothiophenol with 3-mercaptopbenzoic acid.

Example 92B10 (3-Carboxamidophenyl)[2-nitro-4-(*E*-((4-acetyl)piperazin-1-yl)carbonyl)ethenyl phenyl sulfide

To a stirred solution of benzoic acid from Example 92A (40 mg, 0.088 mmol) in 1 mL of anhydrous DMF with HOBT (15 mg, 0.097 mmol) was added EDAC (19 mg, 0.097 mmol), followed by ammonium chloride (large excess). The pH of the solution was adjusted to 6 with addition of triethylamine. The resulting mixture was then stirred at ambient temperature for 6 h. Water was added to quenched the reaction. The product precipitated out after stirring for 30 min, which was then isolated by filtration and dried in vacuum oven to give a light yellow solid (25 mg, 63% yield). ¹H NMR (d⁶-DMSO, 300 MHz) δ 2.04 (s, 3H), 3.43-3.82 (m, 8H), 6.84 (d, *J* = 8.7 Hz, 1H), 7.43 (d, *J* = 15.6 Hz, 1H), 7.53 (d, *J* = 15.6 Hz, 1H), 7.56 (d, *J* = 1.8 Hz, 1H), 7.66 (t, *J* = 7.65 Hz, 1H), 8.06 (d, *J* = 7.80 Hz, 1H), 8.12 (s, 2H), 8.67 (d, *J* = 2.1 Hz, 1H). MS (ESI⁺) (M+Na)⁺ at *m/z* 477.

Example 93

(3-(Hydroxymethyl)phenyl)[2-nitro-4-(*E*-((4-acetyl)piperazin-1-yl)carbonyl)ethenyl] phenyl] sulfide

- 5 To a stirred solution of benzoic acid from Example 92A (255 mg, 0.56 mmol) in 5 mL of anhydrous THF at 0 °C was added in turn Et₃N (102 mL, 0.73 mmol) and ethyl chloroformate (70 mL, 0.73 mmol). After 60 min, the reaction mixture was filtered through celite plug into a stirred solution of NaBH₄ in water at 0 °C. The resulting reaction mixture stirred at 0 °C for 2 h before it was extracted with EtOAc
- 10 (2×20 mL). The combined organic layers was washed with 3N HCl, brine, dried over Na₂SO₄, filtered, concentrated under reduced pressure. The crude product was purified using Gilson Preparative HPLC as described in Example 38B to give the title compound (80 mg, 32% yield) as a light-yellow solid. ¹H NMR (d⁶-DMSO, 300 MHz) δ 2.04 (s, 3H), 3.40-3.79 (m, 8H), 4.56 (s, 2H), 5.38 (br s, 1H), 6.85 (d, *J* = 8.7 Hz,
- 15 1H), 7.42 (d, *J* = 15.6 Hz, 1H), 7.52 (br s, 3H), 7.57 (br s, 2H), 7.91 (dd, *J* = 2.1, 8.7 Hz, 1H), 8.66 (d, *J* = 2.1 Hz, 1H). MS (APCI⁺) (M+NH₄)⁺ at *m/z* 459.

Example 94

Phenyl[2-trifluoromethyl-4-(*E*-((4-*tert*-butoxycarbonyl)piperazin-1-yl)carbonyl)ethenyl] phenyl] sulfide

20

The title compound was obtained as a reductive side product from the reaction mixture described in Example 91, as a colorless oil. ¹H NMR (CDCl₃, 300 MHz) δ

1.49 (s, 9H), 3.43-3.56 (br s, 4H), 3.56-3.82 (m, 4H), 6.85 (d, $J = 15.6$ Hz, 1H), 7.06 (d, $J = 8.4$ Hz, 1H), 7.37-7.50 (m, 4H), 7.63 (d, $J = 15.6$ Hz, 1H), 7.67 (d, $J = 8.4$ Hz, 1H), 7.76 (d, $J = 11.7$ Hz, 1H), 7.80 (s, 1H). MS (APCI) $(M+Cl)^+$ at m/z 527.

5

Example 95

(2-Isopropylphenyl)[2-trifluoromethyl-4-(*E*-(2-carbomethoxy-4-(*tert*-butoxycarbonyl)piperazin-1-yl)carbonyl)ethenyl) phenyl] sulfide

The title compound was prepared according to the procedures of Example 71.

¹H NMR (CDCl₃, 300 MHz) δ 7.79 (s, 1H), 7.62 (d, 1H, $J = 15.0$ Hz), 7.48 (d, 1H, $J = 7.2$ Hz), 7.43 (m, 2H), 7.38 (d, 1H, $J = 8.1$ Hz), 7.22 (m, 1H), 6.86 (d, 1H, $J = 15.4$ Hz), 6.80 (d, 1H, $J = 8.4$ Hz), 5.30 (br, 1H), 4.62 (br d, 2H, $J = 14.0$ Hz), 3.89 (br m, 1H), 3.76 (s, 3H), 3.49 (dq, 1H, $J_1 = J_2 = 6.9$ Hz), 3.12 (m, 2H), 2.94 (br, 1H), 1.46 (s, 9H), 1.17 (d, 6H, $J = 6.6$ Hz). MS (ESI) m/z -591, -627, -677.

15

Example 96

(2-Isopropylphenyl)[2-nitro-4-(*E*-(3-(pyridine-4-methylaminocarbonyl)-4-*tert*-butoxycarbonyl)piperazin-1-yl)carbonyl)ethenyl) phenyl] sulfide

Prepared according to the procedures of Example 71, giving a yellow solid.

¹H NMR (DMSO-*d*₆, 300MHz) δ 1.14 (d, $J = 6.6$ Hz, 6H); 1.38 (s, 9H); 2.83-3.85 (br m, 5H); 4.09-4.51 (br m, 4H); 4.91-5.09 (br m, 1H); 6.64 (d, $J = 8.5$ Hz, 1H); 7.12-7.62 (m, 8H); 7.82-7.96 (m, 1H); 8.26-8.48 (m, 2H); 8.63-8.75 (m, 2H). MS (APCI)

(M+H)⁺ at m/z 646. Anal calcd for C₃₄H₃₉N₅S₂O₆: C, 63.24; H, 6.09; N, 10.84. Found: C, 63.07; H, 6.43; N, 10.54.

Example 97

5 (2-Ethoxyphenyl)-[2-chloro-4(E-[(morpholin-1-yl)carbonyl]ethenyl)phenyl]sulfide

Example 97A

2-Ethoxybenzenethiol

To 7.82g of ethoxybenzene and 7.41g of tetramethylethylenediamine in 75 ml ether, cooled in an ice bath, a solution of 25.6 ml of a 2.5 M n-butyllithium solution in
 10 hexane, was added dropwise under a nitrogen atmosphere. The mixture was stirred for 1 hour at room temperature and then cooled to -65 degrees. Sulfur (2.28 g) was added in portions. The mixture was stirred for 3 hours at room temperature and then cooled in ice. LiAlH₄ (0.6 g) was added and the mixture was stirred 1 hour at room temperature. The mixture was again cooled in ice while 5 ml water was added
 15 dropwise followed by 15% HCl in water until all salts. The aqueous phase was separated and washed with ether. The combined ether layers was washed with HCl, then water. After drying with Na₂SO₄, the ether was evaporated to give 9.66 g of product. NMR analysis showed 70% pure material with 30% of a diaryl sulfide impurity. This mixture was carried forward to the next step.

20

Example 97B

(2-Ethoxyphenyl)-[2-chloro-4(E-[(morpholin-1-yl)carbonyl]ethenyl)phenyl]sulfide

The title compound was prepared according to the procedures of Example 1, substituting the thiol of Example 97A, giving a white solid, m.p. 125-127°C. ¹H NMR (CDCl₃, 300 MHz) δ 1.25 (t, J=7Hz, 3H), 3.60-3.78 (m, 8H), 4.05 (q, J=7Hz, 2H), 6.76 (d, J=15Hz, 1H), 6.82 (d, J=9H, 1H), 6.94-7.00 (m, 2H), 7.16 (dd, J=9Hz, 2H, 5 1H), 7.34-7.45 (m, 2H), 7.54 (d, J=2Hz, 1H), 7.58 (d, J=15Hz, 1H). Anal. Calcd. for C₂₁H₂₂ClNO₃S: C, 62.44; H, 5.49; N, 3.47. Found: C, 62.14; H, 5.70; N, 3.22.

Example 98

(2-Methoxyphenyl)[2-nitro-4-(E-((4-acetyl)piperazin-1-yl)carbonyl)ethenyl)phenyl]sulfide

10

The title compound was prepared by the procedures described in Example 83 substituting 3,4-dimethylthiophenol with 2-methoxythiophenol, giving a yellow solid (40 mg, 77%). ¹H-NMR (CDCl₃, 400 MHz) δ 2.14 (s, 3H), δ 3.54 (br, m, 2H), δ 3.68 (br, m, 6H), δ 3.79 (s, 3H), δ 6.81 (d, J = 21 Hz, 1H), δ 6.89 (d, J = 39 Hz, 1H), δ 7.03 15 (d, J = 21 Hz, 1H), δ 7.08 (m, 1H), δ 7.41 (br, d, J = 21 Hz, 1H), δ 7.53 (m, 1H), δ 7.60 (m, 1H), δ 7.65 (br, s, 1H), δ 8.42 (br, s, 1H). MS (APCI) (M+H)⁺ at m/z 442.

Example 99

(2-(Azetidin-1-yl)phenyl)[2-trifluoromethyl-4-(E-((4-tert-butoxycarbonyl)piperazin-1-yl)carbonyl)ethenyl) phenyl] sulfide

20

The title compound was prepared by the procedures described in Example 69 substituting pyrrolidine with azetidine hydrochloride, and the bromide from Example

12 with bromide from Example 90, giving a white solid. ¹H NMR (CDCl₃, 300 MHz) δ 1.48 (s, 9H), 2.18 (pentet, *J* = 7.43 Hz, 2H), 3.40-3.53 (m, 4H), 3.53-3.77 (m, 4H), 4.02 (t, *J* = 7.43 Hz, 4H), 6.54 (d, *J* = 8.7 Hz, 1H), 6.72 (d, *J* = 8.7 Hz, 1H), 6.78 (tt, *J* = 1.5, 7.35 Hz, 1H), 6.81 (d, *J* = 15.6 Hz, 1H), 7.29-7.42 (m, 3H), 7.61 (d, *J* = 15.6 Hz, 1H), 7.75 (br s, 1H). MS (APCI⁺) (M+H)⁺ at *m/z* 548.

Example 100

(2-(Piperidin-1-yl)phenyl)[2-trifluoromethyl-4-(*E*-((4-(*tert*-butoxycarbonyl)piperazin-1-yl)carbonyl)ethenyl) phenyl] sulfide

10 The title compound was prepared by the procedures described in Example 69 substituting pyrrolidine with piperidine, and the bromide from Example 12 with bromide from Example 90, and isolated as a white solid. ¹H NMR (CDCl₃, 300 MHz) δ 1.48 (s, 9H), 1.54 (br s, 6H), 2.96 (br s, 4H), 3.48 (br s, 4H), 3.55-3.78 (m, 4H), 6.86 (d, *J* = 15.6 Hz, 1H), 6.99 (td, *J* = 1.8, 7.5 Hz, 1H), 7.08 (d, *J* = 8.4 Hz, 1H), 7.19 (dd, *J* = 1.8, 8.1 Hz, 1H), 7.25 (br m, 1H), 7.31 (td, *J* = 1.8, 7.5 Hz, 1H), 7.42 (dd, *J* = 1.8, 8.4 Hz, 1H), 7.65 (d, *J* = 15.6 Hz, 1H), 7.71 (d, *J* = 1.8 Hz, 1H). MS (APCI⁺) (M+H)⁺ at *m/z* 576.

Example 101

20 (3-Chloro-2-formylphenyl)[2-chloro-4-(*E*-((4-acetyl)piperazin-1-yl)carbonyl)ethenyl) phenyl] sulfide

The title compound was prepared by the procedures described in Example 65B substituting 2,3-dichlorobenzaldehyde with 2,6-dichlorobenzaldehyde, isolated as a white solid. ^1H NMR (CDCl_3 , 300 MHz) δ 2.05 (s, 3H), 3.56 (br s, 2H), 3.61-3.86 (m, 6H), 6.68 (q, J = 3.0 Hz, 1H), 6.93 (d, J = 15.6 Hz, 1H), 7.23 (d, J = 3.0 Hz, 1H), 7.25 (m, 1H), 7.45 (dd, J = 2.1, 8.4 Hz, 1H), 7.62 (d, J = 8.4 Hz, 1H), 7.67 (d, J = 15.6 Hz, 1H), 7.69 (d, J = 2.1 Hz, 1H). MS (APCI $^+$) ($\text{M}+\text{H}$) $^+$ at m/z 463, 465, 467.

Example 102

(2-Trifluoromethylphenyl)[2-trifluoromethyl-4-(*E*-((4-acetylpiperazin-1-yl)carbonyl)ethenyl) phenyl] sulfide

The title compound was prepared according to the procedures of Example 1. ^1H NMR (CDCl_3 , 300 MHz) δ 7.84 (s, 1H), 7.80 (m, 1H), 7.66 (d, 1H, J = 15.4 Hz), 7.49 (m, 3H), 7.40 (m, 1H), 7.06 (d, 1H, J = 8.0 Hz), 6.87 (d, 1H, J = 15.4 Hz), 3.62-3.80 (m, 6H), 3.53 (m, 2H), 2.15 (s, 3H). MS (ESI) m/z 503, 525, 1027.

Example 103

(3-Bromophenyl)[2-trifluoromethyl-4-(*E*-((4-acetylpiperazin-1-yl)carbonyl)ethenyl) phenyl] sulfide

The title compound was prepared according to the procedures of Example 1. ^1H NMR (CDCl_3 , 300 MHz) δ 7.83 (s, 1H), 7.66 (d, 1H, J = 15.4 Hz), 7.57 (t, 1H, J = 1.9 Hz), 7.49 (m, 2H), 7.36 (dt, 1H, J = 1.6, 7.8 Hz), 7.24 (m, 1H), 7.18 (d, 1H, J =

8.1 Hz), 6.87 (d, 1H, J = 15.2 Hz), 3.62-3.82 (m, 6H), 3.54 (m, 2H), 2.15 (s, 3H). MS (ESI) m/z 514, 515, 535, 537.

Example 104

5 (3,5-Dimethylphenyl)[2-trifluoromethyl-4-(*E*-((4-acetyl)piperazin-1-yl)carbonyl)ethenyl] phenyl] sulfide

The title compound was prepared according to the procedures of

Example 1. ^1H NMR (CDCl_3 , 300 MHz) δ 7.79 (s, 1H), 7.64 (d, 1H, J = 15.1 Hz), 7.42 (d, 1H, J = 8.8 Hz), 7.49 (m, 2H), 7.13 (s, 2H), 7.04 (s, 2H), 6.84 (d, 1H, J = 15.2 Hz), 3.62-3.82 (m, 6H), 3.54 (m, 2H), 2.32 (s, 6H), 2.15 (s, 3H). MS (ESI) m/z 463, 485, 925, 947.

Example 105

15 (2-Isopropylphenyl)[2-nitro-4-(*E*-((3-dimethylaminocarbonyl-4-(pyridine-4-carbonyl)piperazin-1-yl)carbonyl)ethenyl] phenyl] sulfide

Prepared according to the procedures of Example 71, giving a yellow solid.

^1H NMR ($\text{DMSO}-d_6$, 300MHz) δ 1.14 (d, J = 6.6 Hz, 6H); 2.50-3.83 (br m, 10H); 4.04-4.66 (br m, 3H); 5.32-5.43 (br m, 1H); 6.60-6.69 (m, 1H); 7.15-7.64 (m, 8H); 7.85-7.93 (m, 1H); 8.59-8.72 (m, 3H). MS (APCI) ($\text{M}+\text{H}^+$) at m/z 588. Anal calcd for $\text{C}_{31}\text{H}_{33}\text{N}_5\text{O}_5 \cdot 0.67\text{H}_2\text{O}$: C, 62.07; H, 5.77; N, 11.68. Found: C, 62.13; H, 6.01; N, 11.48.

Example 106

(2-Isopropylphenyl)[2-nitro-4-(*E*-((3-dimethylaminocarbonyl)-4-carbomethoxypiperazin-1-yl)carbonyl)ethenyl] phenyl] sulfide

Prepared according to the procedures of Example 71, giving a yellow solid.

- 5 ¹H NMR (DMSO-d₆, 300MHz) δ 1.14 (d, J = 6.6 Hz, 6H); 2.50-3.83 (br m, 14H); 4.16-4.63 (br m, 2H); 4.98 (br s, 1H); 6.60-6.69 (m, 1H); 7.20-7.61 (m, 6H); 7.85-7.93 (m, 1H); 8.59-8.65 (m, 1H). MS (APCI) (M+H)⁺ at m/z 541.

Example 107

- 10 (2-Isopropylphenyl)[2-nitro-4-(*E*-((3-dimethylaminocarbonyl)-4-acetylpiperazin-1-yl)carbonyl)ethenyl] phenyl] sulfide

Prepared according to the procedures of Example 71, giving a yellow solid.

- ¹H NMR (DMSO-d₆, 300MHz) δ 1.14 (d, J = 6.6 Hz, 6H); 1.88, 2.04 (s, s, 3H); 2.50-3.83 (br m, 11H); 4.16-4.59 (br m, 2H); 5.04-5.25 (br m, 1H); 6.60-6.69 (m, 1H);
15 7.21-7.62 (m, 6H); 7.85-7.93 (m, 1H); 8.58-8.65 (m, 1H). MS (APCI) (M+H)⁺ at m/z 525. Anal calcd for C₂₇H₃₂N₄S₂O₅: C, 61.81; H, 6.15; N, 10.68. Found: C, 61.93; H, 6.75; N, 9.67.

Example 108

- 20 (2-Isopropylphenyl)[2-nitro-4-(*E*-((3-(1-morpholinocarbonyl)-4-*tert*-butoxycarbonylpiperazin-1-yl)carbonyl)ethenyl] phenyl] sulfide

- Prepared according to the procedures of Example 71, giving a yellow solid.
- ¹H NMR (DMSO-d₆, 300MHz) δ 1.11-1.16 (br m, 6H); 1.35, 1.40 (br s, br s, 9H); 2.67-5.0 (br m, 16H); 6.60-6.69 (m, 1H); 7.28-7.62 (m, 6H); 7.87-7.92 (m, 1H); 8.63-8.67 (br m, 1H). MS (APCI) (M+H)⁺ at m/z 625. Anal calcd for C₃₂H₄₀N₄S₁O₇: C, 61.52; H, 6.45; N, 8.97. Found: C, 61.10; H, 6.65; N, 8.60.

Example 109

(2-Isopropylphenyl)[2-nitro-4-(*E*-((3-(pyridine-4-methylaminocarbonyl)piperazin-1-yl)carbonyl)ethenyl) phenyl] sulfide

- Prepared according to the procedures of Example 71, giving a yellow solid.
- ¹H NMR (DMSO-d₆, 300MHz) δ 1.14 (d, J = 6.6 Hz, 6H); 2.50-4.46 (br m, 10H); 6.63 (d, J = 8.5 Hz, 1H); 7.20-7.64 (m, 8H); 7.85-7.93 (m, 1H); 8.43-8.65 (m, 4H). MS (APCI) (M+H)⁺ at m/z 546. Anal calcd for C₂₉H₃₁N₃O₄·0.46CH₃COOCH₂CH₃: C, 63.20; H, 5.96; N, 11.95. Found: C, 63.29; H, 6.27; N, 11.97.

Example 110

(2-Isopropylphenyl)[2-nitro-4-(*E*-((3-dimethylaminocarbonyl)piperazin-1-yl)carbonyl)ethenyl) phenyl] sulfide

- Prepared according to the procedures of Example 71, giving a yellow solid.
- ¹H NMR (DMSO-d₆, 300MHz) δ 1.14 (d, J = 6.6 Hz, 6H); 2.50-3.20 (br m, 4H); 2.82 (s, 3H); 3.04 (s, 3H); 3.26-3.49 (m, 1H); 3.52-3.59 (m, 1H); 4.08-4.47 (br m, 2H); 6.63 (d, J = 8.5 Hz, 1H); 7.31-7.62 (m, 6H); 7.86-7.92 (m, 1H); 8.61 (br m, 1H). MS

(APCI) (M+H)⁺ at m/z 483. Anal calcd for C₂₅H₃₄N₄S₂O₄·0.39CH₃COOCH₂CH₃: C, 61.71; H, 6.46; N, 10.84. Found: C, 61.96; H, 6.69; N, 10.73.

Example 111

- 5 (2-Isopropylphenyl)[2-nitro-4-(E-((3-(benzylaminocarbonyl)-4-tert-butoxycarbonylpiperazin-1-yl)carbonyl)ethenyl)phenyl] sulfide

Prepared according to the procedures of Example 71, giving a yellow solid.

- ¹H NMR (DMSO-d₆, 300MHz) δ 1.14 (d, J = 6.6 Hz, 6H); 1.33, 1.42 (br s, br s, 9H); 2.75-4.77 (br m, 10H); 6.60-6.66 (br m, 1H); 7.02-7.94 (br m, 12H); 8.47-8.67 (m, 2H). MS (APCI) (M+H)⁺ at m/z 645.

Example 112

- (2-Isopropylphenyl)[2-nitro-4-(E-((3-(dimethylaminocarbonyl)-4-tert-butoxycarbonylpiperazin-1-yl)carbonyl)ethenyl)phenyl] sulfide

- 15 Prepared according to the procedures of Example 71, giving a yellow solid.

- ¹H NMR (DMSO-d₆, 300MHz) δ 1.14 (d, J = 6.6 Hz, 6H); 1.35, 1.40 (br s, br s, 9H); 2.50-4.99 (br m, 14H); 6.60-6.69 (m, 1H); 7.21-7.62 (m, 6H); 7.86-7.92 (m, 1H); 8.59-8.63 (br m, 1H). MS (APCI) (M+H)⁺ at m/z 583. Anal calcd for C₃₀H₃₈N₄S₂O₆·0.21C₆H₁₄: C, 62.50; H, 6.87; N, 9.32. Found: C, 62.28; H, 7.15; N, 9.11.

Example 113

(2-Bromophenyl)[2-chloro-4-(E-((3-(5*S*-hydroxymethyl-pyrrolidin-2-on-1-yl)prop-1-ylamino)carbonyl) ethenyl)phenyl]sulfide

(2-Bromophenyl)[2-chloro-4-(2-carboxy-E-ethenyl) phenyl]sulfide was

prepared by the procedures described in Example 1 substituting 2,4

- 5 dichlorothiophenol with 2-bromothiophenol, 2-chlorobenzaldehyde with 3,4 dichlorobenzaldehyde. 1-(3-aminopropyl)-5-((*S*)-thexyldimethylsilyloxymethyl)-2-pyrrolidinone (0.2818g, 0.8959 mmol) was added to a solution of this cinnamic acid (0.3312g, 0.8959 mmol), 1-[3-(dimethylamino)propyl]-3-ethyl carbodiimide hydrochloride (0.3435g, 1.79 mmol), and 1-hydroxybenzotriazole hydrate (0.1816g,
- 10 1.34 mmol) in DMF (4.0 mL). After stirring for 12h the reaction mixture was diluted with EtOAc (250 mL), extracted with sat. NH_4Cl (1x75 mL), extracted with H_2O (2x75 mL), rinsed with brine (75mL), and dried over Na_2SO_4 . The resultant thexyldimethylsilyl alcohol was purified by flash chromatography (EtOAc) on silica gel (.4974 g, 83%). Tetrabutylammonium fluoride (.68 mL of 1.0 M solution in THF)
- 15 was added dropwise to a solution of this protected alcohol (0.4544 g, 0.682 mmol) in THF (1.7 mL). After 2h the reaction was diluted with EtOAc (50 mL) and extracted with sat. NH_4Cl (1x25 mL), extracted with H_2O (2x25 mL), rinsed with brine (25mL), and dried over Na_2SO_4 . Flash chromatography (EtOAc \rightarrow 9:1 CH_2Cl_2 :MeOH) on silica gel yielded the title compound (.3144g, 88%). $^1\text{H-NMR}$ (DMSO-d_6 , 300MHz) δ
- 20 8.14 (t, $J = 5.5$ Hz, 1H), 7.81 (m, 2H), 7.53 (dd, $J = 8.3, 1.7$ Hz, 1H), 7.44 (dt, $J = 7.7, 1.5$, 1H), 7.40 (dt, $J = 7.7, 1.8$, 1H), 7.39 (d, $J = 15.6$ Hz, 1H), 7.28 (dd, $J = 7.7, 1.8$

0011705.03100

Hz, 1H), 7.05 (d, $J = 8.1$ Hz, 1H), 6.67 (d, $J = 15.6$ Hz, 1H), 4.84 (t, $J = 5.1$ Hz, 1H), 2.94-3.62 (m, 8H), 1.54-2.29 (m, 6H), MS(APCI) (M+H)⁺ at m/z 523, 525, 527, 529.

Example 114

5 (2-Bromophenyl)[2-chloro-4-(E-((3-(pyrrolidin-2-on-1-yl)prop-1-ylamino)carbonyl)ethenyl)phenyl]sulfide

The title compound was prepared by the procedures described in Example 1 substituting 2,4 dichlorothiophenol with 2-bromothiophenol, 2-chlorobenzaldehyde with 3,4 dichlorobenzaldehyde, and 6-amino-1-hexanol with 1-(3-aminopropyl)-2-pyrrolidinone. ¹H-NMR (DMSO-d₆, 300MHz) δ 8.12 (t, $J = 5.9$ Hz, 1H), 7.81 (m, 2H), 7.52 (dd, $J = 8.1, 2.0$ Hz, 1H), 7.44 (dt, $J = 7.5, 1.4$, 1H), 7.34 (dt, $J = 7.5, 2.0$, 1H), 7.39 (d, $J = 15.8$ Hz, 1H), 7.28 (dd, $J = 7.6, 1.9$ Hz, 1H), 7.05 (d, $J = 8.1$ Hz, 1H), 6.67 (d, $J = 15.8$ Hz, 1H), 4.02 (d, $J = .7$ Hz, 1H), 3.29-3.35 (m, 2H), 3.11-3.25 (m, 4H), 2.21 (t, $J = 8.1$ Hz, 1H), 1.94 (m, 2H), 1.64 (m, 2H), MS(APCI) (M+H)⁺ at m/z 493, 495, 497, 499.

Example 115

20 (2-Bromophenyl)[2-chloro-4-(E-(N-methyl-N-(3-(pyrrolidin-2-on-1-yl)prop-1-yl)amino)carbonyl)ethenyl]phenyl]sulfide

The title compound was prepared by the procedures described in Example 1 substituting 2,4 dichlorothiophenol with 2-bromothiophenol, 2-chlorobenzaldehyde with 3,4 dichlorobenzaldehyde, and 6-amino-1-hexanol with 1-(3-

methylaminopropyl)-2-pyrrolidinone. ¹H-NMR (DMSO-d₆, 300MHz) δ 8.06 (d, *J* = 1.5 Hz, 1H), 7.80 (dd, *J* = 7.7, 1.1 Hz, 1H), 7.64 (dd, *J* = 8.5, 1.7 Hz, 1H), 7.25-7.46 (m, 5H), 7.04 (d, *J* = 8.1, 1.1, 1H), 3.14-5.30 (m, 6H), 3.14 (s, 1H), 2.91 (s, 2H), 2.19 (m, 2H), 1.92 (m, 2H), 1.68 (m, 2H), MS(APCI) (M+H)⁺ at *m/z* 507, 509, 511, 513.

5

Example 116

(2-[2-Methoxy]ethoxyphenyl)-[2-chloro-4(E-[(morpholin-1-yl)carbonyl]ethenyl)phenyl] sulfide

The title compound was prepared according to the procedures of Example 97, substituting 2-methoxyethoxybenzene, giving a white solid. ¹H NMR (CDCl₃, 300 MHz) δ 3.29 (s, 3H), 3.60 (t, *J* = 7Hz, 2H), 3.60-3.78 (m, 8H), 4.12 (t, *J* = 7Hz, 2H), 6.78 (d, *J* = 15Hz, 1H), 6.82 (d, *J* = 9Hz, 1H), 6.95-7.03 (m, 2H), 7.18 (dd, *J* = 9Hz, 2H, 1H), 7.36-7.45 (m, 2H), 7.52 (d, *J* = 2Hz, 1H), 7.57 (d, *J* = 15Hz, 1H). Anal. Calcd. for C₂₂H₂₄ClNO₄S: C, 60.85; H, 5.57; N, 3.22. Found: C, 60.65; H, 5.59; N, 3.12.

15

Example 117

(2-Isopropylphenyl)[2-nitro-4-(E-[(3-(morpholinocarbonyl)piperazin-1-yl)carbonyl]ethenyl) phenyl] sulfide

Prepared according to the procedures of Example 71, giving a yellow solid. ¹H NMR (DMSO-d₆, 300MHz) δ 1.14 (d, *J* = 6.6 Hz, 6H); 2.50-3.40 (br m, 6H); 3.42-3.64 (br m, 8H); 4.07-4.44 (br m, 2H); 4.08-4.47 (br m, 2H); 6.64 (d, *J* = 8.5 Hz, 1H); 7.31-7.62 (m, 6H); 7.87-7.92 (m, 1H); 8.61 (br m, 1H). MS (APCI) (M+H)⁺ at *m/z*

0054736.03100 00120.5074500

525. Anal calcd for $C_{27}H_{32}N_4S_2O_5 \cdot 1.57H_2O$: C, 58.64; H, 6.41; N, 10.13. Found: C, 58.69; H, 6.36; N, 9.78.

Example 118

5 (2-Isopropylphenyl)[2-nitro-4-(E-((4-tert-butoxycarbonyl)piperazin-1-yl)carbonyl)ethenyl] phenyl] sulfide

Prepared according to the procedures of Example 71, giving a yellow solid.

1H NMR (DMSO- d_6 , 300MHz) δ 1.14 (d, J = 7.0 Hz, 6H); 1.41 (s, 9H); 3.30-3.40 (m, 1H); 3.50-3.72 (br m, 8H); 6.64 (d, J = 8.5 Hz, 1H); 7.34-7.62 (m, 6H); 7.87-7.92 (dd, J = 8.5, 1.5 Hz, 1H); 8.65 (d, J = 1.5 Hz, 1H). MS (APCI) (M+H) $^+$ at m/z 512. Anal calcd for $C_{27}H_{33}N_3S_2O_5$: C, 63.38; H, 6.50; N, 8.21. Found: C, 63.69; H, 6.62; N, 7.87.

Example 119

15 (2-Isopropylphenyl)[2-nitro-4-(E-((4-methoxycarbonyl)piperazin-1-yl)carbonyl)ethenyl] phenyl] sulfide

Prepared according to the procedures of Example 71, giving a yellow solid.

1H NMR (DMSO- d_6 , 300MHz) δ 1.14 (d, J = 6.8 Hz, 6H); 3.62 (s, 3H); 3.30-3.38 (m, 1H); 3.38-3.72 (br m, 8H); 6.64 (d, J = 8.8 Hz, 1H); 7.34-7.62 (m, 6H); 7.87-7.92 (dd, J = 8.8, 2.0 Hz, 1H); 8.64 (d, J = 2.0 Hz, 1H). MS (APCI) (M+H) $^+$ at m/z 470. Anal calcd for $C_{24}H_{27}N_3S_2O_5 \cdot 0.34C_6H_{14}$: C, 62.77; H, 6.27; N, 8.44. Found: C, 62.70; H, 6.33; N, 8.27.

Example 120

(2-Isopropylphenyl)[2-nitro-4-(E-(4-(pyridine-4-carbonyl)piperazin-1-yl)carbonyl)ethenyl] phenyl] sulfide

Prepared according to the procedures of Example 71, giving a yellow solid.

5 ¹H NMR (DMSO-d₆, 300MHz) δ 1.14 (d, J = 6.6 Hz, 6H); 3.30-3.40 (m, 1H); 3.52-3.86 (br m, 8H); 6.61-6.66 (br m, 1H); 7.30-7.62 (m, 8H); 7.83-7.96 (br m, 1H); 8.60-8.71 (m, 3H). MS (APCI) (M+H)⁺ at m/z 517. Anal calcd for C₂₈H₂₈N₄S₂O₆·0.38CH₃COOCH₂CH₃: C, 64.46; H, 5.69; N, 10.19. Found: C, 64.52; H, 5.94; N, 10.21.

10

Example 121

(2-Isopropylphenyl)[2-nitro-4-(E-((3-(pyridine-3-methylaminocarbonyl)-4-tert-butoxycarbonyl)piperazin-1-yl)carbonyl)ethenyl] phenyl] sulfide

Yellow solid; ¹H NMR (DMSO-d₆, 300MHz) δ 1.14 (d, J = 6.8 Hz, 6H); 1.31-1.46 (br m, 9H); 3.30-3.41 (m, 1H); 3.15-4.78 (br m, 9H); 6.61-6.67 (br m, 1H); 7.05-7.95 (br m, 9H); 8.20-8.65 (br m, 4H). MS (APCI) (M+H)⁺ at m/z 646. Anal calcd for C₃₃H₃₉N₅S₂O₆·0.13H₂O: C, 62.97; H, 6.49; N, 10.79. Found: C, 62.66; H, 6.26; N, 10.60.

20

Example 122

(2-Isopropylphenyl)[2-nitro-4-(E-((3-(pyridine-2-methylaminocarbonyl)piperazin-1-yl)carbonyl)ethenyl] phenyl] sulfide

Prepared according to the procedures of Example 71, giving a yellow solid. ¹H NMR (DMSO-d₆, 300MHz) δ 1.14 (d, J = 7.0Hz, 6H); 3.30-3.41 (m, 1H); 2.50-4.46 (br m, 9H); 6.64 (d, J = 8.5 Hz, 1H); 7.21-7.93 (br m, 10H); 8.45-8.65 (br m, 3H). MS (APCI) (M+H)⁺ at m/z 546.

5

Example 123

(2-Isopropylphenyl)[2-nitro-4-(*E*-(3-(pyridine-3-methylaminocarbonyl)piperazin-1-yl)carbonyl)ethenyl)phenyl] sulfide

Prepared according to the procedures of Example 71, giving a yellow solid. ¹H NMR (DMSO-d₆, 300MHz) δ 1.14 (d, J = 6.6 Hz, 6H); 2.50-4.41 (br m, 10H); 6.61-6.67 (br m, 1H); 7.26-7.70 (br m, 8H); 7.86-7.94 (br m, 1H); 8.40-8.67 (br m, 4H). MS (APCI) (M+H)⁺ at m/z 546.

10

Example 124

(4-Hydroxyphenyl)[2-nitro-4-(*E*-((4-acetyl)piperazin-1-yl)carbonyl)ethenyl)phenyl]sulfide

15

The title compound was prepared by the procedures described in Example 83 substituting 3,4-dimethylthiophenol with 4-hydroxythiophenol. Yellow solid (23 mg, 45%); ¹H-NMR (Pyridine-d₅, 500 MHz) δ 2.08 (s, 3H), 3.42 (br, m, 2H), 3.76 (br, m, 6H), 7.01 (d, J = 17 Hz, 1H), 7.26 (m, 2H), 7.37 (d, J = 31 Hz, 1H), 7.59 (m, 3H), 8.02 (d, J = 31 Hz, 1H), 8.60 (d, J = 4 Hz, 1H). MS (APCI) (M+H)⁺ at m/z 428. FAB

20

High Resolution MS calculated m/z for $C_{21}H_{22}N_3O_3S$ ($M+H$)⁺: 428.1280. Observed m/z : 428.1296.

Example 125

5 (3,5-Dichlorophenyl)[2-nitro-4-(*E*-(4-acetylpiperazin-1-yl)carbonyl)ethenyl]phenyl]sulfide

The title compound was prepared by the procedures described in Example 83 substituting 3,4-dimethylthiophenol with 3,5-dichlorothiophenol. Yellow solid (12 mg, 21%); ¹H-NMR (CDCl₃, 400 MHz) δ 2.04 (s, 3H), 3.43 (br, m, 2H), 3.62 (br, m, 10 6H), 6.82 (d, *J* = 22 Hz, 1H), 6.82 (d, *J* = 38 Hz, 1H), 7.37 (s, 1H), 7.38 (s, 1H), 7.40 (m, 1H), 7.43 (dd, *J* = 3, 21 Hz, 1H), 7.55 (d, *J* = 38 Hz, 1H), 8.29 (d, *J* = 4 Hz, 1H). MS (APCI) ($M+H$)⁺ at m/z 480. FAB High Resolution MS calculated m/z for $C_{21}H_{20}N_3O_4Cl_2S$ ($M+H$)⁺: 480.0552. Observed m/z : 480.0553.

15 Example 126

(2-Bromophenyl)[2-chloro-4-(*E*-(3-(5S-acetoxymethyl-pyrrolidin-2-on-1-yl)prop-1-ylamino)carbonyl)ethenyl]phenyl]sulfide

To a solution of the compound of Example 113 (0.0466g, 0.0889 mmol) in CH₂Cl₂ (5 mL) was added triethylamine (0.024 mL, 0.18 mmol) and acetic anhydride 20 (0.0088 mL, 0.0933 mmol). After 12 h the reaction was diluted with MeOH (1.5 mL) and purified by preparative HPLC to provide the title compound (0.0458 g, 91%). ¹H-NMR (DMSO-*d*₆, 300MHz) δ 8.14 (t, *J* = 5.7 Hz, 1H), 7.80 (m, 2H), 7.53 (dd, *J* = 8.5,

1.5 Hz, 1H), 7.45 (dt, $J = 7.7, 1.5$, 1H), 7.35 (dt, $J = 7.7, 1.8$, 1H), 7.39 (d, $J = 15.6$ Hz, 1H), 7.29 (dd, $J = 7.7, 1.8$ Hz, 1H), 7.05 (d, $J = 8.1$ Hz, 1H), 6.67 (d, $J = 15.6$ Hz, 1H), 4.20 (dd, $J = 11.8, 3.7$ Hz, 1H), 4.03 (dd, $J = 11.8, 4.0$ Hz, 1H), 3.85 (m, 1H), 3.45 (m, 2H), 3.15 (m, 2H), 2.95 (m, 2H), 2.00-2.48 (m, 2H), 2.02 (s, 3H), 1.51-1.82 (m, 2H), MS(APCI) (M+H)⁺ at m/z 565, 567, 569, 571.

Example 127

(2-Bromophenyl)[2-chloro-4-(E-(3-(5S-methoxymethyl-pyrrolidin-2-on-1-yl)prop-1-ylamino)carbonyl) ethenyl]phenyl]sulfide

Sodium hydride (0.0088g, 0.22 mmol, 60% dispersion) was added to a solution of the compound of Example 113 (0.0524g, 0.1 mmol) in DMF (0.5 mL). After 15 min, iodomethane (0.025 mL, 0.4 mmol) was added and the reaction was stirred for 12 h. The reaction was diluted with EtOAc (7 mL) and extracted with sat. NH₄Cl (1x2.5 mL), extracted with H₂O (2x2.5 mL), rinsed with brine (2.5mL), dried over Na₂SO₄, filtered, and concentrated *in vacuo*. The crude products were diluted with MeOH (1.5 mL) and purified by preparative HPLC to provide the title compound (0.0408 g, 74%). ¹H-NMR (DMSO-d₆, 300MHz) δ 8.07 (2, 1H), 7.80 (dd, $J = 7.9, 1.3$ Hz, 1H), 7.64 (dd, $J = 8.3, 1.6$ Hz, 1H), 7.23-7.46 (m, 5H), 7.04 (d, $J = 8.1, 1H$), 3.74 (m, 1H), 4.4-3.52 (m, 6H), 3.27 (s, 1.5H), 3.22 (s, 1.5H), 3.14 (s, 1.5H), 2.91 (s, 1.5H), 1.5-2.3 (m, 6H), MS(APCI) (M+H)⁺ at m/z 551, 553, 555.

Example 128

(2-Bromophenyl)[2-chloro-4-(E-((3-(4*R*-hydroxymethyl-pyrrolidin-2-on-1-yl)prop-1-ylamino)carbonyl) ethenyl)phenyl]sulfide

The title compound was prepared by the procedures described for Example 113 substituting 1-(3-aminopropyl)-5-((*S*)-thexyldimethylsilyloxymethyl)-2-pyrrolidinone with 1-(3-aminopropyl)-4-((*R*)-thexyldimethylsilyloxy)-2-pyrrolidinone. ¹H-NMR (DMSO-d₆, 300MHz) δ 8.13 (t, *J* = 5.5 Hz, 1H), 7.80 (m, 2H), 7.53 (dd, *J* = 8.5, 1.7 Hz, 1H), 7.27-7.44 (m, 4H), 7.05 (d, *J* = 8.1 Hz, 1H), 6.67 (d, *J* = 15.8 Hz, 1H), 5.19 (d, *J* = 3.7 Hz, 1H), 4.28 (br s, 1H), 3.10-3.62 (m, 8H), 2.06 (dd, 1H), 1.63 (m, 1H), MS(APCI) (M+H)⁺ at *m/z* 509, 511, 513.

10

Example 129

Phenyl[2-nitro-4-(E-((4-acetyl)piperazin-1-yl)carbonyl)ethenyl]phenyl]sulfide

The title compound was prepared by the procedures described in Example 83 substituting 3,4-dimethylthiophenol with thiophenol. Yellow solid (36 mg, 73%); ¹H-NMR (CDCl₃, 400 MHz) δ 2.20 (s, 3H), 3.59 (br, m, 2H), 3.78 (br, m, 6H), 6.92 (d, *J* = 21 Hz, 1H), 6.95 (d, *J* = 39 Hz, 1H), 7.49 (br, d, *J* = 21 Hz, 1H), 7.56 (m, 3H), 7.65 (m, 2H), 7.69 (d, *J* = 38 Hz, 1H), 8.46 (d, *J* = 4 Hz, 1H). MS (APCI) (M+H)⁺ at *m/z* 412. FAB High Resolution MS calculated *m/z* for C₂₁H₂₂N₃O₄S (M+H)⁺: 412.1331. Observed *m/z*: 412.1342.

20

Example 130

(2-Dimethylaminophenyl)[2-nitro-4-(E-((4-acetyl)piperazin-1-yl)carbonyl)

ethenyl) phenyl] sulfide

To a stirred solution of aniline from Example 47 (21 mg, 0.049 mmol) in 1 mL of ethanol was added Me₂SO₄ (14.0 mL, 0.15 mmol) followed by sat. Na₂CO₃ (25 mL). The mixture was then refluxed for one day. The reaction mixture was allowed to cool down to ambient temperature, partitioned between EtOAc and water. The organic layer was washed with brine, dried over Na₂SO₄, filtered, concentrated under reduced pressure. The residue was then purified on a Gilson Preparative HPLC as described in Example 38B to give the title compound (10 mg, 45% yield), as a light yellow solid.

¹H NMR (CDCl₃, 300 MHz) δ 2.16 (s, 3H), 2.83 (s, 3H), 3.32 (br s, 3H), 3.47-3.85 (m, 8H), 6.75 (d, *J* = 8.4 Hz, 1H), 6.78 (d, *J* = 8.4 Hz, 1H), 6.82 (d, *J* = 8.4 Hz, 1H), 6.89 (d, *J* = 15.6 Hz, 1H), 7.40-7.51 (m, 3H), 7.64 (d, *J* = 15.6 Hz, 1H), 8.45 (d, *J* = 1.8 Hz, 1H). MS (APCI⁺) (M+H)⁺ at *m/z* 454.

Example 131

(3-((2-Hydroxyethyl)aminocarbonyl)phenyl)[2-nitro-4-(*E*-((4-acetylpiperazin-1-yl)carbonyl)ethenyl) phenyl] sulfide

The title compound was prepared by the procedures described in Example 92B, substituting ammonium chloride with ethanolamine, to give a light yellow solid.

¹H NMR (d⁶-DMSO, 300 MHz) δ 2.04 (s, 3H), 3.30-3.79 (m, 12H), 4.75 (t, *J* = 5.7 Hz, 1H), 6.85 (d, *J* = 8.7 Hz, 1H), 7.42 (d, *J* = 15.6 Hz, 1H), 7.54 (d, *J* = 15.6 Hz, 1H), 7.66 (t, *J* = 7.8 Hz, 1H), 7.79 (d, *J* = 8.1 Hz, 1H), 7.92 (dd, *J* = 2.1, 8.1 Hz, 1H), 8.04

(d, $J = 8.4$ Hz, 1H), 8.11 (s, 1H), 8.62 (t, $J = 5.7$ Hz, 1H), 8.66 (d, $J = 2.1$ Hz, 1H). MS (APCI⁺) (M+Cl)⁺ at m/z 533, 535.

Example 132

5 (3-((3-(1-Imidazolyl)propyl)aminocarbonyl)phenyl)[2-nitro-4-(*E*-((4-acetyl)piperazin-1-yl)carbonyl)ethenyl] phenyl] sulfide

The title compound was prepared by the procedures described in Example 92B, substituting ammonium chloride with 3-aminopropyl-1-imidazole, as a light yellow solid. ¹H NMR (d⁶-DMSO, 300 MHz) δ 1.96 (quintet, $J = 6.98$ Hz, 2H), 2.04
 10 (s, 3H), 3.24 (q, $J = 6.98$ Hz, 2H), 3.35-3.95 (m, 8H), 4.02 (t, $J = 6.98$ Hz, 2H), 6.87 (d, $J = 8.4$ Hz, 1H), 6.88 (s, 1H), 7.19 (s, 1H), 7.41 (d, $J = 15.6$ Hz, 1H), 7.54 (d, $J = 15.6$ Hz, 1H), 7.64 (s, 1H), 7.68 (d, $J = 7.8$ Hz, 1H), 7.79 (dt, $J = 1.8, 7.8$ Hz, 1H), 7.91 (dd, $J = 1.8, 8.7$ Hz, 1H), 8.03 (d, $J = 7.8$ Hz, 1H), 8.09 (t, $J = 1.8$ Hz, 1H), 8.65 (d, $J = 1.8$ Hz, 1H). MS (APCI⁺) (M+Cl)⁺ at m/z 597, 599.

15

Example 133

(3-((2-(1-Morpholinyl)ethyl)aminocarbonyl)phenyl)[2-nitro-4-(*E*-((4-acetyl)piperazin-1-yl)carbonyl)ethenyl] phenyl] sulfide

The title compound was prepared by the procedures described in Example 92B, substituting ammonium chloride with 2-aminoethyl-1-morpholine, as a light
 20 yellow solid. ¹H NMR (d⁶-DMSO, 300 MHz) δ 2.04 (s, 3H), 2.44 (br s, 4H), 3.20-3.80 (m, 16H), 6.87 (d, $J = 8.4$ Hz, 1H), 7.41 (d, $J = 15.6$ Hz, 1H), 7.54 (d, $J = 15.6$ Hz,

1H), 7.68 (d, $J = 8.4$ Hz, 1H), 7.79 (d, $J = 8.4$ Hz, 1H), 7.91 (dd, $J = 2.1, 8.4$ Hz, 1H), 8.02 (d, $J = 8.4$ Hz, 1H), 8.07 (s, 1H), 8.58 (t, $J = 6.0$ Hz, 1H), 8.65 (d, $J = 2.1$ Hz, 1H). MS (APCI) $(M+H)^+$ at m/z 568.

5

Example 134

(2-Isopropylphenyl)[2-nitro-4-(*E*-((3-hydroxymethyl-4-*tert*-butoxycarbonyl)piperazin-1-yl)carbonyl)ethenyl] phenyl] sulfide

Prepared according to the procedures of Example 71, giving a yellow solid.

¹H NMR (DMSO-*d*₆, 300MHz) δ 1.14 (d, $J = 7.0$ Hz, 6H); 1.41 (s, 9H); 2.62-3.20 (br m, 4H); 3.30-3.40 (m, 1H); 3.72-4.44 (br m, 4H); 4.72-4.98 (br m, 1H); 6.62-6.66 (br m, 1H); 7.25-7.63 (m, 6H); 7.83-7.93 (br m, 1H); 8.57-8.66 (br m, 1H). MS (APCI) $(M+H)^+$ at m/z 542. Anal calcd for C₂₃H₃₃N₃O₆·0.21C₆H₁₄: C, 62.78; H, 6.83; N, 7.51. Found: C, 62.65; H, 6.99; N, 7.36.

15

Example 135

(2-Isopropylphenyl)[2-nitro-4-(*E*-((4-formylpiperazin-1-yl)carbonyl)ethenyl) phenyl] sulfide

Prepared according to the procedures of Example 71, giving a yellow solid.

¹H NMR (DMSO-*d*₆, 300MHz) δ 1.14 (d, $J = 7.1$ Hz, 6H); 3.30-3.38 (m, 1H); 3.38-3.77 (br m, 8H); 6.64 (d, $J = 8.5$ Hz, 1H); 7.34-7.62 (m, 6H); 7.88-7.92 (dd, $J = 8.5, 1.7$ Hz, 1H); 8.08 (s, 1H); 8.65 (d, $J = 1.7$ Hz, 1H). MS (APCI) $(M+H)^+$ at m/z 440.

Anal calcd for $C_{23}H_{23}N_3S_2O_4$: C, 62.85; H, 5.73; N, 9.56. Found: C, 63.05; H, 5.98; N, 9.47.

Example 136

5 (2-Isopropylphenyl)[2-nitro-4-(E-((2-hydroxymethyl)-4-tert-butoxycarbonyl)piperazin-1-yl)carbonyl]ethenyl] phenyl] sulfide

Prepared according to the procedures of Example 71, giving a yellow solid. 1H NMR (DMSO- d_6 , 300MHz) δ 1.14 (d, J = 6.8 Hz, 6H); 1.41 (s, 9H); 2.72-3.50 (br m, 4H); 3.30-3.40 (m, 1H); 3.85-4.52 (br m, 4H); 4.74-4.91 (br m, 1H); 6.62-6.66 (br m, 1H); 7.28-7.62 (m, 6H); 7.81-7.91 (br m, 1H); 8.57-8.66 (br m, 1H). MS (APCI) (M+H) $^+$ at m/z 542. Anal calcd for $C_{38}H_{33}N_3S_2O_6 \cdot 0.17C_6H_{14}$: C, 62.65; H, 6.77; N, 7.55. Found: C, 62.54; H, 6.83; N, 7.33.

Example 137

15 (2-Ethoxyphenyl)-[2-chloro-4(E-[(3-ethoxycarbonyl)piperidin-1-yl)carbonyl]ethenyl] phenyl]sulfide

The title compound was prepared according to the procedures of Example 97. 1H NMR (CDCl $_3$, 300 MHz) δ 1.25 (t, J = 7 Hz, 6H), broad peaks totaling 9 protons at 1.50-1.62, 1.65-1.92, 2.01-2.15, 2.45-2.55, 2.95-3.05, 3.13-3.30, 3.55-3.68, 3.90-4.10, 4.05 (q, J=7Hz, 2H), 4.15 (q, J=7Hz, 2H), 6.84 (d, J=9Hz, 1H), 6.80-6.95 (broad, 1H), 6.94-6.99 (m, 2H), 7.18 (dd, J=9Hz, 2Hz, 1H), 7.34-7.41 (m, 2H), 7.52 (d, J=15Hz, 1H), 7.55 (d, J=2Hz, 1H). Anal. Calcd. for

$C_{25}H_{28}ClNO_4S$: C, 63.35; H, 5.95; N, 2.95. Found: C, 63.17; H, 6.02; N, 26.02; N, 2.81.

Example 138

5 (3- Aminophenyl)[2-nitro-4-(E-((4-acetyl)piperazin-1-yl)carbonyl)ethenyl]phenyl]sulfide

The title compound was prepared by the procedures described in Example 83 substituting 3,4-dimethylthiophenol with 3-aminothiophenol. Yellow solid (2.9 mg, 5.6%); 1H -NMR ($CDCl_3$, 500 MHz) δ 2.20 (s, 3H), 3.60 (br, m, 2H), 3.77 (br, m, 6H), 10 4.03 (br, s, 2H), 6.85 (dd, J = 4, 16 Hz, 1H), 6.90 (m, 3H), 7.04 (d, J = 17 Hz, 1H), 7.30 (t, J = 16 Hz, 1H), 7.52 (d, J = 17 Hz, 1H), 7.68(d, J = 31 Hz, 1H), 8.44 (d, J = 4 Hz, 1H). MS (APCI) (M+H) $^+$ at m/z 427. FAB High Resolution MS calculated m/z for $C_{21}H_{23}N_4O_4S$ (M+H) $^+$: 427.1440. Observed m/z: 427.1440.

Example 139

15 (4-Aminophenyl)[2-nitro-4-(E-((4-acetyl)piperazin-1-yl)carbonyl)ethenyl]phenyl]sulfide

The title compound was prepared by the procedures described in Example 83 substituting 3,4-dimethylthiophenol with 4-aminothiophenol. Yellow solid (2.5 mg, 4.9%); 1H -NMR ($CDCl_3$, 500 MHz) δ 2.19 (s, 3H), 3.58 (br, m, 2H), 3.76 (br, m, 6H), 20 4.03 (br, s, 2H), 6.80 (m, 1H), 6.93 (m, 3H), 7.37 (m, 1H), 7.46 (d, J = 17 Hz, 1H), 7.67 (d, J = 31 Hz, 1H), 8.43 (d, J = 3 Hz, 1H). MS (APCI) (M+H) $^+$ at m/z 427. FAB

High Resolution MS calculated m/z for $C_{21}H_{23}N_4O_4S$ ($M+H$)⁺: 427.1440. Observed m/z : 427.1441.

Example 140

5 (2,4-Dimethylphenyl)[2- nitro-4-(*E*-((4-acetyl)piperazin-1-yl)carbonyl)ethenyl]phenyl]sulfide

The title compound was prepared by the procedures described in Example 83 substituting 3,4-dimethylthiophenol with 2,4-dimethylthiophenol. Yellow solid (40 mg, 76%); ¹H-NMR (CDCl₃, 400 MHz) δ 1.54 (br, s, 2H), 2.14 (s, 3H), 3.53 (br, m, 10 2H), 3.71 (br, m, 6H), 6.58 (d, *J* = 21 Hz, 1H), 6.76 (d, *J* = 38 Hz, 1H), 7.03 (m, 1H), 7.09 (m, 1H), 7.28 (br, d, *J* = 19 Hz, 1H), 7.33 (d, *J* = 20 Hz, 1H), 7.51 (d, *J* = 38 Hz, 1H), 8.30 (d, *J* = 5 Hz, 1H). MS (APCI) ($M+H$)⁺ at m/z 440. FAB High Resolution MS calculated m/z for $C_{23}H_{26}N_3O_4S$ ($M+H$)⁺: 440.1644. Observed m/z : 440.1656.

15 Example 141

(2,5-Dimethylphenyl)[2- nitro-4-(*E*-((4-acetyl)piperazin-1-yl)carbonyl)ethenyl]phenyl]sulfide

The title compound was prepared by the procedures described in Example 83 substituting 3,4-dimethylthiophenol with 2,5-dimethylthiophenol. Yellow solid (34 20 mg, 64%); ¹H-NMR (CDCl₃, 400 MHz) δ 2.07 (s, 3H), 2.23 (s, 3H), 2.28 (s, 3H), 3.46 (br, m, 2H), 3.64 (br, m, 6H), 6.65 (d, *J* = 21 Hz, 1H), 6.81 (d, *J* = 39 Hz, 1H), 7.19 (m, 2H), 7.34 (m, 2H), 7.56 (d, *J* = 38 Hz, 1H), 8.35 (d, *J* = 5 Hz, 1H). MS (APCI)

(M+H)⁺ at m/z 440. FAB High Resolution MS calculated m/z for C₂₃H₂₆N₃O₄S

(M+H)⁺: 440.1644. Observed m/z: 440.1656.

Example 142

5 (4-Methoxyphenyl)[2-nitro-4-(E-((4-acetylpiperazin-1-yl)carbonyl)ethenyl)phenyl]sulfide

The title compound was prepared by the procedures described in Example 83 substituting 3,4-dimethylthiophenol with 4-methoxythiophenol. Yellow solid (44 mg, 83%); ¹H-NMR (CDCl₃, 400 MHz) δ 2.09 (s, 3H), 3.48 (br, m, 2H), 3.66 (br, m, 6H), 10 3.83 (s, 3H), 6.79 (d, J = 22 Hz, 1H), 6.83 (d, J = 40 Hz, 1H), 6.95 (m, 1H), 6.98 (m, 1H), 7.37 (br, d, J = 20 Hz, 1H), 7.43 (m, 1H), 7.46 (m, 1H), 7.58 (d, J = 38 Hz, 1H), 8.35 (d, J = 4 Hz, 1H). MS (APCI) (M+H)⁺ at m/z 442. FAB High Resolution MS calculated m/z for C₂₂H₂₄N₃O₅S (M+H)⁺: 442.1437. Observed m/z: 442.1434.

Example 143

15 (3-Chlorophenyl)[2-nitro-4-(E-((4-acetylpiperazin-1-yl)carbonyl)ethenyl)phenyl]sulfide

The title compound was prepared by the procedures described in Example 83 substituting 3,4-dimethylthiophenol with 3-chlorothiophenol. Yellow solid (43 mg, 80%); ¹H-NMR (CDCl₃, 400 MHz) δ 2.23 (s, 3H), 3.62 (br, m, 2H), 3.80 (br, m, 6H), 20 6.97 (d, J = 21 Hz, 1H), 6.99 (d, J = 39 Hz, 1H), 7.28 (d, J = 19 Hz, 1H), 7.57 (m, 3H), 7.675 (t, J = 4 Hz, 1H), 7.73 (d, J = 39 Hz, 1H), 8.48 (d, J = 4 Hz, 1H). FAB

High Resolution MS calculated m/z for $C_{21}H_{21}N_3O_4ClS$ ($M+H$)⁺: 446.0941. Observed m/z : 446.0953.

Example 144

5 (2-Chloro, 4,5-diaminophenyl)[2-chloro-4-(*E*-((4-acetyl)piperazin-1-yl)carbonyl)ethenyl) phenyl] sulfide

Example 144A

10 (2-Chloro, 4-nitro, 5-aminophenyl)[2-chloro-4-(*E*-((4-acetyl)piperazin-1-yl)carbonyl)ethenyl) phenyl] sulfide

The title compound was prepared by the procedures described in Example 65B substituting 2,3-dichlorobenzaldehyde with 4,5-dichloro-2-nitroaniline.

Example 144B

15 (2-Chloro, 4,5-diaminophenyl)[2-chloro-4-(*E*-((4-acetyl)piperazin-1-yl)carbonyl)ethenyl) phenyl] sulfide

To a stirred solution of nitrobenzene from Example 144A (170 mg, 0.34 mmol) in 2 mL of EtOH was added $SnCl_2$ (325 mg, 1.72 mmol). The mixture was then refluxed under nitrogen atmosphere for 2 h. The reaction was allowed to cool
20 down to ambient temperature, quenched with sat. $NaHCO_3$, extracted with EtOAc(2×20 mL). The combined organic layer was washed with brine, dried over Na_2SO_4 , concentrated in vacuo. The residue was then purified on Gilson preparative

HPLC as described in Example 38B to give the title compound (70 mg, 44% yield) as a light yellow solid. ¹H NMR (d⁶-DMSO, 300 MHz) δ 2.04 (s, 3H), 3.42-3.80 (m, 8H), 4.84 (s, 2H), 5.32 (s, 2H), 6.51 (d, *J* = 8.4 Hz, 1H), 6.78 (d, *J* = 8.4 Hz, 2H), 7.26 (d, *J* = 15.6 Hz, 1H), 7.41 (d, *J* = 15.6 Hz, 1H), 7.48 (d, *J* = 8.4 Hz, 1H), 7.95 (d, *J* = 1.8 Hz, 1H). MS (APCI⁺) (M+H)⁺ at *m/z* 465, 467, 469, 471.

Example 145

(3,4-Diaminophenyl)[2-chloro-4-(*E*-((4-acetyl)piperazin-1-yl)carbonyl)ethenyl]phenyl sulfide

The title compound was prepared by the procedures described in Example 144, substituting 4,5-dichloronitroaniline with 5-chloronitroaniline, resulting in a light brown solid. ¹H NMR (d⁶-DMSO, 300 MHz) δ 2.04 (s, 3H), 3.31-3.80 (m, 8H), 4.75 (s, 2H), 5.01 (s, 2H), 6.61 (t, *J* = 4.2 Hz, 3H), 6.68 (s, 1H), 7.26 (d, *J* = 15.6 Hz, 1H), 7.40 (d, *J* = 15.6 Hz, 1H), 7.46 (d, *J* = 8.4 Hz, 1H), 7.94 (s, 1H). MS (APCI⁺) (M+H)⁺ at *m/z* 431, 433.

Example 146

(6-Chlorobenzimidazol-2-on-5-yl)[2-chloro-4-(*E*-((4-acetyl)piperazin-1-yl)carbonyl)ethenyl]phenyl sulfide

A mixture of dianiline from Example 144 (35 mg, 0.075 mmol) and CDI (13 mg, 0.075 mmol) in THF was stirred at ambient temperature for one day. Solvent was then removed under reduced pressure. The crude product then purified on a Gilson

preparative HPLC as described in Example 38B to give the title compound (12 mg, 32% yield) as a white solid. ¹H NMR (d⁶-DMSO, 300 MHz) δ 2.04 (s, 3H), 3.40-3.80 (m, 8H), 6.63 (d, *J* = 8.4 Hz, 1H), 7.11 (d, *J* = 2.4 Hz, 1H), 7.12 (s, 1H), 7.23 (s, 1H), 7.32 (d, *J* = 15.6 Hz, 1H), 7.43 (d, *J* = 15.6 Hz, 1H), 7.50 (d, *J* = 8.4 Hz, 1H), 8.03 (br s, 1H). MS (APCI⁺) (M-CO+H)⁺ at *m/z* 465, 467.

Example 147

(1-Methylindol-7-yl)[2-chloro-4-(*E*-((4-acetyl)piperazin-1-yl)carbonyl)ethenyl)phenyl sulfide

The title compound was prepared by the procedures described in 85, substituting 5-iodoindole with *N*-methyl-7-bromoindole, giving a light brown solid. ¹H NMR (CDCl₃, 300 MHz) δ 2.14 (s, 3H), 3.47-3.56 (m, 2H), 3.56-3.83 (m, 6H), 3.96 (s, 3H), 6.42 (d, *J* = 8.4 Hz, 1H), 6.55 (d, *J* = 3.6 Hz, 1H), 6.76 (d, *J* = 15.6 Hz, 1H), 6.99 (d, *J* = 3.6 Hz, 1H), 7.09 (dd, *J* = 2.1, 8.4 Hz, 1H), 7.15 (t, *J* = 7.65 Hz, 1H), 7.42 (dd, *J* = 0.9, 7.5 Hz, 1H), 7.53 (d, *J* = 1.8 Hz, 1H), 7.55 (dd, *J* = 15.6 Hz, 1H), 7.77 (dd, *J* = 0.9, 7.5 Hz, 1H). MS (APCI⁺) (M+H)⁺ at *m/z* 454, 456.

Example 148

(2-Hydroxy, 4-aminophenyl)[2-chloro-4-(*E*-((4-acetyl)piperazin-1-yl)carbonyl)ethenyl)phenyl sulfide

The title compound was prepared by the procedures described in Example 144, substituting 4,5-dichloronitroaniline with 5-chloronitrophenol, giving a light brown

solid. ^1H NMR ($\text{d}^6\text{-DMSO}$, 300 MHz) δ 2.04 (s, 3H), 3.41-3.80 (m, 8H), 5.09 (s, 2H), 6.61 (d, $J = 8.4$ Hz, 1H), 6.70 (d, $J = 7.8$ Hz, 1H), 6.79 (s, 1H), 6.80 (dd, $J = 2.1, 7.8$ Hz, 1H), 7.26 (d, $J = 15.6$ Hz, 1H), 7.40 (d, $J = 15.6$ Hz, 1H), 7.46 (d, $J = 8.4$ Hz, 1H), 7.94 (br s, 1H). MS (APCI) $(\text{M}+\text{H})^+$ at m/z 432, 434.

5

Example 149

(2-Isopropylphenyl)[2-nitro-4-(*E*-((4-methylpiperazin-1-yl)carbonyl)ethenyl) phenyl]

sulfide

Prepared according to the procedures of Example 71, giving a yellow solid.

- 10 ^1H NMR ($\text{DMSO-}d_6$, 300MHz) δ 1.14 (d, $J = 7.0$ Hz, 6H); 2.19 (s, 3H); 2.25-2.36 (br m, 4H); 3.30-3.40 (m, 1H); 3.51-3.72 (br m, 4H); 6.63 (d, $J = 8.5$ Hz, 1H); 7.24-7.63 (m, 6H); 7.88-7.92 (dd, $J = 8.8, 1.8$ Hz, 1H); 8.64 (d, $J = 1.8$ Hz, 1H). MS (APCI) $(\text{M}+\text{H})^+$ at m/z 426. Anal calcd for $\text{C}_{23}\text{H}_{27}\text{N}_3\text{S}_2\text{O}_3 \cdot 0.26\text{H}_2\text{O}$: C, 64.19; H, 6.45; N, 9.76. Found: C, 64.21; H, 6.59; N, 9.70.

15

Example 150

(2-Isopropylphenyl)[2-nitro-4-(*E*-((4-(pyridine-2-carbonyl)piperazin-1-yl)carbonyl)ethenyl) phenyl] sulfide

Prepared according to the procedures of Example 71, giving a yellow solid.

- 20 ^1H NMR ($\text{DMSO-}d_6$, 300MHz) δ 1.14 (d, $J = 6.8$ Hz, 6H); 3.30-3.40 (m, 1H); 3.51-3.83 (br m, 8H); 6.61-6.66 (br m, 1H); 7.30-7.65 (m, 8H); 7.83-7.97 (m, 2H); 8.57-

8.67 (m, 2H). MS (APCI) (M+H)⁺ at m/z 517. Anal calcd for C₂₈H₂₈N₄S₁O₄·0.45H₂O: C, 64.07; H, 5.53; N, 10.67. Found: C, 64.04; H, 5.77; N, 10.97.

Example 151

5 (2-Isopropylphenyl)[2-nitro-4-(E-((4-(pyridine-3-carbonyl)piperazin-1-yl)carbonyl)ethenyl) phenyl] sulfide

Prepared according to the procedures of Example 71, giving a yellow solid.

¹H NMR (DMSO-d₆, 300MHz) δ 1.14 (d, J = 7.0 Hz, 6H); 3.30-3.40 (m, 1H); 3.52-3.87 (br m, 8H); 6.64 (d, J = 8.5 Hz, 1H); 7.30-7.64 (m, 7H); 7.83-7.95 (m, 2H); 8.61-8.70 (m, 3H). MS (APCI) (M+H)⁺ at m/z 517. Anal calcd for C₂₈H₂₈N₄S₁O₄·0.42H₂O: C, 64.16; H, 5.55; N, 10.69. Found: C, 64.18; H, 5.64; N, 10.59.

Example 152

15 (2-Isopropylphenyl)[2-nitro-4-(E-((2-carbomethoxy-4-methoxycarbonylpiperazin-1-yl)carbonyl)ethenyl) phenyl] sulfide

Prepared according to the procedures of Example 71, giving a yellow solid.

¹H NMR (DMSO-d₆, 300MHz) δ 1.14 (d, J = 7.1 Hz, 6H); 2.70-3.95 (br m, 4H); 3.30-3.40 (m, 1H); 3.61, 3.61 (s, s, 3H); 3.65, 3.67 (s, s, 3H); 4.16-4.50 (br m, 2H); 5.08-5.39 (br m, 1H); 6.64 (dd, J = 8.5, 5.1 Hz, 1H); 7.30-7.63 (m, 6H); 7.83-7.94 (m, 1H); 8.62-8.67 (m, 1H). MS (APCI) (M+H)⁺ at m/z 528. Anal calcd for C₂₆H₂₅N₃S₁O₇·0.19C₆H₁₄: C, 59.94; H, 5.87; N, 7.72. Found: C, 59.87; H, 5.94; N, 7.59.

Example 153

(2-Isopropylphenyl)[2-nitro-4-(*E*-(2-carboxy-4-methoxycarbonylpiperazin-1-yl)carbonyl)ethenyl] phenyl] sulfide

5 Prepared according to the procedures of Example 71, giving a yellow solid.
 ¹H NMR (DMSO-*d*₆, 300MHz) δ 1.14 (d, J = 6.8 Hz, 6H); 2.70-3.95 (br m, 4H); 3.30-3.40 (m, 1H); 3.61, 3.61 (s, s, 3H); 4.16-4.51 (br m, 2H); 5.01-5.28 (br m, 1H); 6.61-6.66 (m, 1H); 7.30-7.63 (m, 6H); 7.83-7.94 (m, 1H); 8.66 (br s, 1H). MS (APCI) (M-H)⁺ at m/z 512.

10

Example 154

(2-Isopropylphenyl)[2-nitro-4-(*E*-(3-carbomethoxy-4-methylpiperazin-1-yl)carbonyl)ethenyl] phenyl] sulfide

 Prepared according to the procedures of Example 71, giving a yellow solid.
15 ¹H NMR (DMSO-*d*₆, 300MHz) δ 1.14 (d, J = 7.0 Hz, 6H); 2.25, 2.26 (s, s, 3H); 2.20-3.98 (br m, 8H); 3.57, 3.63 (s, s, 3H); 6.63 (d, J = 8.5 Hz, 1H); 7.30-7.63 (m, 6H); 7.91 (dd, J = 8.5, 1.5 Hz, 1H); 8.60-8.68 (br m, 1H). MS (APCI) (M-H)⁺ at m/z 484.

Example 155

20 (2-Ethoxyphenyl)-[2-chloro-4-(*E*-(3-carboxypiperidin-1-yl)carbonyl)ethenyl]phenyl]
 sulfide

The compound of Example 137 was hydrolyzed using an excess of aqueous 10% NaOH in methanol, stirring overnight. The reaction mixture was concentrated in vacuo, water was added, and the solution was extracted with ether. The mixture was acidified; the resultant solid was collected by filtration and dried overnight in a vacuum oven, giving a white solid, m.p. 166-171C. ¹H-NMR (DMSO 300 MHz) δ 1.17 (t, J=7Hz, 3H), broad peaks totaling 9 protons at 1.32-1.48, 1.51-1.78, 1.90-2.04, 2.25-2.50, 2.80-2.90, 2.95-3.17, 3.45-3.51, 3.95-4.19, 4.41-4.51, 4.06 (q, J=7Hz, 1H), 6.80 (d, J=9Hz, 1H), 7.01 (t, J=7Hz, 1H), 7.15 (d, J=8Hz, 1H), 7.26-7.40 (m, 2H), 7.40-7.48 (m, 1H), 7.51 (dd, J=9Hz, 2Hz, 1H), 7.99 (d, J=9Hz, 1H). Anal. Calcd. for C₂₃H₂₄ClNO₄S: C, 61.94; H, 5.42; N, 3.14. Found: C, 61.75; H, 5.65; N, 3.15. The resultant acid (303 mg, 0.631 mmol) was dissolved in 3 ml MeOH. A KOH solution (38 mg, 0.595 mmol, of 87.6% KOH) in 1 ml MeOH was added. The resulting solution was concentrated in vacuo, and 5 ml. ether was added. The mixture was stirred for one hour to form a powder, which was filtered and dried in the vacuum oven at 60C to yield 307 mg of a solid, water soluble product.

Example 155

(2-Ethoxyphenyl)-[2-chloro-4(E-[(3-carboxypiperidin-1-yl)carbonyl]ethenyl)phenyl]

sulfide

The compound of Example 137 was hydrolyzed using an excess of aqueous 10% NaOH in methanol, stirring overnight. The reaction mixture was concentrated *in vacuo*, water was added, and the solution was extracted with ether, giving a white

solid, m.p. 166-171. ^1H NMR (DMSO, 300 MHz) δ 1.17 (t, $J=7\text{Hz}$, 3H), broad peaks totaling 9 protons at 1.32-1.48, 1.51-1.78, 1.90-2.04, 2.25-2.50, 2.80-2.90, 2.95-3.17, 3.45-3.51, 3.95-4.19, 4.41-4.51, 4.06 (q, $J=7\text{Hz}$, 1H), 6.80 (d, $J=9\text{Hz}$, 1H), 7.01 (t, $J=7\text{Hz}$, 1H), 7.15 (d, $J=8\text{Hz}$, 1H), 7.26-7.40 (m, 2H), 7.40-7.48 (m, 1H), 7.51 (dd,

- 5 $J=9\text{Hz}$, 2H, 1H), 7.99 (d, $J=9\text{Hz}$, 1H). Anal. Calcd. for $\text{C}_{23}\text{H}_{24}\text{ClNO}_4\text{S}$: C, 61.94; H, 5.42; N, 3.14. Found: C, 61.75; H, 5.65; N, 3.15. The resultant acid (303 mg, 0.631 mmol) was dissolved in 3 ml MeOH. A KOH solution (38 mg, 0.595 mmol, of 87.6% KOH) in 1 ml MeOH was added. The resulting solution was concentrated in vacuo, and 5 ml. ether was added. The mixture was stirred for one hour to form a
- 10 powder, which was filtered and dried in the vacuum oven at 60°C to yield 307 mg of a solid, water soluble product.

Example 156

(2-Ethoxyphenyl)-[2-chloro-4(E-[(2-ethoxycarbonyl)piperidin-1-yl]carbonyl]ethenyl
phenyl]sulfide

- 15 The title compound was prepared according to the procedures of Example 97.
- ^1H NMR (CDCl_3 , 300 MHz) δ 1.24 (t, $J=7\text{Hz}$, 3H), 1.28 (t, $J=7\text{Hz}$, 3H), broad peaks totaling 9 protons at 1.35-1.55, 1.65-1.80, 2.25-2.38, 3.33-3.45, 3.95-4.05, 4.15-4.28, 4.60-4.80, 5.44-5.50, 4.05 (q, $J=7\text{Hz}$, 2H), 4.20 (q, $J=7\text{Hz}$, 2H), 6.80-6.98 (m, 4H),
- 20 7.12-7.20 (m, 1H) 7.35-7.43 (m, 2H), 7.50-7.58 (m, 2H). Anal. Calcd. for $\text{C}_{25}\text{H}_{28}\text{ClNO}_4\text{S}$: C, 63.35; H, 5.95; N, 2.95. Found: C, 63.51; H, 6.22; N, 2.61.

Example 157

(2-Ethoxyphenyl)[2-trifluoromethyl-4-(*E*-[(1-(*tert*-butoxycarbonyl)-4-hydroxypyrrolidin-3-ylamino)carbonyl]ethenyl) phenyl] sulfide

The title compound was prepared according to the procedures of Example 1.

- 5 ¹H NMR (CDCl₃, 300 MHz) δ 7.76 (s, 1H), 7.60 (d, 1H, J = 15.1 Hz), 7.46 (dd, 1H, J = 1.7, 7.5 Hz), 7.38 (m, 2H), 7.01 (d, 1H, J = 15.4 Hz), 6.98 (d, 1H, J = 7.8 Hz), 6.93 (d, 1H, J = 8.3 Hz), 6.42 (d, 1H, J = 15.0 Hz), 4.30 (br, 2H), 3.98 (q, 2H, J = 7.0 Hz), 3.87 (m, 1H), 3.71 (m, 1H), 3.33 (br, 2H), 1.47 (s, 9H), 1.17 (t, 3H, J = 7.0 Hz). MS (ESI) *m/z* -551, -1103. Anal. Calcd for C₂₇H₃₁F₃N₂O₅S · 0.61 EtOAc: C, 58.32; H, 5.96; N, 4.62. Found: C, 58.07; H, 5.88; N, 4.76.
- 10

Example 158

(2-Ethoxyphenyl)-[2-chloro-4(*E*-[(2-carboxypiperidin-1-yl)carbonyl]ethenyl)phenyl] sulfide

- 15 The compound of Example 156 was hydrolyzed, and the salt formed, according to the procedures of Example 155. m.p. 170-171°C. ¹H-NMR (DMSO 300 MHz) δ 1.16 (t, J=7Hz, 3H), broad peaks totaling 9 protons at 1.20-1.49, 1.51-1.75, 2.10-2.27, 2.55-2.65, 3.10-3.21, 4.20-4.29, 4.35-4.45, 5.13-5.25, 4.05 (q, J=7Hz, 2H), 6.80 (d, J=9Hz, 1H), 6.97-7.07 (m, 1H), 7.15 (d, J=9Hz, 1H), 7.29-7.57 (m, 5H), 8.02
- 20 (s, 1H). Anal. Calcd. for C₂₃H₂₄ClNO₄S: C, 61.94; H, 5.42; N, 3.14. Found: C, 61.91; H, 5.48; N, 2.90.

Example 159

(2-Ethoxyphenyl)[2-trifluoromethyl-4-(*E*-(((pyrrol-3-yl)carbonyl)ethenyl)phenyl] sulfide

The title compound was prepared according to the procedures of Example 1.

¹H NMR (CDCl₃, 300 MHz) δ 7.81 (s, 1H), 7.68 (d, 1H, J = 15.4 Hz), 7.35-7.47 (m, 3H), 7.04 (d, 1H, J = 8.4 Hz), 6.97 (dd, 1H, J = 1.3, 7.5 Hz), 6.91 (d, 1H, J = 8.5 Hz), 6.70 (d, 1H, J = 15.4 Hz), 5.94 (m, 1H), 5.85 (m, 1H), 4.47 (br, 2H), 4.38 (br, 2H), 3.98 (q, 2H, J = 7.0 Hz), 1.19 (t, 3H, J = 7.0 Hz). MS (ESI) *m/z* 420, 839, 861.

Example 160

(2-Ethoxyphenyl)[2-trifluoromethyl-4-(*E*-((3-(pyrrolidin-2-on-1-yl)prop-1-ylamino)carbonyl)ethenyl)phenyl]sulfide

The title compound was prepared according to the procedures of Example 1.

¹H NMR (CDCl₃, 300 MHz) δ 7.78 (s, 1H), 7.54 (d, 1H, J = 15.8 Hz), 7.42 (dd, 1H, J = 1.7, 7.5 Hz), 7.34-7.39 (m, 2H), 7.13 (br, 1H), 7.03 (d, 1H, J = 8.5), 6.97 (dd, 1H, J = 1.1, 7.7 Hz), 6.91 (d, 1H, J = 8.1 Hz), 6.46 (d, 1H, J = 15.8 Hz), 3.98 (q, 2H, J = 7.0 Hz), 3.43 (m, 4H), 3.34 (q, 2H, J = 6.0 Hz), 2.45 (t, 2H, J = 8.1 Hz), 2.08 (m, 2H), 1.75 (m, 2H), 1.18 (t, 3H, J = 7.0 Hz). MS (ESI) *m/z* 493, 515, 985, 1007.

Example 161

(2-Ethoxyphenyl)[2-trifluoromethyl-4-(*E*-((4-acetyl)piperazin-1-yl)carbonyl)ethenyl)phenyl] sulfide

001220.56214560

The title compound was prepared according to the procedures of Example 1.

¹H NMR (CDCl₃, 300 MHz) δ 7.79 (s, 1H), 7.62 (d, 1H, J = 15.6 Hz), 7.44 (dd, 1H, J = 1.7, 7.5 Hz), 7.38 (m, 2H), 7.04 (d, 1H, J = 8.1), 6.97 (dd, 1H, J = 1.4, 7.5 Hz), 6.92 (d, 1H, J = 8.1 Hz), 6.84 (d, 1H, J = 15.6 Hz), 3.98 (q, 2H, J = 7.0 Hz), 3.63-78 (m, 5 6H), 3.53 (m, 2H), 2.14 (s, 3H), 1.19 (t, 3H, J = 7.0 Hz). MS (ESI) *m/z* 479, 501, 957, 979.

Example 162

(2-Ethoxyphenyl)[2-trifluoromethyl-4-(E-((4-(ethoxycarbonyl)piperazin-1-yl)carbonyl)ethenyl) phenyl] sulfide

10

The title compound was prepared according to the procedures of Example 71.

¹H NMR (CDCl₃, 300 MHz) δ 7.79 (d, 1H, J = 1.7 Hz), 7.63 (d, 1H, J = 15.3 Hz), 7.43 (dd, 1H, J = 1.7, 7.7 Hz), 7.38 (m, 2H), 7.04 (d, 1H, J = 8.5), 6.97 (dd, 1H, J = 1.4, 7.5 Hz), 6.92 (d, 1H, J = 8.1 Hz), 6.84 (d, 1H, J = 15.3 Hz), 4.18 (q, 2H, J = 7.1 15 Hz), 3.98 (q, 2H, J = 6.9 Hz), 3.68 (m, 4H), 3.53 (m, 4H), 1.29 (t, 3H, J = 7.1 Hz) 1.19 (t, 3H, J = 6.9 Hz). MS (ESI) *m/z* 509, 531, 1017, 1039.

Example 163

(2-Ethoxyphenyl)[2-trifluoromethyl-4-(E-((4-(2-furylcarbonyl)piperazin-1-yl)carbonyl)ethenyl) phenyl] sulfide

20

The title compound was prepared according to the procedures of Example 71.

¹H NMR (CDCl₃, 300 MHz) δ 7.80 (d, 1H, J = 1.5 Hz), 7.66 (d, 1H, J = 15.4 Hz),

7.52 (s, 1H), 7.45 (dd, 1H, $J = 1.6, 7.5$ Hz), 7.40 (m, 2H), 7.08 (d, 1H, $J = 4.0$ Hz), 7.04 (d, 1H, $J = 8.1$), 6.98 (dd, 1H, $J = 1.1, 7.3$ Hz), 6.93 (d, 1H, $J = 8.5$ Hz), 6.88 (d, 1H, $J = 15.4$ Hz), 6.52 (dd, 1H, $J = 1.6, 3.5$ Hz), 3.98 (q, 2H, $J = 7.0$ Hz), 3.73-3.90 (m, 8H), 1.19 (t, 3H, $J = 7.0$ Hz). MS (ESI) m/z 531, 553, 1061, 1083.

5

Example 164

(2-Ethoxyphenyl)-[2-chloro-4(*E*-[(3-ethoxycarbonylpiperidin-1-yl)carbonyl]ethenyl)phenyl]sulfide

The title compound was prepared according to the procedures of Example 97.

- 10 $^1\text{H-NMR}$ (CDCl_3) δ 1.25 (t, $J=7\text{Hz}$, 6H), broad peaks totaling 9 protons at 1.65-1.80, 1.95-2.04, 2.51-2.63, 2.90-3.00, 3.15-3.30, 2.95-4.05, 4.42-4.55, 4.14 (q, $J=7\text{Hz}$, 2H), 4.15 (q, $J=7\text{Hz}$, 2H), 6.82 (d, $J=15$ Hz, 1H), 6.84 (d, $J=9\text{Hz}$, 1H), 6.93-6.99 (m, 2H), 7.17 (dd, $J=9\text{Hz}$, 2Hz, 1H), 7.34-7.41 (m, 2H), 7.52 (d, $J=15\text{Hz}$, 1H), 7.55 (d, $J=2\text{Hz}$, 1H). Anal. Calcd. for $\text{C}_{25}\text{H}_{28}\text{ClNO}_4\text{S}$: C, 63.35; H, 5.95; N, 2.95. Found: C, 63.09; H, 6.24; N, 2.77.
- 15

Example 165

(2-Ethoxyphenyl)-[2-chloro-4(*E*-[(4-carboxypiperidin-1-yl)carbonyl]ethenyl)phenyl]sulfide

- 20 The compound of Example 164 was hydrolyzed, and the salt formed, according to the procedures of Example 155. m.p. 165-166C. $^1\text{H-NMR}$ (DMSO 300 MHz) δ 1.25 (t, $J=7\text{Hz}$, 3H), 1.35-1.58 (m, 2H), 1.80-1.95 (m, 2H), 2.50-2.60 (m,

1H), 1.78-1.91 (m, 1H), 3.13-3.24 (m, 1H), 4.05 (q, J=7Hz, 2H), 4.12-4.35 (m, 2H), 6.80 (d, J=9Hz, 1H), 6.96-7.05 (t, J=8 Hz, 1H), 7.15 (d, J=9Hz, 1H), 7.28-7.48 (m, 4H), 7.51 (dd, J=9Hz, 2Hz, 1H), 8.00 (d, J=2Hz).

5

Example 166

(Benzodioxan-6-yl)[2-chloro-4-(E-((4-acetyl)piperazin-1-yl)carbonyl)ethenyl] phenyl]
sulfide

The title compound was prepared by the procedures described in Example 85, substituting 5-iodoindole with 6-iodobenzenedioxane, giving a white solid. ¹H NMR

10 (CDCl₃, 300 MHz) δ 2.14 (s, 3H), 3.44-3.57 (m, 2H), 3.57-3.86 (m, 6H), 4.25-4.35 (m, 4H), 6.75 (d, J = 8.4 Hz, 1H), 6.78 (d, J = 15.6 Hz, 1H), 6.93 (d, J = 8.4 Hz, 1H), 7.03 (dd, J = 2.1, 8.4 Hz, 1H), 7.08 (d, J = 2.1 Hz, 1H), 7.18 (dd, J = 2.1, 8.4 Hz, 1H), 7.51 (d, J = 2.1 Hz, 1H), 7.57 (d, J = 15.6 Hz, 1H). MS (APCI⁺) (M+H)⁺ at m/z 459, 461.

15

Example 167

(2-Isopropylphenyl)[2-nitro-4-(E-((4-ethoxycarbonyl)piperazin-1-yl)carbonyl)ethenyl]
phenyl] sulfide

Prepared according to the procedures of Example 71, giving a yellow solid.

20 ¹H NMR (DMSO-d₆, 300MHz) δ 1.14 (d, J = 7.0 Hz, 6H); 1.19 (t, J = 7.0Hz, 3H); 3.30-3.40 (m, 1H); 3.30-3.73 (br m, 8H); 4.06 (q, J = 7.0 Hz, 2H); 6.64 (d, J = 8.5 Hz, 1H); 7.32-7.63 (m, 6H); 7.90 (dd, J = 8.8, 1.8 Hz, 1H); 8.65 (d, J = 1.8 Hz, 1H). MS

(APCI) (M+H)⁺ at m/z 484. Anal calcd for C₂₅H₂₉N₃O₅: C, 62.09; H, 6.04; N, 8.69.

Found: C, 61.89; H, 6.13; N, 8.51.

Example 168

5 (2-Isopropylphenyl)[2-nitro-4-(*E*-((4-isopropoxycarbonylpiperazin-1-yl)carbonyl)ethenyl) phenyl] sulfide

Prepared according to the procedures of Example 71, giving a yellow solid.

¹H NMR (DMSO-d₆, 300MHz) δ 1.14 (d, J = 6.8 Hz, 6H); 1.20 (d, J = 6.4 Hz, 3H); 3.30-3.40 (m, 1H); 3.32-3.73 (br m, 8H); 4.79 (hept, J = 6.1 Hz, 2H); 6.64 (d, J = 8.5 Hz, 1H); 7.32-7.63 (m, 6H); 7.89 (dd, J = 8.5, 1.7 Hz, 1H); 8.64 (d, J = 1.7 Hz, 1H).
10 MS (APCI) (M+H)⁺ at m/z 498. Anal calcd for C₂₆H₃₁N₃O₅: C, 62.76; H, 6.28; N, 8.44. Found: C, 62.57; H, 6.43; N, 8.33.

Example 169

15 (2-Isopropylphenyl)[2-nitro-4-(*E*-((4-isobutoxycarbonylpiperazin-1-yl)carbonyl)ethenyl) phenyl] sulfide

Prepared according to the procedures of Example 71, giving a yellow solid.

¹H NMR (DMSO-d₆, 300MHz) δ 0.90 (d, J = 6.6 Hz, 6H); 1.14 (d, J = 7.0 Hz, 6H); 1.88 (hept, J = 6.6 Hz, 1H); 3.30-3.40 (m, 1H); 3.30-3.73 (br m, 8H); 3.81 (d, J = 6.3 Hz, 2H); 6.64 (d, J = 8.5 Hz, 1H); 7.32-7.63 (m, 6H); 7.90 (dd, J = 8.5, 1.5 Hz, 1H); 8.65 (d, J = 1.5 Hz, 1H). MS (APCI) (M+H)⁺ at m/z 512. Anal calcd for C₂₇H₃₃N₃O₅: C, 63.38; H, 6.50; N, 8.21. Found: C, 63.15; H, 6.55; N, 8.13.

Example 170

(2-Isopropylphenyl)[2-nitro-4-(*E*-((4-((1-propen-2-oxyl)carbonyl)piperazin-1-yl)carbonyl)ethenyl) phenyl] sulfide

5 Prepared according to the procedures of Example 71, giving a yellow solid.

¹H NMR (DMSO-d₆, 300MHz) δ 1.14 (d, J = 6.8 Hz, 6H); 1.88 (s, 3H); 3.30-3.40 (m, 1H); 3.30-3.78 (br m, 8H); 4.65 (s, 1H); 4.69 (m, 1H); 6.64 (d, J = 8.5 Hz, 1H); 7.32-7.63 (m, 6H); 7.90 (dd, J = 8.5, 1.5 Hz, 1H); 8.65 (d, J = 1.5 Hz, 1H). MS (APCI) (M+NH₄)⁺ at m/z 513. Anal calcd for C₂₆H₂₉N₃O₄: C, 63.01; H, 5.90; N, 8.48.

10 Found: C, 62.98; H, 6.06; N, 8.27.

Example 171

(2-Isopropylphenyl)[2-nitro-4-(*E*-((4-propionylpiperazin-1-yl)carbonyl)ethenyl) phenyl] sulfide

15 Prepared according to the procedures of Example 71, giving a yellow solid.

¹H NMR (DMSO-d₆, 300MHz) δ 1.00 (t, J = 7.3 Hz, 3H); 1.14 (d, J = 7.0 Hz, 6H); 2.35 (q, J = 7.5 Hz, 2H); 3.30-3.40 (m, 1H); 3.41-3.76 (br m, 8H); 6.64 (d, J = 8.5 Hz, 1H); 7.32-7.63 (m, 6H); 7.90 (dd, J = 8.5, 1.5 Hz, 1H); 8.64 (d, J = 1.5 Hz, 1H). MS (APCI) (M+ NH₄)⁺ at m/z 485. Anal calcd for C₂₃H₂₉N₃O₄: C, 64.22; H, 6.25; N, 8.99.

20 Found: C, 64.04; H, 6.44; N, 8.80.

Example 172

00160-567450

(2-Isopropylphenyl)[2-nitro-4-(E-((4-carboxamidopiperazin-1-yl)carbonyl)ethenyl)phenyl] sulfide

Prepared according to the procedures of Example 71, giving a yellow solid.

¹H NMR (DMSO-d₆, 300MHz) δ 1.14 (d, J = 7.0 Hz, 6H); 3.30-3.40 (m, 1H); 3.30-3.73 (br m, 8H); 6.10 (s, 2H); 6.64 (d, J = 8.5 Hz, 1H); 7.32-7.63 (m, 6H); 7.91 (dd, J = 8.5, 1.8 Hz, 1H); 8.65 (d, J = 1.8 Hz, 1H). MS (APCI) (M+NH₄)⁺ at m/z 470. Anal calcd for C₂₃H₂₈N₆S₂O₄·0.26CH₃COOCH₂CH₃; C, 60.48; H, 5.93; N, 11.73. Found: C, 60.10; H, 5.84; N, 11.90.

Example 173

(2-Isopropylphenyl)[2-nitro-4-(E-((4-methylaminocarbonylpiperazin-1-yl)carbonyl)ethenyl)phenyl] sulfide

Prepared according to the procedures of Example 71, giving a yellow solid.

¹H NMR (DMSO-d₆, 300MHz) δ 1.14 (d, J = 6.8 Hz, 6H); 2.58 (d, J = 4.4 Hz, 3H); 3.30-3.40 (m, 1H); 3.28-3.70 (br m, 8H); 6.52 (q, J = 4.4 Hz, 1H); 6.64 (d, J = 8.5 Hz, 1H); 7.32-7.62 (m, 6H); 7.90 (dd, J = 8.5, 1.8 Hz, 1H); 8.64 (d, J = 1.8 Hz, 1H). MS (APCI) (M+NH₄)⁺ at m/z 486. Anal calcd for C₂₄H₂₈N₆S₂O₄·0.36CH₃COOCH₂CH₃; C, 61.07; H, 6.22; N, 11.19. Found: C, 61.14; H, 6.41; N, 11.19.

Example 174

(2-Isopropylphenyl)[2-nitro-4-(E-((4-(pyrimidin-2-yl)piperazin-1-yl)carbonyl)ethenyl)phenyl] sulfide

Prepared according to the procedures of Example 71, giving a yellow solid.

¹H NMR (DMSO-d₆, 300MHz) δ 1.15 (d, J = 6.6 Hz, 6H); 3.30-3.40 (m, 1H); 3.28-3.85 (br m, 8H); 6.64 (d, J = 8.5 Hz, 1H); 6.68 (d, J = 4.8 Hz, 1H); 7.33-7.63 (m, 6H); 7.92 (dd, J = 8.5, 1.8 Hz, 1H); 8.40 (d, J = 4.8 Hz, 2H); 8.67 (d, J = 1.8 Hz, 1H). MS (APCI) (M+H)⁺ at m/z 490. Anal calcd for C₂₆H₂₇N₃S₁O₃: C, 63.78; H, 5.56; N, 14.30. Found: C, 63.83; H, 5.54; N, 14.11.

Example 175

(2-Isopropylphenyl)[2-nitro-4-(E-((4-hydroxyacetyl)piperazin-1-yl)carbonyl)ethenyl)phenyl sulfide

Prepared according to the procedures of Example 71, giving a yellow solid.

¹H NMR (DMSO-d₆, 300MHz) δ 1.15 (d, J = 6.8 Hz, 6H); 3.30-3.40 (m, 1H); 3.28-3.78 (br m, 8H); 4.12 (d, J = 5.8 Hz, 2H); 4.61-4.69 (br m, 1H); 6.64 (d, J = 8.5 Hz, 1H); 7.33-7.63 (m, 6H); 7.90 (dd, J = 8.5, 1.8 Hz, 1H); 8.65 (d, J = 1.8 Hz, 1H). MS (APCI) (M+H)⁺ at m/z 470. Anal calcd for C₂₄H₂₇N₃O₇·0.38CH₃COOCH₂CH₃: C, 60.93; H, 6.02; N, 8.35. Found: C, 60.95; H, 6.06; N, 8.35.

Example 176

(2-Isopropylphenyl)[2-nitro-4-(E-((4-(pyrazine-2-carbonyl)piperazin-1-yl)carbonyl)ethenyl)phenyl sulfide

Prepared according to the procedures of Example 71, giving a yellow solid.

¹H NMR (DMSO-d₆, 300MHz) δ 1.14 (d, J = 6.6 Hz, 6H); 3.30-3.40 (m, 1H); 3.28-

3.88 (br m, 8H); 6.61-6.66 (br m, 1H); 7.31-7.63 (m, 6H); 7.85-7.96 (br m, 1H); 8.61-8.92 (m, 4H). MS (APCI) (M+H)⁺ at m/z 518. Anal calcd for C₂₇H₂₇N₃O₄·0.24CH₃COOCH₂CH₃: C, 62.34; H, 5.41; N, 13.01. Found: C, 62.23; H, 5.50; N, 13.10.

5

Example 177

(2-Isopropylphenyl)[2-trifluoromethyl-4-(*E*-((2-carboxypyrrol-3-in-1-yl)carbonyl)ethenyl) phenyl] sulfide

The title compound was prepared according to the procedures of Example 71.

10 ¹H NMR (CDCl₃, 300 MHz) δ 7.79 (s, 1H), 7.68 (d, 1H, J = 15.4 Hz), 7.48 (d, 1H, J = 7.4 Hz), 7.45 (m, 2H), 7.38 (d, 1H, J = 8.3 Hz), 7.23 (m, 1H), 6.80 (d, 1H, J = 8.5 Hz), 6.70 (d, 1H, J = 15.4 Hz), 6.04 (m, 1H), 5.88 (m, 1H), 5.31 (m, 1H), 4.60 (m, 1H), 4.50 (m, 1H), 3.76 (s, 3H), 3.50 (m, 1H), 1.22 (d, 6H, J = 7.0 Hz). MS (ESI) m/z 476, 498, 951, 973. Anal. Calcd for C₂₅H₂₄F₃NO₃S · 0.38 EtOAc: C, 62.58; H, 5.35; N, 15 2.75. Found: C, 62.53; H, 5.27; N, 2.76.

Example 178

(2-Isopropylphenyl)[2-nitro-4-(*E*-((3-hydroxymethyl-4-methylpiperazin-1-yl)carbonyl)ethenyl) phenyl] sulfide

20

Prepared according to the procedures of Example 71, giving a yellow solid.

¹H NMR (DMSO-d₆, 300MHz) δ 1.14 (d, J = 6.8 Hz, 6H); 2.22 (s, 3H); 1.82-4.63 (br

m, 9H); 3.30-3.40 (m, 1H); 6.62-6.66 (br m, 1H); 7.25-7.63 (m, 6H); 7.86-7.92 (br m, 1H); 8.57-8.65 (br m, 1H). MS (APCI) (M+H)⁺ at m/z 456.

Example 179

5 (2-Isopropylphenyl)[2-trifluoromethyl-4-(E-(((2-carboxypyrrol-3-in-1-yl)carbonyl)ethenyl) phenyl] sulfide

The title compound was prepared according to the procedures of Example 1.

¹H NMR (CDCl₃, 300 MHz) δ 7.79 (s, 1H), 7.72 (d, 1H, J = 15.5 Hz), 7.49 (d, 1H, J = 7.4 Hz), 7.36-7.46 (m, 3H), 7.23 (m, 1H), 6.82 (d, 1H, J = 8.5 Hz), 6.74 (d, 1H, J = 10 15.4 Hz), 6.00 (br, 2H), 4.48 (br, 1H), 4.51 (br, 2H), 3.48 (m, 1H), 1.18 (d, 6H, J = 7.0 Hz). MS (ESI) m/z -460, -492, -921.

Example 180

15 (2-Isopropylphenyl)[2-trifluoromethyl-4-(E-(((2-hydroxymethylpyrrolidin-1-yl)carbonyl)ethenyl) phenyl] sulfide

The title compound was prepared according to the procedures of Example 1.

¹H NMR (CDCl₃, 300 MHz) δ 7.79 (s, 1H), 7.68 (d, 1H, J = 15.4 Hz), 7.48 (d, 1H, J = 7.4 Hz), 7.45 (m, 2H), 7.38 (d, 1H, J = 8.3 Hz), 7.23 (m, 1H), 6.80 (d, 1H, J = 8.5 Hz), 6.70 (d, 1H, J = 15.4 Hz), 5.82 (m, 1H), 5.70 (m, 1H), 4.92 (m, 1H), 4.18 (br s, 2H), 20 3.76 (s, 3H), 3.78 (d, 1H, J = 11.5 Hz), 3.50 (m, 2H), 3.01 (t, 2H, J = 7.5 Hz), 2.58 (t, 2H, J = 7.6 Hz), 1.19 (d, 6H, J = 7.1 Hz). MS (ESI) m/z 450, 472, 921.

Example 181

(2-Isopropylphenyl)[2-nitro-4-(E-((3-methylaminocarbonyl)piperazin-1-yl)carbonyl)ethenyl] phenyl] sulfide

Prepared according to the procedures of Example 71, giving a yellow solid.

- 5 ¹H NMR (DMSO-d₆, 300MHz) δ 1.14 (d, J = 7.0 Hz, 6H); 2.60 (d, J = 4.4 Hz, 3H); 2.50-4.45 (br m, 7H); 3.30-3.40 (m, 1H); 6.62-6.66 (br m, 1H); 7.32-7.62 (m, 6H); 7.81-7.92 (m, 2H); 8.59-8.65 (br m, 1H). MS (APCI) (M+H)⁺ at m/z 469.

Example 182

- 10 (2-Isopropylphenyl)[2-nitro-4-(E-(((3-cyclopropylaminocarbonyl)piperazin-1-yl)carbonyl)ethenyl] phenyl] sulfide

Prepared according to the procedures of Example 71, giving a yellow solid.

- ¹H NMR (DMSO-d₆, 300MHz) δ 0.40-0.62 (br m, 4H); 1.14 (d, J = 6.8 Hz, 6H); 2.50-4.41 (br m, 8H); 3.30-3.40 (m, 1H); 6.62-6.67 (br m, 1H); 7.32-7.62 (m, 6H); 7.87-7.92 (m, 2H); 8.59-8.64 (br m, 1H). MS (APCI) (M+H)⁺ at m/z 495.
- 15

Example 183

(2-Isopropylphenyl)[2-nitro-4-(E-((3-carboxamidopiperazin-1-yl)carbonyl)ethenyl] phenyl] sulfide

- 20 Prepared according to the procedures of Example 71, giving a yellow solid.

¹H NMR (DMSO-d₆, 300MHz) δ 1.14 (d, J = 7.0 Hz, 6H); 2.50-4.42 (br m, 7H); 3.30-

3.40 (m, 1H); 6.62-6.67 (br m, 1H); 7.12-7.62 (m, 8H); 7.87-7.92 (m, 1H); 8.60-8.65 (br m, 1H). MS (APCI) (M+H)⁺ at m/z 455.

Example 184

5 (2-Isopropylphenyl)[2-nitro-4-(*E*-((3-carbomethoxy-4-oxopiperidin-1-yl)carbonyl)ethenyl) phenyl] sulfide

Prepared according to the procedures of Example 71, giving a yellow solid.

¹H NMR (DMSO-d₆, 300MHz) δ 1.14 (d, J = 6.8 Hz, 6H); 2.32-2.55 (br m, 2H); 3.30-3.40 (m, 1H); 3.64, 3.76 (s, s, 3H); 3.68-4.58 (br m, 5H); 6.64 (d, J = 8.5 Hz, 1H); 7.32-7.63 (m, 6H); 7.88-7.96 (m, 1H); 8.60-8.68 (m, 1H). MS (APCI) (M+H)⁺ at m/z 483. Anal calcd for C₂₃H₂₆N₂S₂O₆·0.17C₆H₁₄: C, 62.86; H, 5.75; N, 5.63. Found: C, 62.81; H, 5.83; N, 5.60.

Example 185

15 (2-Isopropylphenyl)[2-nitro-4-(*E*-((3,5-dimethylpiperazin-1-yl)carbonyl)ethenyl) phenyl] sulfide

Prepared according to the procedures of Example 71, giving a yellow solid. ¹H

NMR (DMSO-d₆, 300MHz) δ 0.96-1.06 (m, 6H); 1.14 (d, J = 6.8 Hz, 6H); 2.07-4.39 (br m, 7H); 6.63 (d, J = 8.5 Hz, 1H); 7.30-7.63 (m, 6H); 7.92 (dd, J = 8.5, 1.7 Hz, 1H); 8.60 (d, J = 1.7 Hz, 1H). MS (APCI) (M+H)⁺ at m/z 440. Anal calcd for C₂₆H₂₉N₃S₂O₃: C, 65.58; H, 6.65; N, 9.56. Found: C, 65.36; H, 6.87; N, 9.27.

Example 186

(1-Ethylindol-7-yl)[2-chloro-4-(*E*-((4-acetyl)piperazin-1-yl)carbonyl)ethenyl] phenyl]
sulfide

The title compound was prepared by the procedures described in Example 85,
 5 substituting 5-iodoindole with *N*-ethyl-7-bromoindole. white solid; ¹H NMR (CDCl₃,
 300 MHz) δ 1.30 (t, *J* = 7.05 Hz, 3H), 2.14 (s, 3H), 3.52 (br s, 2H), 3.58-3.84 (m, 6H),
 4.42 (q, *J* = 7.05 Hz, 2H), 6.42 (d, *J* = 8.4 Hz, 1H), 6.59 (d, *J* = 3.0 Hz, 1H), 6.76 (d, *J*
 = 15.6 Hz, 1H), 7.08 (d, *J* = 8.4 Hz, 1H), 7.10 (d, *J* = 3.0 Hz, 1H), 7.16 (t, *J* = 7.65 Hz,
 1H), 7.42 (dd, *J* = 0.9, 7.5 Hz, 1H), 7.53 (d, *J* = 1.8 Hz, 1H), 7.54 (d, *J* = 15.6 Hz, 1H),
 10 7.78 (dd, *J* = 0.9, 7.5 Hz, 1H). MS (APCI⁺) (M+H)⁺ at *m/z* 468, 470.

Example 187

(3-[2-Methoxy]ethoxyphenyl)-[2-chloro-4(*E*-[(morpholin-1-
yl)carbonyl]ethenyl)phenyl] sulfide

15 The title compound was prepared according to the procedures of Example 85.
¹H-NMR (CDCl₃, 300 MHz) δ 3.45 (s, 3H), 3.65-3.80 (m, 10H), 4.09-4.13 (m, 2H),
 6.82 (broad d, *J*=15, 1H), 6.88 (d, *J*=9Hz, 1H), 6.87 (dd, *J*=9Hz, 2Hz, 1H), 7.03-7.10
 (m, 2H), 7.20 (d, *J*=9Hz, 1H), 7.31 (t, *J*=8 Hz, 1H), 7.52 (s, 1H), 7.56 (broad d, *J*=15,
 1H).

Example 188

(2-Bromophenyl)[2-chloro-4-(E-((4,4'-S-dioxythiomorpholin-1-yl)carbonyl)ethenyl)phenyl]sulfide

- 4-Methylmorpholine *N*-oxide (0.0935 g, 0.798 mmol) and 4Å molecular
 5 sieves (0.0333g) were added to a solution of (2-Bromophenyl)[2-chloro-4-(E-
 ((thiomorpholin-1-yl)carbonyl) ethenyl)phenyl]sulfide (0.1230g, 0.27 mmol; prepared
 according to the procedures described in Example 1). After 15 min,
 tetrapropylammonium perruthenate (0.0058g, 0.0166 mmol) was added and after 4h
 had elapsed the starting material was consumed by TLC and the crude products were
 10 passed through a plug of silica with 5:2 hexane:ethyl acetate→ 9:1 CH₂Cl₂: MeOH.
 The mixture was then purified by preparative HPLC to provide the title compound
 (0.0138 g, 10%). ¹H-NMR (DMSO-d₆, 300MHz) δ 8.12 (d, *J* = 1.47 Hz, 1H), 7.81
 (dd, *J* = 7.9, 1.3, 2H), 7.65 (dd, *J* = 8.0, 1.5 Hz, 1H), 7.47 (d, *J* = 9.0 Hz, 1H), 7.27-
 7.53 (m, 4H), 7.03 (d, *J* = 9.0 Hz, 1H), 4.12 (br s, 2H), 3.98 (br s, 2H), 3.26 (br s, 2H),
 15 3.19 (br s, 2H), 1.54-2.29 (m, 6H), MS(APCI) (M+H)⁺ at *m/z* 486, 488, 490.

Example 189

(2-Bromophenyl)[2-chloro-4-(E-(N-carbomethoxymethyl-N-(3-(pyrrolidin-2-on-1-yl)prop-1-yl)amino)carbonyl)ethenyl)phenyl]sulfide

Example 189A

N-Carbomethoxymethyl-N-(3-(pyrrolidin-2-on-1-yl)prop-1-yl)amine

Methyl bromoacetate (1.35 mL, 14.3 mmol) was added dropwise to a solution of 3-aminopropyl-2-pyrrolidinone (2.0 mL, 14.3 mmol) and diisopropylethylamine (2.7 mL) in CH₂Cl₂. The reaction was stirred for 12h and was then concentrated *in vacuo*, and carried forward without further purification.

5

Example 189B

(2-Bromophenyl)[2-chloro-4-(E-(N-carbomethoxymethyl-N-(3-(pyrrolidin-2-on-1-yl)prop-1-yl)amino)carbonyl) ethenyl]phenyl]sulfide

The title compound was prepared by the procedures described for Example 10 113, substituting 2,4 dichlorothiophenol with 2-bromothiophenol, 2-chlorobenzaldehyde with 3,4 dichlorobenzaldehyde, and 1-(3-aminopropyl)-5-((S)-hydroxymethyl)-2-pyrrolidinone with the compound from Example 189A. ¹H-NMR (DMSO-d₆, 300MHz) δ 8.07 (dd, *J* = 9.4, 1.7 Hz, 1H), 7.81 (m, 1H), 7.64 (m, 1H), 7.24-7.49 (m, 5H), 7.05 (m, 1H), 4.53 (s, 1H), 4.14 (s, 1H), 3.68 (s, 1H), 3.64 (s, 2H), 15 3.54 (m, 2H), 3.13-3.43 (m, 4H), 2.39 (m, 2H), 1.91 (m, 2H), 1.72 (m, 2H), MS(APCI) (M+H)⁺ at *m/z* 565, 567, 569.

Example 190

(2-Bromophenyl)[2-chloro-4-(E-((4-S-oxythiomorpholin-1-yl)-2-pyrrolidinone)carbonyl) ethenyl]phenyl]sulfide

20

The title compound (0.0178g, 14%) was isolated from the same reaction mixture as described in Example 188. ¹H-NMR (DMSO-d₆, 300MHz) δ 8.12 (d, *J* =

1.8 Hz, 1H), 7.81 (dd, $J = 7.9, 1.3$ Hz, 1H), 7.65 (dd, $J = 8.3, 1.7$ Hz, 1H), 7.46 (d, $J = 7.4$ Hz, 1H), 7.26-7.48 (m, 4H), 7.04 (d, $J = 7.4$ Hz, 1H), 4.29 (br m, 2H), 3.97 (br m, 1H), 3.61 (br m, 1H), 2.80 (br m, 4H), MS(APCI) (M+H)⁺ at m/z 470, 472, 474.

5

Example 191

(2-Methoxy-5-chlorophenyl)[2-nitro-4-(*E*-((4-acetylpiperazin-1-yl)carbonyl)ethenyl)phenyl] sulfide

The title compound was prepared according to the procedures of Example 1.

¹H NMR (CDCl₃, 300 MHz) δ 8.44 (s, 1H), 7.66 (d, 1H, $J = 15.1$ Hz), 7.58 (d, 1H, $J = 2.6$ Hz), 7.48 (dd, 1H, $J = 2.6, 8.8$ Hz), 7.44 (m, 1H), 6.97 (d, 1H, $J = 8.8$ Hz), 6.92 (d, 1H, $J = 15.5$ Hz), 6.82 (d, 1H, $J = 8.5$ Hz), 3.78 (s, 3H), 3.70 (m, 6H), 3.54 (m, 2H), 2.15 (s, 3H). MS (ESI) m/z 476, 498, 951, 973. Anal. Calcd for C₂₂H₂₂ClN₃O₅S · 0.48 EtOAc: C, 55.44; H, 5.03; N, 8.11. Found: C, 54.36; H, 4.90; N, 8.50.

15

Example 192

(2-Isopropylphenyl)[2-nitro-4-(*E*-((3-acetoxymethyl)piperazin-1-yl)carbonyl)ethenyl)phenyl] sulfide

Prepared according to the procedures of Example 71, giving a yellow solid.

¹H NMR (DMSO-d₆, 300MHz) δ 1.14 (d, $J = 6.8$ Hz, 6H); 2.04 (s, 3H); 3.30-3.40 (m, 1H); 2.50-4.46 (br m, 9H); 6.64 (d, $J = 8.8$ Hz, 1H); 7.30-7.62 (m, 6H); 7.87-7.93 (m, 1H); 8.58-8.63 (br m, 1H). MS (APCI) (M+H)⁺ at m/z 484. Anal calcd for C₂₅H₂₈N₂S₂O₆·0.2H₂O: C, 61.60; H, 6.09; N, 8.62. Found: C, 61.63; H, 6.21; N, 8.41.

00111795.03100

Example 193

(2-Isopropylphenyl)[2-nitro-4-(*E*-((3,5-dimethyl-4-acetyl-piperazin-1-yl)carbonyl)ethenyl) phenyl] sulfide

5 Prepared according to the procedures of Example 71, giving a yellow solid.

¹H NMR (DMSO-d₆, 300MHz) δ 1.00-1.20 (br m, 6H); 1.15 (d, *J* = 6.8Hz, 6H); 2.04 (s, 3H); 2.76-4.58 (br m, 7H); 6.64 (d, *J* = 8.5 Hz, 1H); 7.32-7.63 (m, 6H); 7.94 (dd, *J* = 8.5, 1.8 Hz, 1H); 8.66 (d, *J* = 1.8 Hz, 1H). MS (APCI) (M+H)⁺ at *m/z* 482. Anal calcd for C₂₈H₃₁N₃O₄·0.3H₂O: C, 64.13; H, 6.54; N, 8.63. Found: C, 64.15; H, 6.61;

10 N, 8.50.

Example 194

(1-Methylindol-5-yl)[2-chloro-4-(*E*-((4-acetyl-piperazin-1-yl)carbonyl)ethenyl) phenyl] sulfide

15 The title compound was prepared by the procedures described in Example 85, substituting 5-iodoindole with *N*-methyl-5-bromoindole, giving a white solid. ¹H NMR (d⁶-DMSO, 300 MHz) δ 2.04 (s, 3H), 3.40-3.80 (m, 8H), 3.86 (s, 3H), 6.49 (d, *J* = 8.4 Hz, 1H), 6.52 (d, *J* = 3.0 Hz, 1H), 7.27 (d, *J* = 15.6 Hz, 1H), 7.31 (dd, *J* = 2.4, 8.4 Hz, 1H), 7.39 (d, *J* = 15.6 Hz, 1H), 7.41 (dd, *J* = 1.8, 8.4 Hz, 1H), 7.48 (d, *J* = 3.0 Hz, 1H), 7.63 (d, *J* = 8.4 Hz, 1H), 7.85 (d, *J* = 1.8 Hz, 1H), 7.99 (br s, 1H). MS (APCI)⁺ (M+H)⁺ at *m/z* 454, 456.

20

Example 195

(Benzodioxan-6-yl)[2-nitro-4-(*E*-((4-acetyl)piperazin-1-yl)carbonyl)ethenyl] phenyl]
sulfide

5

Example 195A6-Mercaptobenzodioxane

The title compound was prepared by the procedures described in Example 97A, substituting 2-ethoxybenzene with 6-iodobenzenedioxane.

10

Example 195B

(Benzodioxan-6-yl)[2-nitro-4-(*E*-((4-acetyl)piperazin-1-yl)carbonyl)ethenyl] phenyl]
sulfide

The title compound was prepared by the procedures described in Example 32, substituting 2,4-dichlorobenzenethiol with 6-mercaptobenzenedioxane, to give a light-yellow solid; ¹H NMR (d⁶-DMSO, 300 MHz) δ 2.04 (s, 3H), 3.41-3.80 (m, 8H), 4.28-4.38 (m, 4H), 6.86 (d, *J* = 8.4 Hz, 1H), 7.05 (d, *J* = 8.4 Hz, 1H), 7.10 (dd, *J* = 2.1, 8.4 Hz, 1H), 7.15 (d, *J* = 2.1 Hz, 1H), 7.40 (d, *J* = 15.6 Hz, 1H), 7.53 (d, *J* = 15.6 Hz, 1H), 7.91 (dd, *J* = 1.8, 8.4 Hz, 1H), 8.62 (d, *J* = 1.8 Hz, 1H). MS (APCI⁺) (M+H)⁺ at *m/z* 470. Anal. Calcd for C₂₃H₂₃N₃O₆S · 0.17 H₂O: C, 58.46; H, 4.98; N, 8.89. Found:

C, 58.47; H, 4.88; N, 8.78.

Example 196035475-03100
01100-5674500

(Benzodioxan-6-yl)[2-nitro-4-(E-((3-(pyrrolidin-2-on-1-yl)prop-1-ylamino)carbonyl)ethenyl)phenyl]sulfide

The title compound was prepared by the procedures described in Example 32, substituting 2,4-dichlorobenzenethiol with 6-mercaptobenzenedioxane, and 1-

- 5 acetylpiiperazine with 3-aminopropyl-1-pyrrolidin-2-one, giving a light-yellow solid.
- ¹H NMR (d⁶-DMSO, 300 MHz) δ 1.64 (p, *J* = 7.2 Hz, 2H), 1.92 (p, *J* = 7.8 Hz, 2H), 2.21 (t, *J* = 7.8 Hz, 2H), 3.13 (t, *J* = 7.2 Hz, 2H), 3.19 (t, *J* = 7.2 Hz, 2H), 3.38-3.46 (overlapping t, *J* = 7.8 Hz, 2H), 4.27-4.37 (m, 4H), 6.70 (d, *J* = 15.6 Hz, 1H), 6.90 (d, *J* = 8.4 Hz, 1H), 7.05 (d, *J* = 8.4 Hz, 1H), 7.09 (dd, *J* = 2.1, 8.4 Hz, 1H), 7.16 (d, *J* = 2.1 Hz, 1H), 7.46 (d, *J* = 15.6 Hz, 1H), 7.77 (dd, *J* = 2.1, 8.4 Hz, 1H), 8.16 (t, *J* = 6.0 Hz, 1H), 8.41 (d, *J* = 2.1 Hz, 1H). MS (APCI⁺) (M+H)⁺ at *m/z* 484. Anal. Calcd for C₂₄H₂₅N₃O₆S · 0.51 CH₂Cl₂ · 0.24 MeOH: C, 55.61; H, 5.09; N, 7.86. Found: C, 55.39; H, 5.48; N, 8.26.
- 10

15

Example 197

(Benzodioxan-6-yl)[2-nitro-4-(E-((3-carboethoxypiperidin-1-yl) carbonyl)ethenyl)phenyl] sulfide

The title compound was prepared by the procedures described in Example 196 substituting N-(3'-aminopropyl)-2-pyrrolidinone with ethyl nipecotate, giving a

- 20 yellow solid, mp 73-75 °C. ¹H NMR (CDCl₃, 300 MHz) δ 1.26 (t, *J*=7.0 Hz, 3H), 1.74 (br, 1H), 1.78 (br, 1H), 2.10 (br, 1H), 2.54 (br, 1H), 2.95-3.70 (br, 2H), 3.90-4.10 (br, 2H), 4.15 (q, *J*=7.0 Hz, 2H), 4.30-4.40 (m, 4H), 4.65 (br, 1H), 6.90 (d, *J*=8.5 Hz,

1H), 6.98 (d, J=8.5 Hz, 1H), 7.06 (dd, J=2.0, 8.0 Hz, 1H), 7.10 (d, J=2.0 Hz, 1H), 7.40-7.50 (m, 1H), 7.58 (d, J=15.0 Hz, 1H), 8.40 (d, J=2.0 Hz, 1H). MS (APCI) m/z 499 (M+H)⁺. Anal. calcd. for C₂₅H₂₆N₂O₇S: C, 60.23; H, 5.26; N, 5.62. Found: C, 60.09; H, 5.43; N, 5.47.

5

Example 198

(Benzodioxan-6-yl)[2-nitro-4-(E-((4-carboethoxypiperidin-1-yl) carbonyl)ethenyl) phenyl] sulfide

The title compound was prepared by the procedure described as in example 10 196 substituting N-(3'-aminopropyl)-2-pyrrolidinone with ethyl isonipecotate, giving a yellow solid, mp 78-88 °C. ¹H NMR (CDCl₃, 300 MHz) δ 1.27 (t, J=7.0 Hz, 3H), 1.65 (m, 2H), 2.00 (m, 2H), 2.60 (m, 1H), 2.80-3.50 (br, 2H), 4.15 (br, 1H), 4.16 (q, J=7.0, 2H), 4.34 (m, 4H), 4.54 (br, 1H), 6.90 (d, J=8.0 Hz, 1H), 6.98 (d, J=8.0 Hz, 1H), 7.05 (dd, J=2.0, 8.0 Hz, 1H), 7.10 (d, J=2.0 Hz, 1H), 7.12 (br, 1H), 7.44 (d, J=8.0 15 Hz, 1H), 7.60 (br, 1H), 8.40 (s, 1H). MS (Cl/NH₃) m/z 499 (M+H)⁺. Anal. calcd. for C₂₅H₂₆N₂O₇S 0.03 H₂O: C, 60.16; H, 5.26; N, 5.61. Found: C, 60.15; H, 5.65; N, 5.40.

Example 199

(2-Ethoxyphenyl)[2-trifluoromethyl-4-(Z-((4-acetylpiperazin-1-yl)carbonyl)ethenyl) phenyl] sulfide

20

Example 199A

(2-Ethoxyphenyl)[2-trifluoromethyl-4-(Z-((4-carbomethoxyethenyl) phenyl] sulfide

Bis-(2,2,2-trifluoroethyl)(methoxycarbonylmethyl)phosphonate (1.20 g, 3.77 mmole), and 18-crown-6 (3.56 g, 13.48 mmol) were dissolved in 22 ml of dry THF. The mixture was cooled to -78 °C and $\text{KN}(\text{SiMe}_3)_2$ (0.5 M in THF, 4.04 mmol) was added and stirred for 30 min. (2-Ethoxyphenyl)[2-trifluoromethyl-4-formyl phenyl] sulfide (1.10 g, 3.77 mmol, prepared according to the procedure of example 1) in 13 ml of THF was added via cannulation. After 1 hr at that temperature, the cooling bath was removed and the mixture allowed to warm to ambient temperature. Saturated NH_4Cl soln. was added and the mixture was extracted with ethyl acetate three times. The combined organics were dried over sodium sulfate, concentrated *in vacuo* and purified by medium pressure chromatography on silica gel to give 772 mg (60% yield) of the *cis*- isomer ($J_{\text{olefinic}} = 12.5 \text{ Hz}$) along with 322 mg (25% yield) of the *trans*- isomer ($J_{\text{olefinic}} = 12.5 \text{ Hz}$).

Example 199B

(2-Ethoxyphenyl)[2-trifluoromethyl-4-(Z-((4-acetyl)piperazin-1-yl)carbonyl)ethenyl] phenyl] sulfide

The compound of Example 199A was converted to the corresponding amide according to the procedures of Example 1. ^1H NMR (CDCl_3 , 300 MHz) δ 7.64 (d, 1H, $J = 16.9 \text{ Hz}$), 7.32-7.4 (m, 2H), 6.98 (m, 2H), 6.93 (m, 2H), 6.65 (d, 1H, $J = 12.1 \text{ Hz}$), 6.08 (d, 1H, $J = 12.2 \text{ Hz}$), 3.98 (q, 2H, $J = 7.0 \text{ Hz}$), 3.68 (m, 2H), 3.62 (m, 2H), 3.44-3.54 (m, 4H), 2.11 and 2.05 (s, 3H), 1.20 (t, 3H, $J = 7.0 \text{ Hz}$). MS (ESI)s m/z 479, 501.

Example 200

(2-Ethoxyphenyl)[2-trifluoromethyl-4-(*E*-((6-methylpyrid-2-ylamino)carbonyl)ethenyl) phenyl] sulfide

The title compound was prepared according to the procedures of Example 1.

¹H NMR (CDCl₃, 300 MHz) δ 8.12 (d, 1H, J = 8.1 Hz), 7.78 (s, 1H, J = 1.7 Hz), 7.70 (d, 1H, J = 15.6 Hz), 7.63 (t, 1H, J = 7.8 Hz), 7.46 (dd, 1H, J = 1.6, 7.8 Hz), 7.36-7.42 (m, 2H), 7.04 (d, 1H, J = 8.1), 6.99 (dd, 1H, J = 1.2, 7.6 Hz), 6.92 (m, 2H), 6.50 (d, 1H, J = 15.6 Hz), 3.99 (q, 2H, J = 6.9 Hz), 2.47 (s, 3H), 1.19 (t, 3H, J = 7.0 Hz). MS (ESI)s *m/z* 459, 481. Anal. Calcd for C₂₄H₂₁F₃N₂O₂S · 1.1 H₂O: C, 60.27; H, 4.89; N, 5.86. Found: C, 60.28; H, 5.05; N, 5.94.

Example 201

(2-Methyl-3-chlorophenyl)[2-nitro-4-(*E*-((4-acetyl)piperazin-1-yl)carbonyl)ethenyl] phenyl] sulfide

The title compound was prepared according to the procedures of Example 1.

¹H NMR (CDCl₃, 300 MHz) δ 8.46 (d, 1H, J = 1.5 Hz), 7.64 (d, 1H, J = 15.4 Hz), 7.56 (d, 1H, J = 2.6 Hz), 7.54 (d, 1H, J = 2.2 Hz), 7.47 (d, 1H, J = 8.5 Hz), 7.27 (m, 1H), 6.92 (d, 1H, J = 15.4 Hz), 6.68 (d, 1H, J = 8.5 Hz), 3.63-3.78 (m, 6H), 3.53 (m, 2H), 2.45 (s, 3H), 2.15 (s, 3H). MS (ESI) *m/z* 460, 482, 919. Anal. Calcd for C₂₂H₂₂Cl₁N₃O₄S: C, 57.45, H, 4.82, N, 9.14. Found: C, 75.54, 5.08, N, 8.82.

Example 202

(Benzodioxan-6-yl)[2-nitro-4-(*E*-(3-carboxamidopiperidin-1-yl) carbonyl)ethenyl]
phenyl sulfide

The title compound was prepared by the procedures described in Example 196, substituting N-(3'-aminopropyl)-2-pyrrolidinone with nipecotamide, giving a light
5 yellow solid, mp 243-245 °C. ¹H NMR (CDCl₃, 500 MHz) δ 1.38-1.50 (m, 2H), 1.77-2.00 (m, 2H), 2.38 (m, 1H), 2.70 (m, 1H), 3.11 (m, 1H), 4.22 (m, 1H), 4.28-4.30 (m, 2H), 4.32-4.36 (m, 2H), 4.42 (m, 1H), 6.85 (d, J=8.5 Hz, 1H), 7.04-7.16 (m, 2H), 7.35 (s, 1H), 7.40 (d, J=13.0 Hz, 1H), 7.48 (d, J=15.5 Hz, 1H), 7.91 (d, J=8.5 Hz, 1H), 8.58 (s, 1H). MS (APCI) m/z 470 (M+H)⁺. Anal. calcd. for C₂₃H₂₃N₃O₆S•0.37 H₂O: C, 58.01; H, 5.03; N, 8.82. Found: C, 58.02; H, 5.13; N, 8.61.

Example 203

(Benzodioxan-6-yl)[2-nitro-4-(*E*-((2-carboethoxypiperidin-1-yl) carbonyl)ethenyl]
phenyl sulfide

The title compound was prepared by the procedures described in Example 196, substituting N-(3'-aminopropyl)-2-pyrrolidinone with ethyl pipecolate, producing a
15 light yellow solid, mp 74-75 °C. ¹H NMR (CDCl₃, 300 MHz) δ 1.28 (t, J=7.0 Hz, 3H), 1.32-1.55 (m, 2H), 1.60-1.82 (m, 3H), 2.33 (m, 1H), 3.40 (m, 1H), 3.98 (m, 1H), 4.23 (q, J=6.5 Hz, 2H), 4.32 (q, J=5.0 Hz, 4H), 5.45 (m, 1H), 6.90 (d, J=8.0 Hz, 1H),
20 6.97 (d, J=8.0 Hz, 1H), 7.0-7.10 (m, 3H), 7.44, (d, J=7.5 Hz, 1H), 7.60 (d, J=15.0 Hz, 1H), 8.38 (m, 1H). MS (APCI) m/z 499 (M+H)⁺. Anal. calcd. for C₂₅H₂₆N₂O₇S•0.11 H₂O: C, 59.99; H, 5.28; N, 5.60. Found: C, 59.98; H, 5.42; N, 5.91.

Example 204

(Benzodioxan-6-yl)[2-nitro-4-(E-((4-carboxamidopiperidin-1-yl) carbonyl)ethenyl)
phenyl] sulfide

5 The title compound was prepared by the procedures described in Example196, substituting N-(3'-aminopropyl)-2-pyrrolidinone with isonipecotamide, giving a light yellow solid, mp >230 °C. ¹H NMR (CDCl₃, 500 MHz) δ 1.35 (m, 1H), 1.60 (m, 1H), 1.72 (m, 1H), 1.68 (m, 1H), 2.20 (m, 1H), 2.75 (m, 1H), 3.04 (m, 1H), 3.20 (m, 1H), 4.20 (m, 1H), 4.32 (m, 4H), 6.85 (d, J=8.5 Hz, 1H), 7.04 (d, J=8.5 Hz, 1H), 7.09 (dd, J=2.0, 8.5 Hz, 1H), 7.26 (s, 1H), 7.37 (d, J=16.0 Hz, 1H), 7.47 (d, J=16.0 Hz, 1H), 8.58 (d, J=2.0 Hz, 1H). MS (APCI) m/z 470 (M+H)⁺. Anal. calcd. for C₂₃H₂₃N₅O₆S•0.13 H₂O: C, 58.55; H, 4.97; N, 8.91. Found: C, 58.41; H, 5.14; N, 9.30.

15

Example 205

(Benzodioxan-6-yl)[2-nitro-4-(E-((4-tert-butoxycarbonylpiperazin-1-yl)
carbonyl)ethenyl) phenyl] sulfide

The title compound was prepared by the procedures described in Example196, substituting N-(3'-aminopropyl)-2-pyrrolidinone with Boc-piperazine, giving a light yellow solid, mp 165-167 °C. ¹H NMR (CDCl₃, 300 MHz) δ 1.48 (s, 9H), 3.50 (m, 4H), 3.65 (br, m, 4H), 4.32 (m, 4H), 6.89 (d, J=5.0 Hz, 1H), 6.92 (m, 1H), 6.97 (d, J=8.0 Hz, 1H), 7.05 (dd, J=2.0, 8.5 Hz, 1H), 7.10 (d, J=2.0 Hz, 1H), 7.45

(m, 1H), 7.63 (d, J=15.5 Hz, 1H), 8.40 (m, 1H). MS (APCI) M/z 528 (M+H)⁺. Anal. calcd. for C₂₆H₂₉N₃O₇S: C, 59.19; H, 5.54; N, 7.96. Found: C, 58.85; H, 5.69; N, 8.20.

Example 206

5 (2-Isopropylphenyl)[2-nitro-4-(E-((syn-3,5-dimethylmorpholin-1-yl)carbonyl)ethenyl) phenyl] sulfide

Prepared according to the procedures of Example 71, giving a yellow solid.

¹H NMR (DMSO-d₆, 300MHz) δ 1.10-1.18 (m, 12H); 2.29-2.39 (m, 1H); 2.67-2.78 (m, 1H); 3.30-3.53 (m, 3H); 4.17-4.38 (m, 2H); 6.63 (d, J = 8.8 Hz, 1H); 7.32-7.63 (m, 6H); 7.92 (dd, J = 8.8, 1.5 Hz, 1H); 8.66 (d, J = 1.8 Hz, 1H). MS (APCI) (M+H)⁺ at m/z 441. Anal calcd for C₂₄H₂₈N₂S₁O₄: C, 65.43; H, 6.41; N, 6.36. Found: C, 65.69; H, 6.70; N, 6.17.

Example 207

15 (2-Isopropylphenyl)[2-nitro-4-(E-((anti-3,5-dimethylmorpholin-1-yl)carbonyl)ethenyl) phenyl] sulfide

Prepared according to the procedures of Example 71, giving a yellow solid.

¹H NMR (DMSO-d₆, 300MHz) δ 1.07-1.12 (m, 6H); 1.15 (d, J = 6.6 Hz, 6H); 3.32-3.48 (m, 3H); 3.60-3.83 (br m, 2H); 3.87-3.98 (m, 2H); 6.63 (d, J = 8.5 Hz, 1H); 7.32-7.63 (m, 6H); 7.93 (dd, J = 8.8, 1.8 Hz, 1H); 8.64 (d, J = 1.8 Hz, 1H). MS (APCI) (M+H)⁺ at m/z 441.

Example 208

(2-Isopropylphenyl)[2-nitro-4-(E-((3-carboethoxypiperazin-1-yl)carbonyl)ethenyl)
phenyl] sulfide

Prepared according to the procedures of Example 71, giving a yellow solid.

- 5 ¹H NMR (DMSO-d₆, 300MHz) δ 1.14 (d, J = 6.8 Hz, 6H); 1.08-1.26 (m, 3H); 2.52-3.16 (br m, 4H); 3.25-3.40 (m, 1H); 3.41-4.26 (br m, 5H); 6.61-6.67 (br m, 1H); 7.30-7.62 (m, 6H); 7.87-7.93 (br m, 1H); 8.58-8.64 (br m, 1H). MS (APCI) (M+H)⁺ at m/z 484. Anal calcd for C₂₅H₂₉N₃O₅: C, 62.09; H, 6.04; N, 8.69. Found: C, 61.96; H, 6.28; N, 8.49.

10

Example 209

(2-Isopropylphenyl)[2-nitro-4-(E-((3-isopropoxycarbonyl)piperazin-1-
yl)carbonyl)ethenyl) phenyl] sulfide

Prepared according to the procedures of Example 71, giving a yellow solid.

- 15 ¹H NMR (DMSO-d₆, 300MHz) δ 1.07-1.21 (br m, 6H); 1.14 (d, J = 7.0 Hz, 6H); 2.52-3.16 (br m, 4H); 3.30-3.40 (m, 1H); 3.41-4.24 (br m, 3H); 4.81-4.97 (m, 1H); 6.61-6.68 (br m, 1H); 7.32-7.63 (m, 6H); 7.87-7.94 (br m, 1H); 8.60-8.66 (br m, 1H). MS (APCI) (M+H)⁺ at m/z 498. Anal calcd for C₂₆H₃₁N₃O₅: C, 62.76; H, 6.28; N, 8.44. Found: C, 62.51; H, 6.52; N, 8.14.

20

Example 210

(2-Isopropylphenyl)[2-nitro-4-(E-((3-(dimethylaminocarbonyl)-4-methylpiperazin-1-yl)carbonyl)ethenyl) phenyl] sulfide

Prepared according to the procedures of Example 71, giving a yellow solid.

¹H NMR (DMSO-d₆, 300MHz) δ 1.14 (d, J = 6.8 Hz, 6H); 2.14 (s, 3H); 2.82, 2.84 (s, 5 s, 3H); 3.12 (s, 3H); 2.12-4.24 (br m, 8H); 6.64 (d, J = 8.5 Hz, 1H); 7.32-7.62 (m, 6H); 7.87-7.94 (br m, 1H); 8.60-8.66 (br m, 1H). MS (APCI) (M+H)⁺ at m/z 497. Anal calcd for C₂₈H₃₂N₄S₁O₄·0.42H₂O: C, 61.94; H, 6.56; N, 11.11. Found: C, 62.00; H, 6.78; N, 10.89.

Example 211

(2-Isopropylphenyl)[2-nitro-4-(E-((3-carbomethoxy-4-hydroxypiperidin-1-yl)carbonyl)ethenyl) phenyl] sulfide

Prepared according to the procedures of Example 71, giving a yellow solid.

¹H NMR (DMSO-d₆, 300MHz) δ 1.14 (d, J = 7.0 Hz, 6H); 1.59-1.75 (br m, 2H); 2.50-15 3.14 (br m, 1H); 3.30-3.40 (m, 1H); 3.60, 3.61 (s, s, 3H); 4.01-4.44 (br m, 4H); 5.05-5.10 (br m, 1H); 6.63 (d, J = 8.5 Hz, 1H); 7.34-7.62 (m, 6H); 7.87-7.94 (br m, 1H); 8.60-8.66 (br m, 1H). MS (APCI) (M+H)⁺ at m/z 485.

Example 212

(2-Isopropylphenyl)[2-nitro-4-(E-((3-hydroxymethyl-4-hydroxypiperidin-1-yl)carbonyl)ethenyl) phenyl] sulfide

Prepared according to the procedures of Example 71, giving a yellow solid. ¹H NMR (DMSO-d₆, 300MHz) δ 1.14 (d, J = 6.8 Hz, 6H); 1.49-1.90 (br m, 2H); 2.75-3.14 (br m, 1H); 3.30-3.40 (m, 1H); 3.40-4.23 (br m, 5H); 4.38-4.52 (m, 1H); 4.60-4.73 (m, 1H); 6.61-6.66 (m, 1H); 7.27-7.61 (m, 6H); 7.84-7.93 (br m, 1H); 8.54-8.63 (br m, 1H). MS (APCI) (M+H)⁺ at m/z 457. Anal calcd for C₂₄H₂₈N₂S₁O₄·0.47H₂O: C, 61.97; H, 6.27; N, 6.02. Found: C, 62.02; H, 6.49; N, 5.90.

Example 213

(2-Ethoxyphenyl)[2-trifluoromethyl-4-(E-((2-carbomethoxy-4-(methoxycarbonyl)piperazin-1-yl)carbonyl)ethenyl) phenyl] sulfide

The title compound was prepared according to the procedures of Example 71. ¹H NMR (CDCl₃, 300 MHz) δ 7.80 (s, 1H), 7.66 (d, 1H, J = 15.4 Hz), 7.45 (dd, 1H, J = 1.6, 7.5 Hz), 7.48 (m, 2H), 7.01 (d, 1H, J = 6.6 Hz), 6.95 (d, 1H, J = 6.8 Hz), 6.90 (m, 2H), 5.34 (br s, 1H), 4.66 (m, 2H), 3.76 (s, 3H), 3.73 (s, 3H), 3.18 (m, 1H), 3.00 (m, 3H). MS (ESI) m/z 553, 575.

Example 214

(2-Ethoxyphenyl)[2-trifluoromethyl-4-(E-((2-carbomethoxy-4-methyl piperazin-1-yl)carbonyl)ethenyl) phenyl] sulfide

The title compound was prepared according to the procedures of Example 71. ¹H NMR (CDCl₃, 300 MHz) δ 7.79 (s, 1H), 7.64 (d, 1H, J = 15.3 Hz), 7.45 (dd, 1H, J = 1.7, 7.8 Hz), 7.4-7.35 (m, 2H), 7.01 (d, 1H, J = 8.1 Hz), 6.97 (dd, 1H, J = 1.2, 7.6

Hz), 6.87-7.91 (m, 2H), 5.36 (br s, 1H), 3.98 (q, 2H, $J = 6.9$ Hz), 3.90 (m, 1H), 3.78 (s, 3H), 3.65 (m, 1H), 3.42 (m, 1H), 2.85 (m, 1H), 2.32 (s, 3H), 2.24 (m, 1H), 2.19 (m, 1H), 1.18 (t, 3H, $J = 6.9$ Hz). MS (ESI) m/z 509, 531.

5

Example 215

(2-Ethoxyphenyl)[2-trifluoromethyl-4-(*E*-((2-carboxy-4-(methoxycarbonyl)piperazin-1-yl)carbonyl)ethenyl) phenyl] sulfide

The title compound was prepared according to the procedures of Example 71.

¹H NMR (DMSO-*d*₆, 300 MHz) δ 8.10 (m, 1H), 7.68 (m, 1H), 7.42 (m, 2H), 7.30 (m, 1H), 7.20 (d, 1H, $J = 15.6$ Hz), 7.10 (d, 1H, $J = 8.1$ Hz), 7.04 (d, 1H, $J = 8.5$ Hz), 6.98 (d, 1H, $J = 7.5$ Hz), 4.65 (br s, 1H), 4.53 (m, 2H), 4.05 (m, 2H), 4.00 (q, 2H, $J = 6.9$ Hz), 3.57 (s, 3H), 1.09 (t, 3H, $J = 6.9$ Hz). MS (ESI) m/z -537, -569.

15

Example 216

(Indol-6-yl)[2-chloro-4-(*E*-((4-acetyl)piperazin-1-yl)carbonyl)ethenyl) phenyl] sulfide

The title compound was prepared by the procedures described in Example 85, substituting 5-iodoindole with 6-bromoindole, isolated as a white solid. ¹H NMR (*d*₆-DMSO, 300 MHz) δ 2.03 (s, 3H), 3.40-3.77 (m, 8H), 6.52-6.55 (m, 1H), 6.60 (d, $J = 8.4$ Hz, 1H), 7.13 (dd, $J = 1.8, 8.4$ Hz, 1H), 7.27 (d, $J = 15.6$ Hz, 1H), 7.40 (d, $J = 15.6$ Hz, 1H), 7.43 (dd, $J = 1.8, 8.4$ Hz, 1H), 7.51 (t, $J = 3.0$ Hz, 1H), 7.64 (m, 1H), 7.70 (d, $J = 8.4$ Hz, 1H), 7.99 (d, $J = 1.8$ Hz, 1H). MS (APCI⁺) (M+H)⁺ at m/z 440, 442.

00544795.033100

Example 217

(1-Ethyl-3-(dimethylaminomethyl)indol-7-yl)[2-chloro-4-(*E*-((4-acetyl)piperazin-1-yl)(carbonyl)ethenyl) phenyl] sulfide

5 The title compound was prepared by the procedures described in Example 85, substituting 5-iodoindole with 7-bromo-3-*N,N*-dimethylmethyl-*N*-ethyl indole, and isolated as a light-brown solid. ¹H NMR (CDCl₃, 300 MHz) δ 1.30 (t, *J* = 7.05 Hz, 3H), 2.14 (s, 3H), 2.41 (s, 6H), 2.93-3.05 (m, 2H), 3.47-3.55 (m, 2H), 3.55-3.87 (m, 6H), 6.42 (d, *J* = 8.4 Hz, 1H), 6.85 (d, *J* = 15.6 Hz, 1H), 7.09 (dd, *J* = 2.1, 8.4 Hz, 1H), 7.17 (d, *J* = 8.4 Hz, 1H), 7.23 (d, *J* = 8.4 Hz, 1H), 7.43 (dd, *J* = 0.9, 7.8 Hz, 1H), 7.52 (d, *J* = 2.1 Hz, 1H), 7.54 (d, *J* = 15.6 Hz, 1H), 7.81 (dd, *J* = 0.9, 7.8 Hz, 1H). MS (ESI⁺) (M+H)⁺ at *m/z* 525, 527.

Example 218

15 (5-Ethoxybenzodioxan-6-yl)[2-chloro-4-(*E*-((4-acetyl)piperazin-1-yl)(carbonyl)ethenyl) phenyl] sulfide

 The title compound was prepared by the procedures described in Example 85, substituting 5-iodoindole with 6-bromo-5-ethoxybenzodioxane, as a white solid. ¹H NMR (CDCl₃, 300 MHz) δ 1.28 (t, *J* = 7.2 Hz, 3H), 2.14 (s, 3H), 3.54 (br s, 2H), 3.60-3.88 (m, 6H), 4.06 (q, *J* = 7.2 Hz, 2H), 4.33 (s, 4H), 6.70 (d, *J* = 8.4 Hz, 1H), 6.73 (d, *J* = 8.4 Hz, 1H), 6.78 (d, *J* = 15.6 Hz, 1H), 6.98 (d, *J* = 8.4 Hz, 1H), 7.17 (dd,

$J = 1.8, 8.4 \text{ Hz}$, 1H), 7.50 (d, $J = 1.8 \text{ Hz}$, 1H), 7.57 (d, $J = 15.6 \text{ Hz}$, 1H). MS (APCI⁺) ($M+H$)⁺ at m/z 503, 505.

Example 219

5 (2-Ethyl-4-bromophenyl)[2-nitro-4-(*E*-((4-acetylpiperazin-1-yl)carbonyl)ethenyl) phenyl] sulfide

The title compound was prepared according to the procedures of Example 32.

¹H NMR (CDCl₃, 300 MHz) δ 8.43 (d, 1H, $J = 2.0 \text{ Hz}$), 7.64 (d, 1H, $J = 15.6 \text{ Hz}$), 7.58
(d, 1H, $J = 2.0 \text{ Hz}$), 7.40-7.48 (m, 3H), 6.90 (d, 1H, $J = 15.2 \text{ Hz}$), 6.90 (d, 1H, $J = 8.5$
10 Hz), 3.63-3.77 (m, 6H), 3.54 (m, 2H), 2.72 (q, 2H, $J = 7.5 \text{ Hz}$), 2.15 (s, 3H), 1.18 (t,
3H, $J = 7.5 \text{ Hz}$). MS (ESI) m/z 518, 520, 542, 627. Anal. Calcd for C₂₃H₂₄BrN₃O₄S
: C, 53.08; H, 4.60; N, 7.93. Found: C, 53.29, H, 4.67, N, 8.11.

Example 220

15 (Benzodioxan-6-yl)[2-nitro-4-(*E*-((2-carboxypiperidin-1-yl) carbonyl)ethenyl) phenyl] sulfide

The title compound was prepared by the hydrolysis of the compound of

Example 203 under basic conditions (aq. NaOH/EtOH), producing a light yellow
solid.; mp 165 °C(dec.). ¹H NMR (DMSO-*d*₆, 300 MHz) δ 1.15-1.52 (m, 3H), 1.46-
20 1.62 (m, 2H), 2.32 (m, 1H), 2.80 (m, 1H), 3.45(br, 1/2H), 4.00 (br, 1/2H), 4.44 (br,
1/2H), 4.800 (br, 1/2H), 6.83 (d, $J=8.0 \text{ Hz}$, 1H), 7.03 (d, $J=8.0 \text{ Hz}$, 1H), 7.09 (dd,
 $J=2.0, 14.0 \text{ Hz}$, 1H), 7.15 (d, $J=2.0 \text{ Hz}$, 1H), 7.20 (d, $J=15.5 \text{ Hz}$, 1H), 7.35 (d, $J=15.5$

Hz, 1H), 7.73 (m, 1H), 8.52 (m, 1H). MS (ESI) m/z 469 (M-H)⁺, 471 (M+H)⁺. Anal. calcd. for C₂₃H₂₁N₂O₇SnNa •NaOH•2.7 H₂O: C, 47.54; H, 4.75; N, 4.82. Found: C, 47.18; H, 4.36; N, 4.89.

5

Example 221

(Benzodioxan-6-yl)[2-nitro-4-(E-((4-carboxymethyl)piperazin-1-yl) carbonyl)ethenyl
phenyl] sulfide

The title compound was prepared by deprotection of the compound 33 with

10 TFA in CH₂Cl₂. The resultant free amine was treated with *tert*-butyl bromoacetate and TEA in acetonitrile at room temperature, and followed by deprotection with TFA in CH₂Cl₂, giving a light solid, mp 120 °C (dec.). ¹H NMR (DMSO-d₆, 300 MHz) δ 3.20-3.45 (m, 4H), 4.20 (s, 2H), 3.50-3.80 (m, 4H), 4.28-4.46 (m, 4H), 6.86 (d, J=8.5 Hz, 1H), 7.04 (m, J=8.0 Hz, 1H), 7.09 (dd, J=2.0 8.0 Hz, 1H), 7.15 (d, J=2.0 Hz, 1H),

15 7.40 (d, J=15.5 Hz, 1H), 7.56 (d, J=15.0 Hz, 1H), 7.90 (dd, J=2.0, 8.5 Hz, 1H), 8.63 (m, 1H). MS (ESI) m/z 484 (M-H)⁺, 486 (M+H)⁺. Calcd. Anal for C₂₃H₂₁N₃O₇S•1.19CF₃COOH•1.34 H₂O: 47.63; H, 4.11; N, 6.89. Found: C, 47.93; H, 4.51; N, 6.49.

20

Example 222

(3-Morpholinophenyl)[2-nitro-4-(E-((4-acetyl)piperazin-1-yl)carbonyl)ethenyl
phenyl] sulfide

The title compound was prepared according to the procedures of Example 62, employing the compound of Example 103 as starting material. ¹H NMR (CDCl₃, 300 MHz) δ 7.80 (s, 1H), 7.64 (d, 1H, J = 15.4 Hz), 7.43 (m, 1H), 7.32 (t, 1H, J = 8.1 Hz), 7.08 (m, 2H), 6.99 (m, 2H), 6.84 (d, 1H, J = 15.4 Hz), 3.87 (t, 4H, J = 4.8 Hz), 3.63-3.79 (m, 6H), 3.50-3.55 (m, 2H), 3.18 (t, 4H, J = 4.8 Hz), 2.10 (s, 3H). MS (ESI) *m/z* 520, 542, 1061.

Example 223

(5-Ethoxybenzodioxan-8-yl)[2-chloro-4-(*E*-(4-acetyl)piperazin-1-yl)carbonyl]ethenyl phenyl sulfide

10

The title compound was prepared by the procedures described in Example 85, substituting 5-iodoindole with 8-bromo-5-ethoxybenzodioxane, giving a white solid. ¹H NMR (CDCl₃, 300 MHz) δ 1.52 (t, *J* = 7.2 Hz, 3H), 2.15 (s, 3H), 3.48-3.59 (m, 2H), 3.59-3.85 (m, 6H), 4.16 (q, *J* = 7.2 Hz, 2H), 4.22-4.30 (m, 2H), 4.30-4.40 (m, 2H), 6.59 (d, *J* = 8.7 Hz, 1H), 6.63 (d, *J* = 8.7 Hz, 1H), 6.78 (d, *J* = 15.6 Hz, 1H), 7.08 (d, *J* = 8.7 Hz, 1H), 7.17 (dd, *J* = 2.1, 8.7 Hz, 1H), 7.51 (d, *J* = 2.1 Hz, 1H), 7.58 (d, *J* = 15.6 Hz, 1H). MS (APCI⁺) (*M*+H)⁺ at *m/z* 503, 505.

15

Example 224

(5-Chloro-8-ethoxyquinolin-7-yl)[2-chloro-4-(*E*-(4-acetyl)piperazin-1-yl)carbonyl]ethenyl phenyl sulfide

20

The title compound was prepared by the procedures described in Example 85, substituting 5-iodoindole with 5-chloro-8-ethoxy-7-iodoquinoline, giving a white solid. ¹H NMR (d⁶-DMSO, 300 MHz) δ 1.37 (t, *J* = 7.2 Hz, 3H), 2.04 (s, 3H), 3.41-3.82 (m, 8H), 4.46 (q, *J* = 7.2 Hz, 2H), 7.29 (s, 1H), 7.37 (d, *J* = 8.4 Hz, 1H), 7.42 (d, *J* = 15.6 Hz, 1H), 7.51 (d, *J* = 15.6 Hz, 1H), 7.68 (dd, *J* = 1.8, 8.4 Hz, 1H), 7.74 (dd, *J* = 3.9, 8.4 Hz, 1H), 8.15 (s, 1H), 8.55 (dd, *J* = 1.8, 8.4 Hz, 1H), 9.05 (dd, *J* = 1.8, 3.9 Hz, 1H). MS (APCI⁺) (M+H)⁺ at *m/z* 530, 532, 534.

Example 225

10 (2-Isopropylphenyl)[2-nitro-4-(*E*-((3-carboethoxypiperidin-1-yl)carbonyl)ethenyl)phenyl] sulfide

Example 225A

(2-Isopropylphenyl)[2-nitro-4-(*E*-(carboxy)ethenyl)phenyl] sulfide

15 To a stirred mixture of 4-chloro-3-nitrocinnamic acid (500 mg, 2.2 mmol) in 5 mL of anhydrous DMF with K₂CO₃ (911 mg, 6.6 mmol) was added 2-isopropylbenzenethiol (372 mL, 2.2 mmol) in 1 mL of DMF dropwise. The resulting mixture was then heated at 70 °C under nitrogen atmosphere over night. Water (25 mL) was then added and the reaction mixture was acidified to pH = 4 with 3N HCl.

20 The cloudy mixture was extracted with EtOAc (2×20 mL). The combined organic layer was washed with brine, dried over Na₂SO₄, concentrated in vacuo to give the

title compound as viscous light-yellow oil, which was used for coupling with further purification.

Example 225B

5 (2-Isopropylphenyl)[2-nitro-4-(*E*-((3-carboethoxypiperidin-1-yl)carbonyl)ethenyl]
 phenyl] sulfide

The title compound was prepared by the procedures described in Example 92, substituting the benzoic acid with cinnamic acid from 225A, and ammonium chloride with ethyl nipecotate, giving a light-yellow solid. ¹H NMR (CDCl₃, 300 MHz) δ 1.18
10 (d, *J* = 6.6 Hz, 6H), 1.27 (t, *J* = 7.2 Hz, 3H), 1.69-1.82 (m, 1H), 1.82-1.99 (m, 1H),
1.99-2.20 (m, 1H), 2.45-2.62 (m, 2H), 3.45 (septet, *J* = 6.6 Hz, 1H), 3.56-3.80 (m,
1H), 3.80-4.10 (m, 2H), 4.16 (q, *J* = 7.2 Hz, 2H), 4.65-4.81 (m, 1H), 6.69 (d, *J* = 8.4
Hz, 1H), 7.00 (br s, 1H), 7.31 (dd, *J* = 2.4, 6.9 Hz, 1H), 7.42 (br d, *J* = 8.4 Hz, 1H),
7.51 (d, *J* = 15.6 Hz, 1H), 7.52 (overlapping d, 2H), 7.58 (d, *J* = 15.6 Hz, 1H), 8.43 (s,
15 1H). MS (APCI⁺) (M+H)⁺ at *m/z* 483.

Example 226

(2-Isopropylphenyl)[2-nitro-4-(*E*-((3-carboxypiperidin-1-yl)carbonyl)ethenyl]
 phenyl] sulfide

20 The title compound was prepared by the procedures described in Example 155,
substituting the ethyl ester from Example 137 with the ethyl ester from Example
225B, and KOH with NaOH, to give a light-yellow solid. ¹H NMR (d⁶-DMSO, 300

MHz) δ 1.15 (d, J = 6.9 Hz, 6H), 1.30-1.50 (m, 1H), 1.50-1.80 (m, 2H), 1.88-2.04 (m, 2H), 2.95-3.17 (m, 1H), 3.94-4.06 (m, 1H), 4.06-4.22 (m, 2H), 4.40-4.52 (m, 1H), 6.63 (d, J = 8.7 Hz, 1H), 7.33-7.53 (m, 3H), 7.56-7.68 (m, 3H), 7.91 (dd, J = 1.8, 8.4 Hz, 1H), 8.63 (d, J = 8.4 Hz, 1H). MS (APCI⁺) (M+H)⁺ at m/z 455.

5

Example 227

(2-Isopropylphenyl)[2-nitro-4-(*E*-(((3-ethanesulfonylaminocarbonyl)piperidin-1-yl)carbonyl)ethenyl) phenyl] sulfide

To a stirred solution of free acid (50 mg, 0.11 mmol) from Example 226 in 1 mL of methylene chloride was added ethyl sulfonamide (18 mg, 0.17 mmol), EDAC (25 mg, 0.13 mmol), and DAMP (2.7 mg, 0.022 mmol) sequentially. The mixture was stirred at ambient temperature for 16 h. The solvent was then removed on a rotavap under reduced pressure and the residue was purified on an Alltech sep-pak, eluting with 1% MeOH in EtOAc to give 30 mg (50 % yield) the title compound as a light yellow solid. ¹H NMR (CDCl₃, 300 MHz) δ 1.18 (d, J = 6.3 Hz, 6H), 1.34 (t, J = 7.5 Hz, 3H), 1.61-1.74 (m, 2H), 1.84-2.04 (m, 1H), 2.13-2.35 (m, 1H), 2.60-2.75 (m, 2H), 3.44 (p, J = 7.5 Hz, 2H), 3.53-3.66 (m, 1H), 3.66-3.85 (m, 2H), 4.00-4.18 (m, 1H), 6.71 (d, J = 8.7 Hz, 1H), 6.88 (d, J = 15.6 Hz, 1H), 7.31 (dd, J = 2.4, 8.4 Hz, 1H), 7.41 (d, J = 1.8, 8.4 Hz, 1H), 7.51 (d, J = 1.8 Hz, 1H), 7.54 (d, J = 8.4 Hz, 1H), 7.67 (d, J = 15.6 Hz, 1H), 8.43 (s, 1H). MS (ESI⁺) (M+H)⁺ at m/z 546.

Example 228

(2-Isopropylphenyl)[2-nitro-4-(*E*-(((3-(4-methylpiperazine)sulfonylaminocarbonyl)piperidin-1-yl)carbonyl)ethenyl)phenyl] sulfide

The title compound was prepared by the procedures described in Example 228, substituting ethyl sulfonamide with *N*-methylpiperazine sulfonamide, giving a light

5 yellow solid. ¹H NMR (CDCl₃, 300 MHz) δ 1.18 (d, *J* = 6.5 Hz, 6H), 1.40-2.10 (m, 9H), 2.60 (s, 3H), 2.60-2.76 (m, 4H), 2.90 (br s, 3H), 3.44 (septet, *J* = 6.5 Hz, 1H), 3.52-4.08 (m, 4H), 6.71 (d, *J* = 8.4 Hz, 1H), 6.95 (d, *J* = 15.6 Hz, 1H), 7.31 (d, *J* = 2.1, 8.4 Hz, 1H), 7.43-7.57 (m, 4H), 7.64 (d, *J* = 15.6 Hz, 1H), 8.44 (s, 1H). MS (ESI⁺) (M+H)⁺ at *m/z* 616. Anal. Calcd for C₂₉H₃₇N₅O₆S₂ · 1.13 H₂O: C, 54.76; H, 6.22; N, 11.01. Found: C, 54.78; H, 6.11; N, 10.87.

10

Example 229

(2-Isopropylphenyl)[2-nitro-4-(*E*-(((3-*p*-toluenesulfonylaminocarbonyl)piperidin-1-yl)carbonyl)ethenyl)phenyl] sulfide

15 The title compound was prepared by the procedures described in Example 228, substituting ethyl sulfonamide with *p*-toluenesulfonamide, giving a light yellow solid.

¹H NMR (CDCl₃, 300 MHz) δ 1.19 (d, *J* = 6.5 Hz, 6H), 1.75-1.94 (m, 2H), 2.05-2.24 (m, 1H), 2.40 (s, 3H), 2.48-2.60 (m, 2H), 3.45 (septet, *J* = 6.5 Hz, 1H), 3.50-3.85 (m, 3H), 3.85-4.12 (m, 1H), 6.72 (d, *J* = 8.4 Hz, 1H), 6.86 (d, *J* = 15.6 Hz, 1H), 7.27-7.34

20 (m, 2H), 7.43 (dd, *J* = 2.1, 8.4 Hz, 1H), 7.50 (overlapping d, 1H), 7.53 (d, *J* = 8.4 Hz, 2H), 7.55 (d, *J* = 8.4 Hz, 1H), 7.92 (d, *J* = 8.4 Hz, 2H), 8.44 (s, 1H). MS (ESI⁺) (M+H)⁺ at *m/z* 608.

Example 230

(2-Isopropylphenyl)[2-nitro-4-(E-((3-methyl-4-acetyl)piperazin-1-yl)carbonyl)ethenyl] phenyl] sulfide

5 Prepared according to the procedures of Example 71, giving a yellow solid.
¹H NMR (DMSO-d₆, 300MHz) δ 0.94-1.18 (m, 3H); 1.14 (d, J = 7.0 Hz, 6H); 1.98-2.08 (br m, 3H); 2.69-3.74 (br m, 4H); 4.02-4.65 (br m, 4H); 6.64 (d, J = 8.5 Hz, 1H); 7.31-7.63 (m, 6H); 7.88-7.96 (br m, 1H); 8.65 (br s, 1H). MS (APCI) (M+H)⁺ at m/z 468. Anal calcd for C₂₅H₂₉N₃S₂O₄·0.1H₂O: C, 63.91; H, 6.70; N, 8.94. Found: C, 63.54; H, 6.41; N, 8.67.

Example 231

(2-Hydroxyphenyl)-[2-chloro-4(E-[(morpholin-1-yl)carbonyl]ethenyl)phenyl]sulfide

The title compound was prepared according to the procedures of Example 1, giving a white solid, m.p. 157-158C. ¹H-NMR (CDCl₃, 300 MHz) δ 3.60-3.76 (m, 8H), 6.42 (s, 1H), 6.57 (d, J=9hz, 1H), 6.76 (d, J=15Hz, 1H), 6.99-7.04 (m, 1H), 7.10-7.20 (m, 2H), 7.42-7.55 (m, 4H). Anal. Calcd. for C₁₉H₁₈ClNO₃S: C, 60.71; h, 4.83; N, 3.73. Found: C, 60.48; H, 5.05; N, 3.69.

Example 232

(1-(Carboxymethyl)indol-5-yl)[2-chloro-4-(E-((4-acetyl)piperazin-1-yl)carbonyl)ethenyl] phenyl] sulfide

To a stirred solution of indole compound from Example 85 (35 mg, 0.080 mmol) in 1 mL of anhydrous DMSO was added crushed KOH (18 mg, 0.32 mmol). After 45 min, *t*-butyl bromoacetate (23.5 mL, 0.16 mmol) was added. The resulting mixture was stirred at ambient temperature for 10 h. Water was then added and the reaction mixture was acidified with 3 N HCl to pH = 3. The title compound (25 mg, 63 %) was collected through filtration and dried in vacuum oven, giving a white solid.

¹H NMR (d⁶-DMSO, 300 MHz) δ 2.04 (s, 3H), 3.38-3.80 (m, 8H), 4.59 (s, 2H), 6.45 (d, *J* = 3.0 Hz, 1H), 6.52 (d, *J* = 8.7 Hz, 1H), 7.21 (dd, *J* = 2.1, 8.7 Hz, 1H), 7.25 (d, *J* = 15.6 Hz, 1H), 7.38 (d, *J* = 15.6 Hz, 1H), 7.40 (d, *J* = 3.0 Hz, 1H), 7.47 (d, *J* = 8.4 Hz, 1H), 7.80 (d, *J* = 2.1 Hz, 1H), 7.97 (s, 1H). MS (ESI⁺) (M-H)⁺ at *m/z* 496, 498.

Example 233

(Benzodioxan-6-yl)[2-trifluoromethyl-4-(*E*-((4-acetyl)piperazin-1-yl)carbonyl)ethenyl] phenyl sulfide

The title compound was prepared by the procedures described in Example 84, substituting 2-bromothiophenol with 6-mercaptobenzenedioxane. white solid; ¹H NMR (CDCl₃, 300 MHz) δ 2.15 (s, 3H), 3.46-3.89 (m, 8H), 4.30 (dd, *J* = 2.1, 6.0 Hz, 4H), 6.84 (d, *J* = 15.0 Hz, 1H), 6.92 (d, *J* = 8.4 Hz, 1H), 6.97-7.10 (m, 3H), 7.42 (d, *J* = 8.4 Hz, 1H), 7.64 (d, *J* = 15.0 Hz, 1H), 7.77 (s, 1H). MS (ESI⁺) *m/z* 493 (M+H)⁺.

Example 234

(2-Isopropylphenyl)[2-nitro-4-(*E*-((3-(1-pyrrolidin-2-onyl)prop-1-ylamino)
carbonyl)ethenyl) phenyl] sulfide

Prepared according to the procedures of Example 71, giving a yellow solid. ¹H NMR (DMSO-d₆, 300MHz) δ 1.14 (d, J = 7.1 Hz, 6H); 1.58-1.68 (m, 2H); 1.85-1.97 (m, 2H); 2.18-2.24 (m, 2H); 3.10-3.22 (m, 4H); 3.30-3.39 (m, 3H); 6.65-6.72 (m, 2H); 7.32-7.45 (m, 2H); 7.57-7.62 (m, 3H); 7.76 (dd, J = 8.8, 2.0 Hz, 1H); 8.11-8.17 (m, 1H); 8.44 (d, J = 2.0 Hz, 1H). MS (APCI) (M+H)⁺ at m/z 468. Anal calcd for C₂₅H₂₉N₃S₂O₄·0.26CH₃COOCH₂CH₃: C, 63.77; H, 6.39; N, 8.57. Found: C, 63.46; H, 6.37; N, 8.90.

10

Example 235

(3-(2-Morpholinoethylamino)phenyl)[2-trifluoromethyl-4-(*E*-((4-acetyl)piperazin-1-yl)carbonyl)ethenyl) phenyl] sulfide

The title compound was prepared according to the procedures of Example 62, employing the compound of Example 103 as starting material. ¹H NMR (CDCl₃, 300 MHz) δ 7.78 (d, 1H, J = 1.4 Hz), 7.64 (d, 1H, J = 15.4 Hz), 7.42 (d, 1H, J = 8.8 Hz), 7.21 (t, 1H, J = 7.9 Hz), 7.12 (d, 1H, J = 8.5 Hz), 6.84 (d, 1H, J = 15.4 Hz), 6.82 (m, 1H), 6.76 (t, 1H, J = 1.8 Hz), 6.66 (m, 1H), 3.72 (m, 10H), 3.51-3.55 (m, 2H), 3.16 (t, 2H, J = 5.9 Hz), 2.64 (t, 2H, J = 5.9 Hz), 2.50 (m, 4H), 2.15 (s, 3H). MS (ESI) m/z 563.

20

Example 236

(2-Pyrrolidin-1-yl)phenyl)[2-nitro-4-(*E*-((4-acetyl)piperazin-1-yl)carbonyl)ethenyl)
phenyl] sulfide

The title compound was prepared according to the procedures of Example 62, employing the compound of Example 103 as starting material. ¹H NMR (CDCl₃, 300 MHz) δ 7.77 (s, 1H), 7.64 (d, 1H, J = 15.4 Hz), 7.40 (m, 1H), 7.22 (d, 1H, J = 7.8 Hz), 7.10 (d, 1H, J = 8.8 Hz), 6.82 (d, 1H, J = 15.3 Hz), 6.76 (d, 1H, J = 7.8 Hz), 6.70 (t, 1H, J = 2.0 Hz), 6.59 (dd, 1H, J = 2.4, 8.1 Hz), 3.61-3.79 (m, 6H), 3.51-3.54 (m, 2H), 3.28 (m, 4H), 2.14 (s, 3H), 2.01 (m, 4H). MS (ESI) *m/z* 504.

10

Example 237

(3-Bromophenyl)[2-nitro-4-(*E*-((3-carboethoxypyrrolidin-1-yl)carbonyl)ethenyl)
phenyl] sulfide

The title compound was prepared according to the procedures of Example 1. ¹H NMR (CDCl₃, 300 MHz) δ 8.40 (d, 1H, J = 1.5 Hz), 7.75 (m, 1H), 7.45 (m, 1H), 7.48-7.56 (m, 2H), 7.38 (t, 1H, J = 7.9 Hz), 7.00 (br, 1H), 6.87 (d, 1H, J = 9.5 Hz), 4.16 (q, 2H, J = 7.1 Hz), 3.99 (br, 2H), 3.70 (br, 1H), 3.30 (br, 1H), 3.00 (br, 1H), 2.55 (s, 1H), 2.10 (m, 1H), 1.89 (br, 1H), 1.85 (br, 1H), 1.27 (t, 3H, J = 7.0 Hz). MS (ESI) *m/z* 519, 521. Anal. Calcd for C₂₃H₂₃BrN₂O₅S · 0.19 H₂O: C, 52.84; H, 4.51; N, 5.36. Found: C, 52.85; H, 4.55; N, 5.28.

20

Example 238

(3-Bromophenyl)[2-nitro-4-(*E*-((4-carboethoxypyrrolidin-1-yl)carbonyl)ethenyl)
phenyl] sulfide

The title compound was prepared according to the procedures of Example 1.

¹H NMR (CDCl₃, 300 MHz) δ 8.41 (s, 1H), 7.75 (m, 1H), 7.62-7.67 (m, 2H), 7.53 (m,
 5 1H), 7.48 (d, 1H, *J* = 8.8 Hz), 7.38 (t, 1H, *J* = 7.9 Hz), 6.98 (br, 1H), 6.88 (d, 1H, *J* =
 8.5 Hz), 4.18 (q, 2H, *J* = 7.1 Hz), 3.64-78 (br, 4H), 3.55 (br, 4H), 1.29 (t, 3H, *J* = 7.0
 Hz). MS (ESI) *m/z* 520, 522.

Example 239

10 (2-(Hydroxymethyl)-benzodioxan-6-yl)[2-chloro-4-(*E*-((4-acetylpiperazin-1-
yl)carbonyl)ethenyl) phenyl] sulfide

The title compound was prepared by the procedures described in Example 85,
 substituting 5-iodoindole with a mixture of 2-hydroxymethyl-6-bromobenzodioxane
 and 2-hydroxymethyl-7-bromobenzodioxane, giving a white solid. ¹H NMR (CDCl₃,
 15 300 MHz, mixture of 3:2 regioisomers) δ 2.15 (s, 3H), 3.46-3.83 (m, 8H), 3.83-4.01
 (m, 2H), 4.10-4.42 (m, 4H), 6.75 (d, *J* = 8.4 Hz, 1H), 6.79 (d, *J* = 15.9 Hz, 1H), [6.95
 (d), 6.98 (d), *J* = 4.8 Hz, 1H in total], [7.04 (t), 7.07 (t), *J* = 1.5 Hz, 1H in total], [7.10
 (d), 7.11 (d), *J* = 2.4 Hz, 1H in total], 7.19 (d, *J* = 8.4 Hz, 1H), 7.53 (s, 1H), 7.58 (d, *J*
 = 15.6 Hz, 1H). MS (APCI⁺) (M+H)⁺ at *m/z* 489.

20

Example 240

(Benzodioxan-6-yl)[2-trifluoromethyl-4-(E-((3-(pyrrolidin-2-on-1-yl)prop-1-ylamino)carbonyl)ethenyl)phenyl]sulfide

The title compound was prepared by the procedures described in Example 233, substituting 1-acetylpipezazine with 3-aminopropyl-1-pyrrolidin-2-one, giving a white
 5 solid. ¹H NMR (CDCl₃, 300 MHz) δ 1.69-1.80 (m, 2H), 2.08 (p, *J* = 7.5 Hz, 2H), 2.44 (t, *J* = 7.5 Hz, 2H), 3.27-3.48 (m, 6H), 4.24-4.34 (m, 4H), 6.44 (d, *J* = 15.6 Hz, 1H), 6.90 (d, *J* = 8.4 Hz, 1H), 7.00 (d, *J* = 8.4 Hz, 1H), 7.01 (dd, *J* = 2.7, 8.4 Hz, 1H), 7.06 (d, *J* = 2.7 Hz, 1H), 7.08 (s, 1H), 7.40 (dd, *J* = 2.1, 8.4 Hz, 1H), 7.53 (d, *J* = 15.6 Hz, 1H), 7.75 (d, *J* = 2.1 Hz, 1H). MS (ESI⁺) (M+H)⁺ at *m/z* 507.

10

Example 241

(3-(Dimethylaminomethyl)indol-5-yl)[2-chloro-4-(E-((4-acetylpipezazin-1-yl)carbonyl)ethenyl)phenyl] sulfide

The title compound was prepared by the procedures described in Example 217,
 15 substituting the indole from 186 with the indole from Example 85, resulting in a white solid. ¹H NMR (CDCl₃, 300 MHz) δ 2.15 (s, 3H), 2.54 (s, 6H), 3.47-3.85 (m, 8H), 4.05 (s, 2H), 6.56 (d, *J* = 8.7 Hz, 1H), 6.77 (d, *J* = 15.6 Hz, 1H), 7.09 (d, *J* = 8.7 Hz, 1H), 7.36 (dd, *J* = 1.5, 8.7 Hz, 1H), 7.50 (d, *J* = 8.7 Hz, 1H), 7.52 (s, 2H), 7.56 (d, *J* = 15.6 Hz, 1H), 7.88 (s, 1H), 9.27 (s, 1H). MS (ESI⁺) (M+H)⁺ at *m/z* 497, 499. Anal.
 20 Calcd for C₂₆H₂₉ClN₄O₂S · 0.46 TFA · 1.72 MeOH: C, 56.89; H, 6.06; N, 9.27.
 Found: C, 56.83; H, 6.15; N, 9.46.

Example 242

(2-Isopropylphenyl)[2-nitro-4-(*E*-((2-carboethoxypiperidin-1-yl)carbonyl)ethenyl)
phenyl] sulfide

The title compound was prepared by the procedures described in Example 225,
 5 substituting ethyl nipecotate with ethyl pipecolate, giving a light-yellow solid. ¹H
 NMR (CDCl₃, 300 MHz) δ 1.18 (d, *J* = 6.9 Hz, 6H), 1.28 (t, *J* = 7.35 Hz, 3H), 1.34-
 1.62 (m, 2H), 1.62-1.84 (m, 3H), 2.32 (br d, *J* = 13.2 Hz, 1H), 3.33-3.54 (m, 1H), 3.45
 (septet, *J* = 6.9 Hz, 1H), 3.99 (br d, *J* = 13.2 Hz, 1H), 4.21 (q, *J* = 7.35 Hz, 2H), 5.46
 (br s, 1H), 6.69 (d, *J* = 8.7 Hz, 1H), 7.01 (d, *J* = 15.6 Hz, 1H), 7.25-7.34 (m, 1H), 7.42
 10 (d, *J* = 8.7 Hz, 1H), 7.46-7.60 (m, 3H), 7.58 (d, *J* = 15.6 Hz, 1H), 8.44 (s, 1H). MS
 (ESI⁺) (M+H)⁺ at *m/z* 483.

Example 243

(2-Isopropylphenyl)[2-nitro-4-(*E*-((2-carboxypiperidin-1-yl)carbonyl)ethenyl)
phenyl] sulfide

15 The title compound was prepared by the procedures described in Example 226,
 substituting the ethyl ester from Example 225 with the ethyl ester from Example 242,
 giving a light-yellow solid. ¹H NMR (CDCl₃, 300 MHz) δ 1.18 (d, *J* = 6.9 Hz, 6H),
 1.40-1.89 (m, 5H), 2.34 (br d, *J* = 11.7 Hz, 1H), 3.31-3.51 (m, 1H), 3.44 (septet, *J* =
 20 6.9 Hz, 1H), 4.01 (d, *J* = 11.7 Hz, 1H), 5.42 (br s, 1H), 6.70 (d, *J* = 7.8 Hz, 1H), 6.99
 (br d, *J* = 15.6 Hz, 1H), 7.29 (td, *J* = 2.7, 6.9 Hz, 1H), 7.41 (d, *J* = 7.8 Hz, 1H), 7.45-
 7.58 (m, 3H), 7.64 (d, *J* = 15.6 Hz, 1H), 8.43 (s, 1H). MS (ESI⁺) (M+H)⁺ at *m/z* 455.

20

20

20

The title compound was prepared by the procedures described in Example 226, substituting the ethyl ester from Example 225 with the ethyl ester from Example 244, producing a light-yellow solid. ¹H NMR (CDCl₃, 300 MHz) δ 1.18 (d, *J* = 6.9 Hz, 6H), 1.65-1.89 (m, 2H), 1.97-2.14 (m, 2H), 2.59-2.74 (m, 1H), 2.93-3.20 (m, 1H),

3.20-3.42 (m, 1H), 3.44 (septet, $J = 6.9$ Hz, 1H), 3.97-4.18 (m, 1H), 4.40-4.65 (m, 1H), 6.70 (d, $J = 8.7$ Hz, 1H), 6.97 (d, $J = 15.6$ Hz, 1H), 7.30 (td, $J = 2.7, 6.9$ Hz, 1H), 7.41 (d, $J = 8.7$ Hz, 1H), 7.46-7.65 (m, 3H), 7.60 (d, $J = 15.6$ Hz, 1H), 8.43 (s, 1H). MS (ESI⁺) (M+H)⁺ at m/z 455.

5

Example 246

(2-Isopropylphenyl)[2-nitro-4-(*E*-((4-*p*-toluenesulfonylamino)carbonyl)piperidin-1-yl)carbonyl)ethenyl] phenyl] sulfide

The title compound was prepared by the procedures described in Example 229, substituting the acid from Example 226 with the acid from Example 245. light-yellow solid; ¹H NMR (d⁶-DMSO, 300 MHz) δ 1.14 (d, $J = 6.9$ Hz, 6H), 1.18-1.39 (m, 2H), 1.67-1.79 (m, 2H), 2.39 (s, 3H), 2.60-2.75 (m, 1H), 2.96-3.14 (m, 1H), 3.26-3.42 (m, 1H), 3.34 (septet, $J = 6.9$ Hz, 1H), 4.10-4.42 (m, 2H), 6.62 (d, $J = 8.4$ Hz, 1H), 7.32-7.43 (m, 4H), 7.45 (d, $J = 15.6$ Hz, 1H), 7.58 (d, $J = 8.4$ Hz, 2H), 7.60 (d, $J = 3.6$ Hz, 1H), 7.78 (d, $J = 8.4$ Hz, 2H), 7.87 (dd, $J = 2.7, 8.4$ Hz, 1H), 8.60 (d, $J = 2.7$ Hz, 1H). MS (ESI⁺) (M+H)⁺ at m/z 606. Anal. Calcd for C₃₁H₃₃N₃O₆S₂ · 0.26 H₂O: C, 60.80; H, 5.52; N, 6.86. Found: C, 60.85; H, 5.84; N, 6.61.

Example 247

(2-Isopropylphenyl)[2-nitro-4-(*E*-((3-carboxy-4-hydroxypiperidin-1-yl)carbonyl)ethenyl] phenyl] sulfide

20

- Prepared according to the procedures of Example 71, giving a yellow solid. ¹H NMR (DMSO-d₆, 300MHz) δ 1.14 (d, J = 6.8 Hz, 6H); 1.53-1.70 (br m, 2H); 2.92-3.52 (br m, 1H); 3.30-3.40 (m, 1H); 3.98-4.44 (br m, 4H); 4.90-5.20 (br m, 1H); 6.63 (d, J = 8.5 Hz, 1H); 7.34-7.62 (m, 6H); 7.87-7.94 (br m, 1H); 8.58-8.64 (br m, 1H).
- 5 MS (APCI) (M+H)⁺ at m/z 471. Anal calcd for C₂₄H₂₆N₂S₂O₆: C, 61.26; H, 5.57; N, 5.95. Found: C, 61.05; H, 5.85; N, 5.73.

Example 248

- (Benzodioxan-6-yl)[2-trifluoromethyl-4-(*E*-((3-carboethoxypiperidin-1-yl) carbonyl)ethenyl) phenyl] sulfide
- 10

- The title compound was prepared by the procedures described in Example 240 substituting N-(3'-aminopropyl)-2-pyrrolidinone with ethyl nipecotate, giving a white hygroscopic solid. ¹H NMR (CDCl₃, 300 MHz) δ 1.26 (t, J=7.0 Hz, 3H), 1.54 (m, 1H), 1.65-1.80 (m, 2H), 2.10 (m, 1H), 2.54 (m, 1H), 2.92-3.40 (m, 2H), 3.60-4.10 (m, 2H), 4.14 (q, J=7.0 Hz, 2H), 4.25-4.32 (m, 4H), 6.91 (d, J=7.5 Hz, 1H), 7.00 (dd, J=2.0, 15.0 Hz, 3H), 7.05 (d, J=2.0 Hz, 1H), 7.40 (d, J=8.0, 1H), 7.56 (d, J=15.0 Hz, 1H), 7.76 (s, 1H). MS (CI/NH₃) m/z 522 (M+H)⁺. Anal. calcd. for C₂₆H₂₆F₃NO₅S: C, 59.88; H, 5.02; N, 2.69. Found: C, 59.92; H, 5.39; N, 2.56.
- 15

20

Example 249

(Benzodioxan-6-yl)[2-trifluoromethyl-4-(*E*-((2-carboethoxypiperidin-1-yl) carbonyl)ethenyl) phenyl] sulfide

The title compound was prepared by the procedures described in Example 240 substituting N-(3'-amino~~propyl~~)~~2~~-pyrrolidinone with ethyl pipercolinate. ¹H NMR (CDCl₃, 300 MHz) δ 1.28 (t, J=7.0 Hz, 3H), 1.35-1.54 (m, 2H), 1.64-1.82 (m, 3H), 2.30 (m, 1H), 3.40 (m, 1H), 4.00 (m, 1H), 4.22 (q, J=7.0 Hz, 2H), 4.26-4.34 (m, 4H), 5.48 (m, 1H), 6.91 (d, J=8.5 Hz, 1H), 6.98 (m, 1H), 7.02 (dd, J=2.0, 8.0 Hz, 2H), 7.06 (d, J=2.0 Hz, 1H), 7.41 (d, J=8.0 Hz, 1H), 7.57 (d, J=15.0 Hz, 1H), 7.77 (s, 1H). MS (Cl/NH₃) m/z 522 (M+H)⁺. Anal. calcd for C₂₆H₂₆F₃NO₅S: C, 59.88; H, 5.02; N, 2.69. Found: C, 60.25; H, 5.12; N, 2.55.

Example 250

(Benzodioxan-6-yl)[2-nitro-4-(*E*-((4-carboxypiperidin-1-yl) carbonyl)ethenyl) phenyl] sulfide

The title compound was prepared by the hydrolysis of compound 198 under basic condition (aq. NaOH/EtOH), and purified by reversed-phase HPLC. ¹H NMR (DMSO-d₆, 300 MHz) δ 1.44 (m, 2H), 1.78 (m, 2H), 2.04 (m, 2H), 2.82 (m, 1H), 4.02-4.20 (m, 2H), 4.420-4.35 (m, 4H), 6.90 (d, J=8.0 Hz, 1H), 6.97 (d, J=8.0 Hz, 1H), 7.05(dd, J=2.0, 8.0 Hz, 1H), 7.10(d, J=2.0 Hz, 1H), 7.15 (br, 1H), 7.44 (m 1H), 7.60 (br, 1H), 8.40 (s, 1H). MS (ESI) m/z 469 (M-1)⁻.

Example 251

(Benzodioxan-6-yl)[2-trifluoromethyl-4-(*E*-((3-carboxypyrrolidin-1-yl)carbonyl)ethenyl) phenyl] sulfide

The title compound was prepared according to the procedures of

Example 1. ¹H NMR (CDCl₃, 300 MHz) δ 7.75 (s, 1H), 7.60 (d, 1H, J = 15.0 Hz) 7.40 (br, 1H), 7.06 (d, 1H, J = 2.2 Hz), 6.96-7.02 (m, 3H), 6.90 (d, 1H, J = 8.5 Hz), 4.30 (m, 5H), 3.99 (br, 2H), 3.29 (br, 2H), 2.60 (br, 2H), 1.85 (br, 2H). MS (ESI) *m/z* -492.

Example 252

(Benzodioxan-6-yl)[2-trifluoromethyl-4-(*E*-(4-carboethoxypiperidin-1-yl) carbonyl)ethenyl] phenyl] sulfide

The title compound was prepared by the procedures described in Example 240 substituting N-(3'-aminopropyl)-2-pyrrolidinone with ethyl isonipecotate, giving a white sticky solid. ¹H NMR (CDCl₃, 300 MHz) δ 1.26 (t, J=7.0 Hz, 3H), 1.68-1.80 (m, 2H), 1.98-2.10 (, 2H), 2.54-2.70 (m, 2H), 3.00-3.30 (br, 2H), 4.15 (m, 3H), 4.26-4.34 (m, 4H), 6.90 (d, J=8.0 Hz, 2H), 7.00 (dd, J=2.0, 8.0 Hz, 2H), 7.06 (d, J=2.0 Hz, 1H), 7.41 (m, 1H), 7.50 (br, 1H), 7.75 (s, 1H). MS (CI/NH₃) *m/z* 522 (M+H)⁺. Anal. calcd. for C₂₄H₂₂F₃NO₅S• 0.1 H₂O: C, 58.20; H, 4.52; N, 2.83. Found: C, 58.14; H, 4.69; N, 2.76.

Example 253

(Benzodioxan-6-yl)[2-trifluoromethyl-4-(*E*-(2-carbomethoxy-4-*tert*-butoxycarbonylpiperazin-1-yl) carbonyl)ethenyl] phenyl] sulfide

The title compound was prepared by the procedures described in Example 240 substituting N-(3'-aminopropyl)-2-pyrrolidinone with 1-Boc-3-carbomethoxypiperazine, giving a white solid, mp 85-87 °C. ¹H NMR (CDCl₃, 300 MHz) δ 1.46 (s, 9H), 2.90-3.00 (m, 2H), 3.08-3.20 (m, 2H), 3.76 (s, 3H), 3.90 (m, 1H), 4.25-4.34 (m, 4H), 4.58-4.66 (m, 2H), 6.92 (d, J=8.0 Hz, 1H), 6.98 (m, 1H), 7.02 (dd, J=2.0, 8.0 Hz, 2H), 7.06 (d, J=2.0 Hz, 1H), 7.40 (m, 1H), 7.62 (br, 1H), 7.76 (s, 1H). MS (APCI) m/z 609 (M+H)⁺. Anal. calcd. for C₂₉H₃₁F₃N₂O₇S: C, 57.23; H, 5.13; N, 4.60. Found: C, 57.09; H, 5.25; N, 4.37.

Example 254

(Benzodioxan-6-yl)[2-trifluoromethyl-4-(E-((2-carbomethoxy-4-methoxycarbonylpiperazin-1-yl) carbonyl)ethenyl) phenyl] sulfide

The title compound was prepared by treating the compound of Example 255 with methyl chloroformate and pyridine in CH₂Cl₂ at room temperature, producing a white foam. ¹H NMR (CDCl₃, 300 MHz) δ 3.00 (m, 1H), 3.18 (m, 1H), 3.60 (m, 1H), 3.72 (s, 3H), 3.76 (s, 3H), 3.90 (m, 1H), 4.10 (br, 1H), 4.28-4.34 (m, 4H), 4.64 (m, 1H), 5.32 (m, 1H), 6.85 (d, J=15.5 Hz, 1H), 6.92 (d, J= 8.0 Hz, 1H), 6.98 (m, 1H), 7.02 (dd, J=2.0, 8.0 Hz, 1H), 7.08 (d, J=2.0 Hz, 1H), 7.40 (d, J=8.0 Hz, 1H), 7.64 (d, J=15.0 Hz, 1H), 7.77 (s, 1H). MS (CI/NH₃) m/z 567 (M+H)⁺. Anal. calcd. for C₂₆H₂₅F₃N₂O₇S: C, 55.12; H, 4.45; N, 4.94. Found: C, 55.18; H, 4.70; N, 4.68.

Example 255

(Benzodioxan-6-yl)[2-trifluoromethyl-4-(*E*-((2-carbomethoxypiperazin-1-yl)carbonyl)ethenyl) phenyl] sulfide

The title compound was prepared by deprotection of compound 253 with TFA in CH₂Cl₂, resulting in a light yellow solid, mp 70-72 °C. ¹H NMR (CDCl₃, 300 MHz)

5 δ 2.90 (m, 1H), 3.05 (m, 1H), 3.35 (m, 1H), 3.68 (m, 1H), 3.80 (s, 3H), 4.00 (m, 1H), 4.25-4.34 (m, 4H), 4.70 (br, 1H), 5.46 (m, 1H), 6.84 (d, J=15.5 Hz, 1H), 6.90 (d, J=8.0 Hz, 1H), 6.96-7.04 (m, 2H), 7.06 (m, 1H), 7.40 (d, J=8.0 Hz, 1H), 7.65 (d, J=15.5 Hz, 1H), 7.77 (s, 1H). MS (CI/NH₃) m/z 509 (M+H)⁺. Anal. calcd. for C₂₄H₂₃F₃N₂O₅S•1.55 H₂O: C, 53.74; H, 4.90; N, 5.22. Found: C, 54.04; H, 4.59; N,

10 4.82.

Example 256

(2-Methyl-3-(carboethoxymethyl)indol-5-yl)[2-trifluoromethyl-4-(*E*-((morpholin-1-yl)carbonyl)ethenyl) phenyl] sulfide

15

Example 256A

(4-Bromophenyl)[2-trifluoromethyl-4-(*E*-((morpholin-1-yl)carbonyl)ethenyl) phenyl] sulfide

The bromide was prepared by the procedure described in Example 12,

20 substituting 2-bromothiophenol with 4-bromothiophenol, and 3,4-dichlorobenzaldehyde with 4-fluoro-3-trifluoromethylbenzaldehyde.

00544795-033100
00100-56774560

Example 256B

(4-Hydrazinophenyl)[2-trifluoromethyl-4-(E-((morpholin-1-yl)carbonyl)ethenyl) phenyl] sulfide, benzophenone hydrazone

To a stirred solution of above-described bromide (1.0 g, 2.12 mmol) in 10 mL
5 of toluene with Pd(OAc)₂ (9.5 mg, 0.04 mmol), BINAP (40 mg, 0.06 mmol), and
benzophenone hydrazone (437 mg, 2.12 mmol) was added NaOt-Bu (285 mg, 2.97
mmol). The reaction mixture was bubbled with N₂ for 2 min before it was heated at 80
°C for 4 h. The reaction mixture was then allowed to cool down to ambient
temperature. Ether was then added and the mixture was filtered through celite, washed
10 with diethyl ether. The filtrate was concentrate in vacuo and the residue was purified
on a SiO₂ flash column chromatography eluting with 10-30% EtOAc/hexanes to give
170 mg (13%) of the title compound as light brown foamy solid.

Example 256C

15 (2-Methyl-3-(carboethoxymethyl)indol-5-yl)[2-trifluoromethyl-4-(E-((morpholin-1-yl)carbonyl)ethenyl) phenyl] sulfide

To a stirred solution of hydrazone (90 mg, 0.15 mmol) in 2 mL of ethanol was
added levulinic acid (24 mL, 23 mmol) and *p*-TsOH (146 mg, 0.75 mmol). The
mixture was then refluxed for 2 days. After cooled down to ambient temperature, the
20 reaction mixture was partitioned between EtOAc and sat. NaHCO₃. The organic layer
was then washed with brine, dried over Na₂SO₄, concentrated in vacuo. The residue
was then purified on Gilson preparative HPLC as described in Example 38B to give

- 6.0 mg (7%) of the title compound. light-brown solid. ^1H NMR (CDCl_3 , 300 MHz) δ 1.20 (t, $J = 7.4$ Hz, 3H), 2.46 (s, 3H), 3.55-3.83 (br m, 8H), 3.67 (s, 2H), 4.12 (q, $J = 7.4$ Hz, 2H), 6.79 (d, $J = 15.3$ Hz, 1H), 6.84 (d, $J = 8.4$ Hz, 1H), 7.23-7.31 (m, 2H), 7.34 (d, $J = 8.4$ Hz, 1H), 7.60 (d, $J = 15.3$ Hz, 1H), 7.76 (s, 1H), 7.80 (s, 1H), 8.04 (s, 1H). MS (ESI $^+$) ($\text{M}+\text{H}$) $^+$ at m/z 533.

Example 257

(1-(2-Methoxyethyl)indol-5-yl)[2-chloro-4-(*E*-((4-acetyl)piperazin-1-yl)(carbonyl)ethenyl) phenyl] sulfide

- 10 The title compound was prepared by the procedures described in Example 232, substituting *t*-butyl bromoacetate with bromoethylmethyl ether. white solid; ^1H NMR (CDCl_3 , 300 MHz) δ 2.14 (s, 2H), 3.35 (s, 3H), 3.46-3.56 (m, 2H), 3.56-3.80 (m, 6H), 3.75 (t, $J = 5.6$ Hz, 2H), 4.33 (t, $J = 5.6$ Hz, 2H), 6.54 (d, $J = 3.3$ Hz, 1H), 6.61 (d, $J = 8.7$ Hz, 1H), 6.75 (d, $J = 15.3$ Hz, 1H), 7.09 (dd, $J = 2.1, 11.7$ Hz, 1H), 7.26
- 15 (overlapping d, 1H), 7.36 (dd, $J = 2.1, 8.7$ Hz, 1H), 7.44 (d, $J = 8.7$ Hz, 1H), 7.51 (d, $J = 2.1$ Hz, 1H), 7.56 (d, $J = 15.3$ Hz, 1H), 7.88 (d, $J = 1.5$ Hz, 1H). MS (ESI $^+$) ($\text{M}+\text{H}$) $^+$ at m/z 498, 500.

Example 258

- 20 (2-Isopropylphenyl)[2-nitro-4-(*E*-((3-acetoxymethyl-4-hydroxypiperidin-1-yl)(carbonyl)ethenyl) phenyl] sulfide

Prepared according to the procedures of Example 71, giving a yellow solid.

¹H NMR (DMSO-d₆, 300MHz) δ 1.14 (d, J = 7.0 Hz, 6H); 1.51-1.90 (br m, 2H); 1.92-2.06 (m, 3H); 2.50-3.21 (br m, 2H); 3.30-3.40 (m, 1H); 3.40-4.44 (br m, 5H); 4.88-4.97 (br m, 1H); 6.63 (d, J = 8.5 Hz, 1H); 7.31-7.62 (m, 6H); 7.87-7.94 (br m, 1H);

5 8.58-8.64 (br m, 1H). MS (APCI) (M+H)⁺ at m/z 499. Anal calcd for C₂₆H₃₀N₂S₇O₆·0.29H₂O: C, 61.98; H, 6.12; N, 5.56. Found: C, 62.00; H, 6.35; N, 5.55.

Example 259

(2-Isopropylphenyl)[2-nitro-4-(*E*-((3-(dimethylaminocarbonyl)-4-hydroxypiperidin-1-yl)carbonyl)ethenyl) phenyl] sulfide

10

Prepared according to the procedures of Example 71, giving a yellow solid.

¹H NMR (DMSO-d₆, 300MHz) δ 1.14 (d, J = 6.8 Hz, 6H); 1.54-1.75 (br m, 2H); 2.81, 2.82 (br s, br s, 3H); 3.00, 3.04 (br s, br s, 3H); 2.75-3.60 (br m, 3H); 3.30-3.40 (m, 1H); 3.90-4.28 (br m, 2H); 4.95-5.28 (br m, 1H); 6.61-6.66 (m, 1H); 7.34-7.62 (m,

15 6H); 7.87-7.94 (br m, 1H); 8.58-8.63 (br m, 1H). MS (ESI) (M+H)⁺ at m/z 498. Anal calcd for C₂₆H₃₁N₅S₇O₅·0.34H₂O: C, 61.99; H, 6.34; N, 8.34. Found: C, 61.96; H, 6.37; N, 8.56.

Example 260

(2-Isopropylphenyl)[2-nitro-4-(*E*-((3-cyanomorpholin-1-yl)carbonyl)ethenyl) phenyl] sulfide

20

Prepared according to the procedures of Example 71, giving a yellow solid.

¹H NMR (DMSO-d₆, 300MHz) δ 1.14 (d, J = 6.8 Hz, 6H); 3.30-3.40 (m, 1H); 3.30-4.16 (br m, 5H); 4.20-4.29 (br m, 1H); 5.07 (t, J = 3.5 Hz, 1H); 6.65(d, J = 8.8 Hz, 1H); 7.32-7.44 (m, 2H); 7.54-7.62 (m, 4H); 7.91 (dd, J = 8.8, 2.0 Hz, 1H); 8.67 (d, J = 2.0 Hz, 1H). MS (APCI) (M+H)⁺ at m/z 438. Anal calcd for C₂₃H₂₃N₃S₁O₄·0.25C₈H₁₄: C, 64.11; H, 5.82; N, 9.15. Found: C, 63.99; H, 6.00; N, 9.12.

Example 261

(2-Isopropylphenyl)[2-nitro-4-(*E*-((3-carboethoxymorpholin-1-yl)carbonyl)ethenyl)phenyl] sulfide

Prepared according to the procedures of Example 71, giving a yellow solid.

¹H NMR (DMSO-d₆, 300MHz) δ 1.14 (d, J = 7.0 Hz, 6H); 1.12-1.27 (m, 3H); 3.30-3.40 (m, 1H); 3.15-4.33 (br m, 9H); 6.64 (d, J = 8.5 Hz, 1H); 7.32-7.42 (m, 2H); 7.50-7.62 (m, 4H); 7.88-7.96 (br m, 1H); 8.65 (br s, 1H). MS (APCI) (M+H)⁺ at m/z 485. Anal calcd for C₂₅H₂₈N₂S₁O₆: C, 61.97; H, 5.82; N, 5.78. Found: C, 61.83; H, 6.07; N, 5.74.

Example 262

(2-Isopropylphenyl)[2-nitro-4-(*E*-((3-(tetrazol-5-yl)morpholin-1-yl)carbonyl)ethenyl)phenyl] sulfide

The compound of Example 260 (160 mg, 0.336), sodium azide (56.6 mg,

0.872 mmol), n-Bu₃SnCl and THF were mixed in a reaction tube, flushed with

nitrogen and heated to reflux overnight. The mixture was then cooled to ambient temperature and 1N HCl soln. was added. The mixture was extracted with ethyl acetate three times and the combined organics were dried over MgSO_4 . The mixture was filtered through a short silica gel plug to give 96 mg (56% yield) of the desired material. ^1H NMR ($\text{DMSO}-d_6$, 300MHz) δ 1.14 (d, J = 6.8 Hz, 6H); 2.96-4.62 (br m, 7H); 4.77 (dd, J = 10.5, 2.7 Hz, 1H); 6.58-6.67 (m, 1H); 7.32-7.62 (m, 6H); 7.92 (dd, J = 8.8, 2.0 Hz, 1H); 8.62-8.67 (br m, 1H). MS (APCI) $(\text{M}+\text{H})^+$ at m/z 481. Anal. calcd for $\text{C}_{23}\text{H}_{24}\text{N}_6\text{S}_4\text{O}_4 \cdot 1.2\text{H}_2\text{O}$: C, 54.93; H, 5.31; N, 16.71. Found: C, 54.97; H, 5.12; N, 16.50.

10

Example 263

(Benzodioxan-6-yl)[2-trifluoromethyl-4-(*E*-((4-carboxypiperidin-1-yl)carbonyl)ethenyl) phenyl] sulfide

The title compound was prepared by hydrolysis of the compound of Example 252 under basic conditions (aq. NaOH/EtOH), giving a white solid, mp 88 °C (dec.). ^1H NMR ($\text{DMSO}-d_6$, 300 MHz) δ 1.40 (m, 2H), 1.98 (m, 2H), 2.95 (m, 1H), 3.15 (m, 1H), 3.45 (m, 1H), 4.20 (m, 2H), 4.35 (m, 4H), 7.00 (m, 4H), 7.20 (m, 2H), 7.90 (m, 1H), 8.20 (m, 1H), 12.30 (s, 1H). MS (APCI) m/z 494 $(\text{M}+\text{H})^+$. Anal. calcd. for $\text{C}_{24}\text{H}_{22}\text{F}_3\text{N}_3\text{O}_5 \cdot 0.1\text{H}_2\text{O}$: C, 58.20; H, 4.52; N, 2.83. Found: C, 58.14; H, 4.69; N, 2.76.

20

Example 264

(Benzodioxan-6-yl)[2-trifluoromethyl-4-(*E*-((2-carboxypiperidin-1-yl)
carbonyl)ethenyl) phenyl] sulfide

- The title compound was prepared by hydrolysis of the compound of Example 249 under basic conditions (aq. NaOH/EtOH), resulting in a white solid, mp 90 °C
- 5 (dec.). ¹H NMR (DMSO-d₆, 300 MHz) δ 1.15-1.50 (m, 2H), 1.50-1.70 (m, 2H), 2.16 (m, 1H), 2.56 (m, 1H), 3.15 (m, 1H), 4.30 (s, 4H), 4.32 (m, 1H), 5.20 (m, 1H), 7.02 (m, 4H), 7.30-7.52 (m, 2H), 7.84 (m, 1H), 8.15 (s, 1H). MS (APCI) *m/z* 494 (M+H)⁺. Anal. calcd. for C₂₄H₂₂F₃NO₅•0.3 H₂O: C, 57.78; H, 4.57; N, 2.81. Found: C, 57.87; H, 4.5
- 10 7; N, 2.76.

Example 265

(Benzodioxan-6-yl)[2-trifluoromethyl-4-(*E*-((4-carbomethoxypiperazin-1-yl)
carbonyl)ethenyl) phenyl] sulfide

- 15 The title compound was prepared according to the procedures of Example 1.
- ¹H NMR (CDCl₃, 300 MHz) δ 7.76 (s, 1H), 7.62 (d, 1H, *J* = 15.0 Hz), 7.40 (d, 1H, *J* = 8.6 Hz), 7.06 (d, 1H, *J* = 2.1 Hz), 6.98-7.04 (m, 2H), 6.91 (d, 1H, *J* = 8.4 Hz), 6.84 (d, 1H, *J* = 15.6 Hz), 4.31 (m, 4H), 4.18 (q, 2H, *J* = 7.1 Hz), 3.68 (br, 4H), 3.54 (br s, 4H), 1.29 (t, 3H, *J* = 7.2 Hz). MS (ESI) *m/z* 523, 545, 1045, 1067.

Example 266

(Benzodioxan-6-yl)[2-trifluoromethyl-4-(E-((3-aza-6,9-dioxaspiro[5.4]decan-1-yl)carbonyl)ethenyl) phenyl] sulfide

The title compound was prepared according to the procedures of Example 1.

¹H NMR (DMSO-d₆, 300 MHz) δ 8.13 (s, 1H), 7.84 (d, 1H, J = 9.0 Hz), 7.48 (d, 1H, J = 15.4 Hz) 7.38 (d, 1H, J = 15.4 Hz), 6.98-7.06 (m, 4H), 4.30 (m, 4H), 3.92 (s, 4H), 3.74 (br, 2H), 2.62 (br, 2H), 1.63 (br, 4H). MS (ESI) *m/z* 508, 1015.

Example 267

(Benzodioxan-6-yl)[2-trifluoro-4-(E-((4-(benzimidazolone-1-yl)piperidin-1-yl)carbonyl)ethenyl) phenyl] sulfide

The title compound was prepared according to the procedures of Example 1.

¹H NMR (CDCl₃, 300 MHz) δ 8.32 (s, 1H), 7.79 (s, 1H), 7.66 (d, 1H, J = 15.4 Hz), 7.44 (d, 1H, J = 8.5 Hz), 7.0-7.12 (m, 6H), 6.94 (d, 1H, J = 9.9 Hz), 6.90 (d, 1H, J = 2.6 Hz), 4.98 (m, 1H), 4.59 (m, 1H), 4.20 (m, 5H), 3.31 (br, 1H), 2.83 (br, 1H), 2.40 (m, 2H), 1.98 (m, 2H). MS (ESI) *m/z* 582, 604, 1163, 1185.

Example 268

(Benzodioxan-6-yl)[2-trifluoromethyl-4-(E-((4-(methylaminocarbonyl)piperidin-1-yl)carbonyl)ethenyl) phenyl] sulfide

The title compound was prepared according to the procedures of Example 1.

¹H NMR (CDCl₃, 300 MHz) δ 7.75 (s, 1H), 7.67 (d, 1H, J = 15.4 Hz) 7.40 (d, 1H, J = 8.1 Hz), 7.06 (d, 1H, J = 2.4 Hz), 6.96-7.02 (m, 2H), 6.90 (d, 1H, J = 8.2 Hz), 4.28

(m, 4H), 3.95 (br, 2H), 3.50 (m, 1H), 2.82 (s, 3H), 2.40 (m, 1H), 2.15 (br, 1H), 1.88 (br, 1H), 1.73 (br, 2H). MS (ESI) m/z 507, 529, 1035.

Example 269

5 (Benzodioxan-6-yl)[2-trifluoromethyl-4-(*E*-((3-carbomethoxy-4-methoxycarbonylpiperazin-1-yl) carbonyl)ethenyl) phenyl] sulfide

The title compound was prepared by the procedures described in Example 240 substituting N-(3'-aminopropyl)-2-pyrrolidinone with 2-carbomethoxy-1-methoxycarbonylpiperazine, producing a light yellow solid, mp 56 °C (dec.). ¹H

10 NMR (CDCl₃, 300 MHz) δ 2.70-3.50 (br, 4H), 3.70 (s, 3H), 3.76 (d, J=9.0 Hz, 3H), 4.00(m, 1H), 4.20 (m, 4H), 4.50-5.00 (br, 2H), 6.91 (d, J=8.5 Hz, 1H), 6.92-7.02 (m, 2H), 7.07 (d, J=2.0 Hz, 1H), 7.25 (m, 1H), 7.40 (m, 1H), 7.60 (m, 1H), 7.72 (s, 1H). MS (APCI) m/z 567 (M+H)⁺. Anal. calcd. for C₂₆H₂₅F₃N₂O₇S: C, 55.12; H, 4.45; N, 4.94. Found: C, 55.33; H, 4.74; N, 4.76.

15

Example 270

(2-Isopropylphenyl)[2-nitro-4-(*E*-((3-carboxymorpholin-1-yl)carbonyl)ethenyl) phenyl] sulfide

Prepared according to the procedures of Example 71, giving a yellow solid. ¹H

20 NMR (DMSO-d₆, 300MHz) δ 1.14 (d, J = 6.8 Hz, 6H); 3.08-4.33 (br m, 7H); 3.30-3.40 (m, 1H); 6.58-6.68 (m, 1H); 7.32-7.66 (m, 6H); 7.87-7.94 (m, 1H); 8.53-8.65

(m, 1H). MS (APCI) (M+H)⁺ at m/z 457. Anal calcd for C₂₃H₂₄N₂S₁O₆: C, 60.51; H, 5.30; N, 6.14. Found: C, 60.33; H, 5.54; N, 5.80.

Example 271

5 (Benzodioxan-6-yl)[2-trifluoromethyl-4-(*E*-(2-carboxy-4-methoxycarbonylpiperazin-1-yl) carbonyl)ethenyl] phenyl] sulfide

The title compound was prepared by treating the compound of Example 255 with methyl chloroformate and pyridine in CH₂Cl₂ at room temperature, and followed by hydrolysis under basic conditions (aq. NaOH/EtOH), producing a white solid, mp
10 102 °C (dec.). ¹H NMR (DMSO-d₆, 300 MHz) δ 2.85 (m, 1H), 3.02 (m, 1H), 3.20 (m, 1H), 3.40 (m, 1H), 3.62 (s, 3H), 3.88 (m, 1H), 4.29 (s, 4H), 4.35 (m, 1H), 5.15 (m, 1H), 6.90-7.10 (m, 3H), 7.30 (d, J=15.0 Hz, 1 H), 7.40 (d, J=15.0 Hz, 1H), 7.54 (d, J=15.0 Hz, 1H), 7.82 (m, 1H), 8.15 (m, 1H). MS (ESI) m/z 553 (M+H)⁺. Anal. calcd. for C₂₅H₂₃F₃N₂O₇S• 0.25 H₂O: C, 53.91; H, 4.25; N, 5.03. Found: 53.91; H, 4.35; N, 15 5.05.

Example 272

20 (Benzodioxan-6-yl)[2-trifluoromethyl-4-(*E*-((morpholin-1-yl)carbonyl)ethenyl] phenyl] sulfide

The title compound was prepared according to the procedures of Example 1. ¹H NMR (CDCl₃, 300 MHz) δ 7.76 (s, 1H), 7.62 (d, 1H, J = 15.6 Hz), 7.40 (dd, 1H, J = 1.8, 8.2 Hz), 7.04 (d, 1H, J = 2.1 Hz), 6.98-7.03 (m, 2H), 6.91 (d, 1H, J = 8.1 Hz),

6.81 (d, 1H, J = 15.3 Hz), 4.30 (m, 4H), 3.65-3.74 (br m, 8H). MS (ESI) m/z 452, 474, 925.

Example 273

5 (Benzodioxan-6-yl)[2-trifluoromethyl-4-(*E*-((4-(pyrrolidin-1-yl)piperidin-1-yl)carbonyl)ethenyl) phenyl] sulfide

The title compound was prepared according to the procedures of Example 1.

¹H NMR (CDCl₃, 300 MHz) δ 7.75 (s, 1H), 7.65 (d, 1H, J = 15.3 Hz), 7.40 (dd, 1H, J = 1.4, 8.3 Hz), 7.06 (d, 1H, J = 2.4 Hz), 6.98-7.02 (m, 2H), 6.90 (d, 1H, J = 8.1 Hz),
10 6.85 (d, 1H, J = 15.3 Hz), 4.68 (m, 1H), 4.20 (m, 4H), 3.10 (m, 1H), 3.14 (m, 1H), 2.81 (s, 4H), 2.58 (br, 1H), 2.02 (s, 4H), 1.88 (s, 4H), 1.64 (m, 1H). MS (ESI) m/z 519, 1037.

Example 274

15 (2-Isopropylphenyl)[2-nitro-4-(*E*-((3-aza-6,9-dioxaspiro[5.4]decan-1-yl)carbonyl)ethenyl) phenyl] sulfide

The title compound was prepared according to the procedures of Example 1.

¹H NMR (CDCl₃, 300 MHz) δ 8.44 (s, 1H), 7.50-7.62 (m, 4H), 7.41 (d, 1H, J = 8.0 Hz), 7.30 (m, 1H), 6.96 (br d, 1H, J = 15.6 Hz), 6.69 (d, 1H, J = 9.4 Hz), 4.00 (s, 4H),
20 3.75 (br m, 4H), 3.44 (m, 1H), 1.75 (br s, 4H), 1.18 (d, 6H, J = 7.0 Hz). MS (ESI) m/z 439, 937.

Example 275

(2-Isopropylphenyl)[2-nitro-4-(E-((2-(dimethylaminomethyl)piperidin-1-yl)carbonyl)ethenyl) phenyl] sulfide

The title compound was prepared according to the procedures of Example 1.

- 5 ¹H NMR (CDCl₃, 300 MHz) δ 8.40 (d, 1H, J = 1.8 Hz), 7.50-7.58 (m, 4H), 7.42 (d, 1H, J = 8.1 Hz), 7.30 (dd, 1H, J = 1.9, 7.0 Hz), 7.00 (d, 1H, J = 15.4 Hz), 6.68 (d, 1H, J = 8.5 Hz), 5.10 (br, 1H), 3.92 (br, 1H), 3.44 (quintet, 1H, J = 6.9 Hz), 3.20 (m, 1H), 2.26-2.50 (m, 7H), 1.62-1.85 (m, 7H), 1.48 (m, 1H), 1.18 (d, 6H, J = 7.0 Hz). MS (ESI) *m/z* 468.

10

Example 276

(2-Isopropylphenyl)[2-nitro-4-(E-((piperidin-1-ylamino)carbonyl)ethenyl) phenyl] sulfide

The title compound was prepared according to the procedures of Example 1.

- 15 ¹H NMR (CDCl₃, 300 MHz) δ 8.44 (d, 1H, J = 1.8 Hz), 7.66 (d, 1H, J = 16.2 Hz), 7.55 (d, 1H, J = 7.4 Hz), 7.47-7.51 (m, 3H), 7.30 (m, 2H), 6.72 (d, 1H, J = 8.5 Hz), 6.37 (s, 1H), 3.48 (m, 2H), 3.10 (m, 2H), 2.63 (m, 1H), 1.81-1.89 (m, 2H), 1.62-1.77 (m, 4H), 1.19 (d, 6H, J = 7.0 Hz). MS (ESI) *m/z* 426, 851.

20

Example 277

(Benzodioxan-6-yl)[2-trifluoromethyl-4-(E-((3-carboxy-4-methoxycarbonyl)piperazin-1-yl) carbonyl)ethenyl) phenyl] sulfide

The title compound was prepared by hydrolysis of the compound of Example 269 under basic conditions (aq. NaOH/EtOH). ^1H NMR (DMSO- d_6 , 300 MHz) δ 2.60-3.30 (m, 3H), 3.40-3.50 (m, 1H), 3.62 (d, $J=12.0$ Hz, 1H), 3.80 (m, 1H), 4.25-4.35 (m, 4H), 4.55 (m, 1H), 7.00 (s, 2H), 7.00-7.06 (m, 1H), 7.25 (m, 2H), 7.5 (m, 1H), 7.80 (m, 1H), 8.10 (m, 1H). MS (APCI) m/z 553 ($M+H$) $^+$. Calcd. Anal. $C_{24}H_{23}F_3N_2O_5 \cdot 1.55 H_2O$: C, 54.35; H, 4.20; N, 5.07. Found: C, 54.16; H, 4.19; N, 4.96.

Example 278

10 (2-(Dimethylaminocarbonyl)-benzodioxan-6-yl)[2-chloro-4-(E-((4-acetyl)piperazin-1-yl)carbonyl)ethenyl] phenyl] sulfide

The title compound was prepared by the procedures described in Example 85, substituting 5-iodoindole with 2-*N,N*-dimethylcarboxamide-6-bromobenzenedioxane and 3-*N,N*-dimethylcarboxamide-6-bromobenzenedioxane, giving a white solid. ^1H NMR (CDCl $_3$, 300 MHz, mixture of regioisomers) δ 1.93 (s, 3H), 2.15 (s, 6H), 3.53 (br s, 2H), 3.59-3.90 (br m, 8H), 4.86-5.01 (m, 1H), 6.74-6.81 (m, 1H), 6.80 (d, $J=15.3$ Hz, 1H), 6.93 (d, $J=8.7$ Hz, 1H), 7.02 (d, CDCl $_3$ 1.8 Hz, 1H), 7.13 (dd, $J=1.8$, 8.4 Hz, 1H), 7.16-7.25 (m, 1H), 7.54 (s, 1H), 7.58 (d, $J=15.6$ Hz, 1H). MS (ESI $^+$) ($M+Na$) $^+$ at m/z 552, 554.

Example 279

00120-5674-560

(2-Isopropylphenyl)[2-nitro-4-(E-((3-(2-(methoxymethyl)tetrazol-5-yl) piperidin-1-yl)carbonyl)ethenyl) phenyl] sulfide

The title compound was prepared by the procedures described in Example 225, substituting ethyl nipecotate with 3-*N*-methoxymethyltetrazolylpiperidine, to give a
 5 light-yellow solid. ¹H NMR (CDCl₃, 300 MHz) δ 1.19 (d, *J* = 6.9 Hz, 6H), 1.62-1.80 (br m, 2H), 1.80-2.20 (br m, 2H), 2.20-2.39 (br m, 2H), 3.12-3.38 (br m, 2H), 3.46 (s, 3H), 4.11 (septet, *J* = 6.9 Hz, 1H), 4.17-4.34 (br m, 1H), 5.79 (s, 2H), 6.70 (br s, 1H), 7.05 (d, *J* = 15.3 Hz, 1H), 7.31 (d, *J* = 7.8 Hz, 1H), 7.35-7.68 (m, 5H), 8.42 (br s, 1H). MS (ESI⁺) (M+H)⁺ at *m/z* 523.

10

Example 280

(2-Isopropylphenyl)[2-nitro-4-(E-((3-(1-(methoxymethyl)tetrazol-5-yl) piperidin-1-yl)carbonyl)ethenyl) phenyl] sulfide

The title compound was prepared by the procedures described in Example 279
 15 and separated from the same reaction mixture via SiO₂ flash column chromatography, to give a light-yellow solid. ¹H NMR (CDCl₃, 300 MHz) δ 1.19 (d, *J* = 6.9 Hz, 6H), 1.62-1.80 (br m, 2H), 1.80-2.20 (br m, 2H), 2.20-2.39 (br m, 2H), 3.12-3.38 (br m, 2H), 3.46 (s, 3H), 4.11 (septet, *J* = 6.9 Hz, 1H), 4.17-4.34 (br m, 1H), 5.79 (s, 2H), 6.70 (br s, 1H), 7.05 (d, *J* = 15.3 Hz, 1H), 7.31 (d, *J* = 7.8 Hz, 1H), 7.35-7.68 (m, 5H),
 20 8.42 (br s, 1H). MS (ESI⁺) (M+H)⁺ at *m/z* 523.

Example 281

(1-Methylindol-5-yl)[2-chloro-4-(*E*-(3-(1-pyrrolidin-2-onyl)propylamino)carbonyl)ethenyl] phenyl] sulfide

Example 281A

5 Triisopropylsilyl(1-methylindol-5-yl) sulfide

To a stirred solution of 5-bromo-*N*-methyl indole (300 mg, 1.43 mmol) in 5 mL of benzene in a sealed tube was charged with Pd(PPh₃)₄ (82 mg, 0.072 mmol), followed by KSTIPS (326 mg, 1.43 mmol). The mixture was flushed with N₂, the tube was capped, and the reaction mixture refluxed for 2 h. The reaction mixture was then
10 allowed to cool down, partitioned between Et₂O and water. The organic layer was washed with brine, dried over Na₂SO₄, concentrated in vacuo. The residue was purified on a SiO₂ flash column chromatography eluting with 5% EtOAc/hexanes to give 400 mg (88 %) of the title compound as colorless oil.

15 Example 281B

3-Chloro-4-((1-methylindol-5-yl)thio) benzaldehyde

To a stirred solution of thiolsilyl ether (1.0 g, 3.13 mmol) in 5 mL of DMF with 3-chloro-4-fluorobenzaldehyde (500 mg, 3.13 mmol) at ambient temperature was added CsF (5.7 mg, 0.38 mmol). The mixture was stirred over night before it was
20 poured in water and extracted with Et₂O (2×25 mL). The combined organic layer was washed with water and brine, dried over Na₂SO₄, concentrated in vacuo. The residue

00544795-033100
001220-56214560

was purified on a SiO₂ flash column chromatography eluting with 5-10 % EtOAc/hexanes to give 650 mg (71 %) of the title compound as white solid.

Example 281C

5 (1-Methylindol-5-yl)[2-chloro-4-(*E*-((3-(1-pyrrolidin-2-onyl)propylamino) carbonyl)ethenyl) phenyl] sulfide

The title compound was prepared by the procedures described in Example 92, substituting the benzoic acid with cinnamic acid prepared from the above-described aldehyde, and ammonium with 3-aminopropyl-1-pyrrolidin-2-one, to give a white
 10 solid. ¹H NMR (CDCl₃, 300 MHz) δ 1.74 (br m, 2H), 2.07 (br m, 2H), 2.44 (br m, 2H), 3.32 (br m, 2H), 3.40 (br m, 4H), 3.85 (s, 3H), 6.36 (d, *J* = 15.3 Hz, 1H), 7.14 (d, *J* = 3.0 Hz, 1H), 7.36 (dd, *J* = 1.5, 9.0 Hz, 1H), 7.41 (d, *J* = 9.0 Hz, 1H), 7.50 (s, 1H), 7.89 (d, *J* = 1.5 Hz, 1H). MS (ESI⁺) (M+H)⁺ at *m/z* 468, 470. Anal. Calcd for C₂₅H₂₆ClN₃O₂S · 1.37 H₂O: C, 60.95; H, 5.88; N, 8.53. Found: C, 60.97; H, 5.98;
 15 N, 8.46.

Example 282

(2-Isopropylphenyl)[2-nitro-4-(*E*-((3-(tetrazol-5-yl) piperidin-1-yl)carbonyl)ethenyl) phenyl] sulfide

20 The compound from Example 279 (75 mg, 0.14 mmol) was dissolved in 1 mL of neat TFA and left at ambient temperature for overnight. The reagent was then removed in vacuo and the residue was evaporated twice with benzene. The crude

product was purified using Gilson Preparative HPLC as described in Example 38B to

give the title compound as a light-yellow solid (50 mg, 72 %); ^1H NMR (CDCl_3 , 300

MHz) δ 1.17 (d, $J = 6.5$ Hz, 6H), 1.25-1.39 (m, 1H), 1.69-1.81 (m, 1H), 2.09 (br s,

1H), 2.14-2.30 (m, 1H), 2.57-2.71 (m, 1H), 3.35-3.66 (m, 3H), 3.90-4.03 (m, 1H),

5 4.66-4.78 (m, 1H), 6.73 (d, $J = 8.7$ Hz, 1H), 6.86 (d, $J = 15.3$ Hz, 1H), 7.32 (dd, $J =$

2.1, 6.9 Hz, 1H), 7.42 (dd, $J = 2.1$, 8.7 Hz, 1H), 7.47-7.57 (m, 3H), 7.76 (d, $J = 15.3$

Hz, 1H), 8.46 (d, $J = 2.1$ Hz, 1H). MS (ESI^+) ($\text{M}+\text{H}^+$) at m/z 479. Anal. Calcd for

$\text{C}_{24}\text{H}_{26}\text{N}_6\text{O}_3\text{S} \cdot 0.28 \text{H}_2\text{O}$: C, 59.61; H, 5.54; N, 17.38. Found: C, 59.71; H, 5.44; N,

16.99.

10

Example 283

(1-Methylindol-5-yl)[2-chloro-4-(*E*-(3-carboethoxypiperidin-1-yl)carbonyl)ethenyl]

phenyl] sulfide

The title compound was prepared by the procedures described in Example

15 281C, substituting aminopropyl pyrrolidinone with ethyl nipecotate, giving a white

solid. ^1H NMR (CDCl_3 , 300 MHz) δ 1.26 (t, $J = 7.5$ Hz, 3H), 1.65-1.96 (m, 2H), 2.00-

2.20 (m, 1H), 2.04 (s, 1H), 2.54 (br m, 1H), 3.12-3.34 (m, 1H), 3.85 (s, 3H), 3.92-4.07

(m, 1H), 4.07-4.20 (m, 1H), 4.15 (q, $J = 7.5$ Hz, 2H), 4.65-4.90 (m, 1H), 6.53 (d, $J =$

3.0 Hz, 1H), 6.57 (d, $J = 8.1$ Hz, 1H), 6.85 (d, $J = 15.3$ Hz, 1H), 7.08 (d, $J = 8.7$ Hz,

20 1H), 7.14 (d, $J = 3.0$ Hz, 1H), 7.37 (dd, $J = 1.5$, 8.7 Hz, 1H), 7.42 (d, $J = 8.7$ Hz, 1H),

7.51 (s, 1H), 7.51 (d, $J = 15.3$ Hz, 1H), 7.89 (d, $J = 1.5$ Hz, 1H). MS (ESI^+) ($\text{M}+\text{H}^+$) at

m/z 483, 485.

00543796-033400
00543796-033400

Example 284

(1-Methylindol-5-yl)[2-chloro-4-(*E*-((3-carboxypiperidin-1-yl)carbonyl)ethenyl)phenyl] sulfide

- 5 The title compound was prepared by the procedures described in Example 155, substituting the ethyl ester from Example 137 with ethyl ester from Example 283, and KOH with NaOH, to provide a white solid. ¹H NMR (CDCl₃, 300 MHz) δ 1.45-1.69 (m, 1H), 1.69-1.98 (m, 2H), 1.98-2.22 (m, 1H), 2.51-2.70 (m, 1H), 3.05-3.47 (m, 1H), 3.80-4.20 (m, 2H), 3.85 (s, 3H), 4.47-4.68 (m, 1H), 6.53 (d, *J* = 3.0 Hz, 1H), 6.57 (d, *J* = 8.1 Hz, 1H), 6.87 (d, *J* = 15.3 Hz, 1H), 7.08 (d, *J* = 8.1 Hz, 1H), 7.14 (d, *J* = 3.0 Hz, 1H), 7.37 (d, *J* = 9.0 Hz, 1H), 7.42 (d, *J* = 9.0 Hz, 1H), 7.51 (s, 1H), 7.52 (d, *J* = 15.3 Hz, 1H), 7.89 (br s, 1H). MS (ESI⁺) (M-H+H)⁺ at *m/z* 453, 455.
- 10

Example 285

- 15 (1-Methylindol-5-yl)[2-chloro-4-(*E*-((4-carboethoxypiperidin-1-yl)carbonyl)ethenyl)phenyl] sulfide

- The title compound was prepared by the procedures described in Example 281C, substituting aminopropyl pyrrolidinone with ethyl isonipecotate, giving a white solid. ¹H NMR (CDCl₃, 300 MHz) δ 1.26 (t, *J* = 7.5 Hz, 3H), 1.64-1.83 (m, 2H), 1.88-2.08 (m, 2H), 2.48-2.67 (m, 1H), 2.86-3.40 (m, 2H), 3.85 (s, 3H), 3.89-4.24 (m, 1H), 4.15 (q, *J* = 7.5 Hz, 2H), 4.24-4.65 (m, 1H), 6.53 (d, *J* = 3.0 Hz, 1H), 6.58 (d, *J* = 8.1 Hz, 1H), 6.81 (d, *J* = 15.3 Hz, 1H), 7.07 (d, *J* = 8.1 Hz, 1H), 7.14 (d, *J* = 3.0 Hz, 1H),
- 20

7.37 (dd, $J = 1.5, 9.0$ Hz, 1H), 7.50 (d, $J = 9.0$ Hz, 1H), 7.50 (d, $J = 15.3$ Hz, 1H), 7.88 (d, $J = 1.5$ Hz, 1H). MS (ESI⁺) (M+H)⁺ at m/z 483, 485.

Example 286

5 (1-Methylindol-5-yl)[2-chloro-4-(*E*-((3-carboxypiperidin-1-yl)carbonyl)ethenyl)phenyl] sulfide

The title compound was prepared by the procedures described in Example 155, substituting the ethyl ester from Example 137 with ethyl ester from Example 285, and KOH with NaOH, giving a white solid. ¹H NMR (CDCl₃, 300 MHz) δ 1.60-1.90 (m, 2H), 1.90-2.10 (m, 2H), 2.57-2.72 (m, 1H), 2.80-3.40 (m, 2H), 3.85 (s, 3H), 3.91-4.20 (m, 1H), 4.30-4.68 (m, 1H), 6.53 (d, $J = 3.0$ Hz, 1H), 6.57 (d, $J = 8.1$ Hz, 1H), 6.80 (d, $J = 15.3$ Hz, 1H), 7.07 (d, $J = 8.1$ Hz, 1H), 7.15 (d, $J = 3.0$ Hz, 1H), 7.37 (dd, $J = 1.5, 9.0$ Hz, 1H), 7.51 (d, $J = 9.0$ Hz, 1H), 7.51 (s, 1H), 7.51 (d, $J = 15.3$ Hz, 1H), 7.89 (br s, 1H). MS (ESI⁺) (M+H)⁺ at m/z 455, 457. Anal. Calcd for C₂₄H₂₃ClN₂O₃S · 0.42 H₂O: C, 62.32; H, 5.20; N, 6.06. Found: C, 62.35; H, 5.30; N, 5.87.

Example 287

(2-Isopropylphenyl)[2-nitro-4-(*E*-((2-(1-methylpyrrolidin-2-yl)ethylamino)carbonyl)ethenyl)phenyl] sulfide

20 The title compound was prepared according to the procedures of Example 1. ¹H NMR (CDCl₃, 300 MHz) δ 8.44 (d, 1H, $J = 1.8$ Hz), 7.56 (d, 1H, $J = 3.7$ Hz), 7.50-7.58 (m, 3H), 7.43 (DD, 1H, $J = 1.84, 8.4$ Hz), 7.30 (dd, 1H, $J = 2.2, 6.8$ Hz),

6.78 (d, 1H, J = 8.5 Hz), 6.52 (d, 1H, J = 15.8 Hz), 3.63 (m, 2H), 3.42 (m, 3H), 3.00 (m, 1H), 3.78 (m, 1H), 2.59 (s, 3H), 2.05 (m, 1H), 2.00 (m, 5H), 1.18 (d, 6H, J = 7.0 Hz). MS (ESI) m/z 454, 490.

5

Example 288

(2-Isopropylphenyl)[2-nitro-4-(E-((4-(pyrrolidin-1-yl)piperidin-1-yl)carbonyl)ethenyl)phenyl] sulfide

The title compound was prepared according to the procedures of Example 1.

¹H NMR (CDCl₃, 300 MHz) δ 8.43 (d, 1H, J = 1.8 Hz), 7.57 (d, 1H, J = 8.5 Hz), 7.51-7.55 (m, 3H), 7.41 (dd, 1H, J = 1.84, 8.8 Hz), 7.31 (dd, 1H, J = 2.4, 7.5 Hz), 6.92 (d, 1H, J = 15.4 Hz), 6.70 (d, 1H, J = 8.5 Hz), 4.70 (m, 1H), 4.10 (m, 1H), 3.44 (pent, 1H, J = 6.8 Hz), 3.16 (m, 1H), 2.80 (br, 4H), 2.55 (br, 1H), 2.03 (m, 4H), 1.90 (m, 4H), 1.65 (m, 1H), 1.18 (d, 6H, J = 7.0 Hz). MS (ESI) m/z 480, 959.

15

Example 289

(2-Isopropylphenyl)[2-nitro-4-(E-((4-sulfopiperidin-1-yl)carbonyl)ethenyl)phenyl] sulfide

The title compound was prepared according to the procedures of Example 1.

¹H NMR (DMSO-d₆, 300 MHz) δ 8.63 (d, 1H, J = 1.8 Hz), 7.92 (dd, 1H, J = 1.8, 8.8 Hz), 7.60 (m, 3H), 7.47 (d, 1H, J = 14.2 Hz), 7.42 (d, 1H, J = 14.2 Hz), 6.62 (d, 1H, J = 8.5 Hz), 4.45 (m, 2H), 4.38 (m, 2H), 3.34 (m, 1H), 3.00 (m, 2H), 2.70 (m, 1H), 2.60 (m, 2H), 1.14 (d, 6H, J = 6.9 Hz). MS (ESI) m/z 491, 981.

Example 290

(2-Isopropylphenyl)[2-nitro-4-(*E*-((3-hydroxypiperidin-1-yl)carbonyl)ethenyl) phenyl]
sulfide

- 5 The title compound was prepared according to the procedures of Example 1.
- ¹H NMR (CDCl₃, 300 MHz) δ 8.43 (s, 1H), 7.50-7.62 (m, 4H), 7.41 (d, 1H, J = 8.1 Hz), 6.97 (m, 1H), 6.69 (d, 1H, J = 8.1 Hz), 3.85 (m, 2H), 3.65 (m, 1H), 3.50 (m, 3H), 1.93 (m, 2H), 1.65 (m, 2H), 1.18 (d, 6H, J = 6.6 Hz). MS (ESI) *m/z* 427, 449, 853, 875.

10

Example 291

(Benzodioxan-6-yl)[2-trifluoromethyl-4-(*E*-((3-
 ((ethanesulfonylamino)carbonyl)piperidin-1-yl) carbonyl)ethenyl) phenyl] sulfide

The title compound was prepared by the procedures described in Example 227.

- 15 The product was purified by reversed-phase HPLC. ¹H NMR (CDCl₃, 300 MHz) δ 1.34 (t, J= 7.0 Hz, 2H), 1.44 (t, J=7.0 Hz, 3H), 1.95 (br, 1/2H), 2.20 (br, 1/2H), 2.68 (br, 1H), 3.14 (q, J=7.0 Hz, 2H), 3.45 (m, 1H), 3.65 (m, 1H), 3.93 (m, 1H), 4.30 (m, 4H), 4.50-4.60 (br, 2H), 6.92 (d, J=8.0 Hz, 1H), 6.98-7.04 (m, 3H), 7.06 (m, 1H), 7.40 (d, J=8.0 Hz, 1H), 7.65 (m, 1H), 7.75 (s, 1H). MS (APCI) *m/z* 585 (M+H)⁺.

20

Example 292

(Benzodioxan-6-yl)[2-trifluoromethyl-4-(*E*-((3-(*p*-toluenesulfonylamino)carbonyl)piperidin-1-yl) carbonyl)ethenyl) phenyl] sulfide

The title compound was prepared by the same procedure described in Example

229. ¹H NMR (CDCl₃, 300 MHz) δ 1.25 (m, 2H), 1.55 (m, 1H), 1.70-2.25 (br, 1H),
 5 2.41 (d, J=13.0 Hz, 3H), 2.55 (br, 1H), 3.50-3.80 (br, 2H), 4.20-4.35 (m, 4H), 4.68-
 4.75 (m, 2H), 6.90 (d, J=8.0 Hz, 1H), 7.00-7.10 (m, 2H), 7.30 (d, J=8.0 Hz, 1H), 7.81
 (d, J=8.0 Hz, 1H), 7.91 (m, 1H). MS (CI/NH₃) m/z 647 (M+H)⁺. Anal. calcd. for
 C₃₁H₂₉F₃N₂O₆S₂ • 0.5 H₂O: C, 56.78; H, 4.61; N, 4.27. Found: C, 56.86; H, 4.69; N,
 4.35.

10

Example 293

((Benzodioxan-6-yl)[2-trifluoromethyl-4-(*E*-((4-(ethanesulfonylamino)carbonyl)piperidin-1-yl) carbonyl)ethenyl) phenyl] sulfide

- The title compound was prepared by the procedures described in Example 227,
 15 giving a white foam. ¹H NMR (CDCl₃, 300 MHz) δ 1.35-1.40 (m, 2H), 1.44 (t, J=7.0
 Hz, 3H), 1.76 (m, 1H), 2.0 (m, 1H), 2.50-3.20 (br, 1H), 3.15 (q, J=7.0 Hz, 2H), 3.40-
 3.55 (m, 2H), 4.25-4.32 (m, 4H), 4.52 (br, 2H), 6.90 (d, J=8.0 Hz, 1H), 6.98-7.05 (dd,
 J=2.0, 8.0 Hz, 2H), 7.06 (d, J=2.0 Hz, 1H), 7.40 (m, 1H), 7.60 (m, 1H), 7.75 (s, 1H),
 8.22 (br, 1H). MS (APCI) m/z 585 (M+H)⁺. Anal. calcd. for C₂₆H₂₇F₃N₂O₆S₂ • 0.8
 20 H₂O: C, 52.13; H, 4.81; N, 4.68. Found: C, 52.14; H, 4.80; N, 4.66.

Example 294

(Benzodioxan-6-yl)[2-trifluoromethyl-4-(*E*-(2-tetrazol-5-yl)morpholin-1-yl)carbonyl)ethenyl] phenyl] sulfide

The corresponding nitrile (160 mg, 0.336 mmol, prepared via the procedures of Example 1), sodium azide (56.6 mg, 0.872 mmol), n-Bu₃SnCl and THF were mixed

5 in a reaction tube, flushed with nitrogen and heated to reflux overnight. The mixture was then cooled to ambient temperature, and 1N HCl soln. was added. The mixture was extracted with ethyl acetate three times and the combined organics were dried over MgSO₄. The mixture was filtered through a short silica gel plug to give 96 mg (56% yield) of the desired material. ¹H NMR (DMSO-d₆, 500 MHz, 100 °C) δ 7.99

10 (d, 1H, J = 1.7 Hz), 7.79 (dd, 1H, J = 2.0, 8.6 Hz), 7.50 (d, 1H, J = 15.3 Hz), 7.24 (d, 1H, J = 15.6 Hz), 7.14 (d, 1H, J = 8.2 Hz), 6.96 (m, 1H), 6.94 (d, 1H, J = 2.1 Hz), 6.92 (m, 1H), 4.60 (dd, 1H, J = 3.0, 9.8 Hz), 4.50 (br d, 1H, J = 12.2 Hz), 4.26 (m, 5H), 4.17 (m, 1H), 4.00 (dt, 1H, J = 3.2, 11.6 Hz), 3.72 (td, 1H, J = 3.0, 11.0 Hz), 3.43 (br m, 1H), 3.29 (br m, 1H). MS (ESI) *m/z* -518. Anal. Calcd for C₂₃H₂₀F₃N₅O₄S ·

15 1.83 HOAc: C, 50.88; H, 4.38; N, 11.13. Found: C, 50.61; H, 4.46; N, 11.4.

Example 295

(2-Isopropylphenyl)[2-nitro-4-(*E*-(2-butyl, 5-(tetrazol-5-yl)morpholin-1-yl)carbonyl)ethenyl] phenyl] sulfide

Example 295A

2-Butyl-5-cyanomorpholine

The title compound was prepared by the procedures described in Example 260A, substituting ethanolamine with 2-aminoethanol.

Example 295B

5 (2-Isopropylphenyl)[2-nitro-4-(*E*-((2-butyl-5-cyanomorpholin-1-yl)carbonyl)ethenyl)phenyl] sulfide

The title compound was prepared by the procedures described in Example 260B, substituting the morpholine from Example 260A with the compound of Example 295A.

10

Example 295C

(2-Isopropylphenyl)[2-nitro-4-(*E*-(2-butyl-5-(tetrazol-5-yl)morpholin-1-yl)carbonyl)ethenyl) phenyl] sulfide

The title compound was prepared by the procedures described in Example 262, substituting the nitrile compound from Example 260 with the compound of Example 295B, giving a light-yellow solid. ¹H NMR (CDCl₃, 300 MHz, 3:2 mixture of diastereomers) δ 0.89 (t, *J* = 7.5 Hz, 1H), 1.01 (br m, 1H), 1.19 (d, *J* = 6.5 Hz, 6H), 1.23-1.43 (m, 4H), 1.68-1.84 (m, 1H), 3.10-3.61 (m, 2H), 3.83-4.17 (m, 2H), 4.40-5.26 (m, 2H), 6.67-6.77 (m, 1H), [6.91 (d), 7.02 (d), *J* = 15.3 Hz, 1H in total], 7.25-7.37 (m, 2H), 7.44-7.60 (m, 3H), [7.67 (d), 7.79 (d), *J* = 15.3 Hz, 1H in total], 8.43-8.50 (m, 1H). MS (ESI⁺) (M-H)⁺ at *m/z* 535.

20

Example 296

(2-(and 3-)(Hydroxymethyl)-benzodioxan-6-yl)[2-nitro-4-(E-((4-acetyl)piperazin-1-yl)carbonyl)ethenyl) phenyl] sulfide

5

Example 296A

Triisopropylsilyl (2-(and 3-)hydroxymethylbenzodioxan-6-yl) sulfide

The title compound was prepared by the procedures described in Example 281A, substituting 5-bromo-*N*-methyl indole with a mixture of 6-bromo-2-hydroxymethylbenzenedioxane and 6-bromo-3-hydroxymethylbenzenedioxane.

10

Example 296B

(2-(and 3-)(Hydroxymethyl)-benzodioxan-6-yl)[2-nitro-4-(E-((4-acetyl)piperazin-1-yl)carbonyl)ethenyl) phenyl] sulfide

The title compound was prepared by the procedures described in Example

- 15 281B, substituting 3-chloro-4-fluorobenzaldehyde with 4-chloro-3-nitrocinnamide, giving a light yellow solid. ¹H NMR (CDCl₃, 300 MHz, 3:2 mixture of diastereomers) δ [2.11 (s), 2.15 (s), 3H in total], 3.48-3.83 (m, 8H), 3.83-4.04 (m, 2H), 4.20 (dd, *J* = 8.4, 11.4 Hz, 1H), 4.26-4.44 (m, 2H), 6.89 (d, *J* = 5.7 Hz, 1H), 6.92 (s, 1H), 6.97-7.11 (m, 1H), 7.04 (d, *J* = 15.0 Hz, 1H), 7.14 (d, *J* = 2.1 Hz, 1H), 7.46 (br d, *J* = 9.0 Hz, 1H), 7.65 (d, *J* = 15.0 Hz, 1H), 8.41 (d, *J* = 2.1 Hz, 1H). MS (ESI⁺) (M+H)⁺ at *m/z* 500.
- 20

00544756-033400
00544756-033400

Example 297

(2-(and 3-)(Hydroxymethyl)-benzodioxan-6-yl)[2-nitro-4-(E-((3-(pyrrolidin-2-on-1-yl)prop-1-ylamino)carbonyl) ethenyl)phenyl]sulfide

- The title compound was prepared by the procedures described in Example 296B, substituting the acetylpipezazine 4-chloro-3-nitrocinnamide with 3-aminopropyl-1-pyrrolidin-3-one 4-chloro-3-nitrocinnamide, giving a light-yellow solid. ¹H NMR (CDCl₃, 300 MHz, 3:2 mixture of diastereomers) δ 1.75 (br m, 2H), 2.08 (p, *J* = 7.5 Hz, 2H), 2.45 (t, *J* = 7.5 Hz, 2H), 3.27-3.48 (m, 6H), 3.82-4.03 (m, 2H), 4.13-4.44 (m, 3H), 6.49 (d, *J* = 15.0 Hz, 1H), 6.88 (d, *J* = 8.4 Hz, 1H), [6.99 (d), 7.01 (d), *J* = 8.4 Hz, 1H in total], [7.06 (dd), 7.08 (dd), *J* = 1.5, 2.4 Hz, 1H in total], [7.13 (d), 7.14 (d), *J* = 2.4 Hz, 1H in total], 7.17 (br s, 1H), 7.46 (d, *J* = 8.4 Hz, 1H), 7.54 (d, *J* = 15.0 Hz, 1H), 8.36 (d, *J* = 1.5 Hz, 1H). MS (ESI⁺) (M+H)⁺ at *m/z* 514.

Example 298

- (2-(and 3-)(Hydroxymethyl)-benzodioxan-6-yl)[2-trifluoromethyl-4-(E-((3-(pyrrolidin-2-on-1-yl)prop-1-ylamino)carbonyl) ethenyl)phenyl]sulfide

- The title compound was prepared by the procedures described in Example 281, substituting 6-thiolsilyl indole with the thiolsilyl ether described in Example 296A, and 3-chloro-4-fluorobenzaldehyde with 4-fluoro-3-trifluoromethylbenzaldehyde, producing a white solid. ¹H NMR (CDCl₃, 300 MHz, 3:2 mixture of diastereomers) δ 1.75 (br m, 2H), 2.09 (br m, 2H), 2.45 (br m, 2H), 3.25-3.60 (m, 6H), 3.80-4.43 (m, 5H), 6.46 (d, *J* = 15.3 Hz, 1H), [6.92 (d), 6.95 (d), *J* = 6.8 Hz, 1H in total], [7.03 (d),

7.04 (d), $J = 8.1$ Hz, 1H in total], 7.06-7.10 (m, 1H), 7.13 (br s, 1H), 7.42 (d, $J = 8.1$ Hz, 1H), 7.54 (d, $J = 15.3$ Hz, 1H), 7.77 (s, 1H). MS (ESI⁺) (M+H)⁺ at m/z 537.

Example 299

- 5 3-(Hydroxymethyl)-benzodioxan-6-yl)[2-nitro-4-(E-((3-(pyrrolidin-2-on-1-yl)prop-1-ylamino)carbonyl) ethenyl)phenyl]sulfide

Example 299A

3-(Hydroxymethyl)-6-bromo-benzodioxane

- 10 To a stirred solution of 5-bromosalicylaldehyde (5.0 g, 24.9 mmol), and epichlorohydrin (5.6 mL, 72.1 mmol) in 20 mL of DMF at 80 °C was added K₂CO₃ slowly in portions. The resulting mixture was then heated at 90 °C for 3 h. Reaction was then stopped, water was added, extracted with diethyl ether. The organic extracts were washed with water, brine, dried over Na₂SO₄, concentrated in vacuo. The residue
- 15 was purified on a SiO₂ flash column chromatography eluting with 15-30 % EtOAc/hexanes to give 2.82 g (44 %) of the title compound as colorless oil.

- To a stirred solution of the aldehyde (2.82 g, 11 mmol) in 35 mL of CHCl₃ was added mCPBA (2.27 g, 13 mmol). The mixture was stirred at ambient temperature for 30 min and then heated at 50 °C for 2 h. The reaction was then
- 20 quenched with aq. Na₂S₂O₃, extracted with Et₂O (2×50 mL). The combined organic layer was washed with aq. NaHCO₃, brine, dried over Na₂SO₄, concentrated in vacuo

to give 2.92 g of crude product which was proceeded to the next step without purification.

To a stirred solution of the above-described crude formate (2.92 g) in 5 mL of THF was added 3N aq. NaOH (3.9 mL, 11.7 mmol). The reaction mixture was then heated at 70 °C for 4 h. The reaction mixture was then partitioned between EtOAc and water. The organic layer was then washed with brine, dried over Na₂SO₄, concentrated in vacuo to give 2.50 g (93% over two steps) of the title compound.

Example 299B

10 Triisopropyl (3-(hydroxymethyl)-benzodioxan-6-yl) sulfide

The title compound was prepared by the procedures described in Example 281A, substituting 5-bromo-*N*-methyl indole with the bromide from Example 299A.

Example 299C

15 (3-Hydroxymethyl)-benzodioxan-6-yl)[2-nitro-4-(E-((3-(pyrrolidin-2-on-1-yl)prop-1-ylamino)carbonyl) ethenyl)phenyl]sulfide

The title compound was prepared by the procedures described in Example 297, substituting the mixture of thioisilyl ethers from Example 296A with the compound of Example 299B, giving a white solid. ¹H NMR (CDCl₃, 300 MHz) δ 1.74 (br m, 2H), 2.08 (t, *J* = 7.5 Hz, 2H), 2.44 (t, *J* = 7.5 Hz, 2H), 3.25-3.53 (m, 6H), 3.88 (dd, *J* = 4.8, 16.8 Hz, 1H), 3.97 (dd, *J* = 4.8, 16.8 Hz, 1H), 4.21 (dd, *J* = 3.1, 12.9 Hz, 1H), 4.26-4.36 (m, 1H), 4.40 (dd, *J* = 2.4, 12.9 Hz, 1H), 6.49 (d, *J* = 15.3 Hz, 1H), 6.88 (d, *J* =

8.7 Hz, 1H), 7.00 (d, $J = 8.7$ Hz, 1H), 7.07 (dd, $J = 2.4, 8.7$ Hz, 1H), 7.14 (d, $J = 2.4$ Hz, 1H), 7.20 (br s, 1H), 7.46 (dd, $J = 0.9, 8.7$ Hz, 1H), 7.54 (d, $J = 15.3$ Hz, 1H), 8.36 (s, 1H). MS (ESI⁺) (M+H)⁺ at m/z 514. Anal. Calcd for C₂₅H₂₇N₃O₇S · 0.82 H₂O: C, 56.83; H, 5.46; N, 7.95. Found: C, 56.84; H, 5.18; N, 7.74.

5

Example 300

(Benzodioxan-6-yl)[2-chloro-4-(*E*-((3-carboxypiperidin-1-yl)carbonyl)ethenyl)phenyl] sulfide

The title compound was prepared by the procedures described in Example 263, substituting 4-fluoro-3-trifluoromethylbenzaldehyde with 3-chloro-4-fluorobenzaldehyde, giving a white solid. ¹H NMR (CDCl₃, 300 MHz) δ 1.64-1.88 (br m, 2H), 1.95-2.09 (br m, 2H), 2.57-2.73 (m, 1H), 2.90-3.17 (m, 1H), 3.17-3.50 (m, 1H), 3.90-4.19 (m, 1H), 4.25-4.36 (m, 4H), 4.39-4.66 (m, 1H), 6.75 (d, $J = 8.4$ Hz, 1H), 6.84 (d, $J = 15.3$ Hz, 1H), 6.93 (d, $J = 8.7$ Hz, 1H), 7.03 (dd, $J = 2.4, 8.7$ Hz, 1H), 7.08 (d, $J = 2.4$ Hz, 1H), 7.18 (d, $J = 8.4$ Hz, 1H), 7.51 (s, 1H), 7.54 (d, $J = 15.3$ Hz, 1H). MS (ESI⁺) (M+H)⁺ at m/z 460, 462.

20

Example 301

(2-(and 3-(Aminomethyl)-benzodioxan-6-yl)[2-trifluoromethyl-4-(*E*-((3-(pyrrolidin-2-on-1-yl)prop-1-ylamino)carbonyl) ethenyl)phenyl]sulfide

Example 301A

00100-56274560

(2-(and 3-)(Mesyloxymethyl)-benzodioxan-6-yl)[2-trifluoromethyl-4-(E-((3-(pyrrolidin-2-on-1-yl)prop-1-ylamino)carbonyl) ethenyl)phenyl]sulfide

To a stirred solution of alcohol from Example 298 (200 mg, 0.37 mmol)) in 2 mL of methylene chloride with Et₃N (104 mL, 0.74 mmol)) was added

- 5 methanesulfonyl chloride (35 mL, 0.56 mmol) dropwise. The mixture was then stirred at ambient temperature for one hour. The reaction mixture was then poured into 3N HCl, extracted with EtOAc (2×10 mL). The combined organic layer was washed with aq. NaHCO₃, brine, dried over Na₂SO₄, concentrated in vacuo to give 275 mg of crude product which was proceeded to the next step without purification.

10

Example 301B

(2-(and 3-)(Azidomethyl)-benzodioxan-6-yl)[2-trifluoromethyl-4-(E-((3-(pyrrolidin-2-on-1-yl)prop-1-ylamino)carbonyl) ethenyl)phenyl]sulfide

To a stirred solution suspension of NaN₃ (44 mg, 0.68 mmol) in 1 mL of

- 15 DMSO was added mesylate (275 mg) in 0.5 mL of DMSO solution. The reaction mixture was then heated at 70 °C for 2 h, then cooled down to room temperature, water was added, extracted with EtOAc (2×10 mL). The combined organic layer was washed with water, brine, dried over Na₂SO₄, concentrated in vacuo. The residue was purified on a SiO₂ flash column chromatography eluting with 5-10% MeOH/EtOAc to
- 20 give 35 (17%, two steps) mg of the title compound as light brown oil.

Example 301C

(2-(and 3-)(Aminomethyl)-benzodioxan-6-yl)[2-trifluoromethyl-4-(E-((3-(pyrrolidin-2-on-1-yl)prop-1-ylamino)carbonyl)ethenyl)phenyl]sulfide

To a stirred solution of azide (230 mg, 0.41 mmol) in 1 mL of THF was added

- 5 PPh₃ (118 mg, 0.45 mmol), followed by one drop of water. The mixture was then stirred at room temperature for one hour. The volatile solvent was then removed in vacuo and the crude product was purified using Gilson Preparative HPLC as described in Example 38B to give 25 mg (11%) of the title compound. Light brown oil; ¹H NMR (CDCl₃, 300 MHz, 3:2 mixture of diastereomers) δ 1.74 (br m, 2H), 1.96-2.16 (m, 2H), 2.35-2.50 (m, 2H), 3.23-3.47 (m, 6H), 3.92-4.63 (m, 5H), 6.41-6.55 (m, 1H), 10 6.83-7.10 (m, 3H), 7.36-7.58 (m, 3H), 7.67-7.67 (m, 2H). MS (ESI⁺) (M+H)⁺ at *m/z* 536. Anal. Calcd for C₂₆H₂₈F₃N₃O₄S · 0 H₂O: C, 58.31; H, 5.27; N, 7.85. Found: C, 58.34; H, 5.48; N, 7.78.

15

Example 302

(2-Isopropylphenyl)[2-nitro-4-(E-((3-(methylaminocarbonyl)morpholin-1-yl)carbonyl)ethenyl) phenyl] sulfide

Prepared according to the procedures of Example 71, giving a yellow solid.

- ¹H NMR (DMSO-d₆, 300MHz) δ 1.14 (d, J = 6.6 Hz, 6H); 2.61 (d, J = 4.8 Hz, 3H); 10 3.14-4.62 (br m, 7H); 3.30-3.40 (m, 1H); 6.63 (d, J = 8.8 Hz, 1H); 7.32-7.62 (m, 6H); 7.80-7.97 (m, 2H); 8.66(d, J = 1.5 Hz, 1H). MS (APCI) (M+H)⁺ at *m/z* 470.

Anal calcd for $C_{24}H_{27}N_3O_3 \cdot 0.8H_2O$: C, 59.58; H, 5.96; N, 8.68. Found: C, 59.57; H, 5.94; N, 8.72.

Example 303

5 (2-Isopropylphenyl)[2-nitro-4-(*E*-(3-(hydroxymethyl)morpholin-1-yl)carbonyl)ethenyl phenyl] sulfide

Prepared according to the procedures of Example 71, giving a yellow solid.

¹H NMR (DMSO-*d*₆, 300MHz) δ 1.14 (d, J = 6.8 Hz, 6H); 2.70-3.51 (br m, 5H); 3.30-3.40 (m, 1H); 3.83-3.93 (m, 1H); 4.03-4.47 (br m, 2H); 4.74-4.82 (m, 1H); 6.64 (d, J = 8.5 Hz, 1H); 7.30-7.62 (m, 6H); 7.86-7.94 (m, 1H); 8.59-8.65 (m, 1H). MS (APCI) (M+H)⁺ at m/z 443. Anal calcd for $C_{23}H_{26}N_2S_1O_3$: C, 62.43; H, 5.92; N, 6.33. Found: C, 62.12; H, 6.20; N, 6.06.

Example 304

15 (2-Isopropylphenyl)[2-nitro-4-(*E*-(3-(acetoxymethyl)morpholin-1-yl)carbonyl)ethenyl phenyl] sulfide

Prepared according to the procedures of Example 71, giving a yellow solid.

¹H NMR (DMSO-*d*₆, 300MHz) δ 1.14 (d, J = 7.1 Hz, 6H); 2.04 (s, 3H); 3.30-3.40 (m, 1H); 2.58-4.41 (br m, 9H); 6.64 (d, J = 8.5 Hz, 1H); 7.30-7.62 (m, 6H); 7.90 (dd, J = 8.5, 1.8 Hz, 1H); 8.59-8.65 (m, 1H). MS (APCI) (M+H)⁺ at m/z 485. Anal calcd for $C_{25}H_{28}N_2S_1O_6$: C, 61.97; H, 5.82; N, 5.78. Found: C, 61.85; H, 5.84; N, 5.68.

Example 305

(2-Isopropylphenyl)[2-nitro-4-(E-((3-(aminomethyl)morpholin-1-yl)carbonyl)ethenyl) phenyl] sulfide

Prepared according to the procedures of Example 71, giving a yellow solid.

- 5 ¹H NMR (DMSO-d₆, 300MHz) δ 1.14 (d, J = 7.0 Hz, 6H); 2.61 (d, J = 5.5 Hz, 2H); 2.49-3.60 (br m, 5H); 3.82-3.93 (m, 1H); 4.13-4.45 (m, 2H); 6.64 (d, J = 8.5 Hz, 1H); 7.32-7.62 (m, 6H); 7.88-7.95 (m, 1H); 8.59-8.67(m, 1H). MS (APCI) (M+H)⁺ at m/z 442. Anal calcd for C₂₃H₂₇N₃S₁O₄·0.4H₂O: C, 61.55; H, 6.25; N, 9.36. Found: C, 61.60; H, 6.25; N, 9.00.

10

Example 306

(2-Isopropylphenyl)[2-nitro-4-(E-((3-(acetamidomethyl)morpholin-1-yl)carbonyl)ethenyl) phenyl] sulfide

Prepared according to the procedures of Example 71, giving a yellow solid.

- 15 ¹H NMR (DMSO-d₆, 300MHz) δ 1.14 (d, J = 6.8 Hz, 6H); 1.82 (s, 3H); 2.70-3.50 (br m, 7H); 3.85-3.94 (m, 1H); 4.13-4.40 (m, 2H); 6.64 (d, J = 8.5 Hz, 1H); 7.32-7.62 (m, 6H); 7.88-8.06 (m, 1H); 8.59-8.67(m, 1H). MS (APCI) (M+H)⁺ at m/z 484. Anal calcd for C₂₅H₂₉N₃S₁O₅·0.27H₂O: C, 61.47; H, 6.10; N, 8.60. Found: C, 61.50; H, 6.34; N, 8.53.

20

Example 307

(Benzodioxan-6-yl)[2-chloro-4-(E-((3-(pyrrolidin-2-on-1-yl)prop-1-ylamino)carbonyl)ethenyl)phenyl]sulfide

The title compound was prepared by the procedures described in Example 300 substituting ethyl isonipecotate with N-(3'-aminopropyl)-2-pyrrolidinone. ¹H NMR (CDCl₃, 300 MHz) δ 1.75 (br s, 2H), 2.02-2.34 (m, 2H), 2.40-2.50 (m, 2H), 3.30-3.50 (m, 6H), 4.28-4.33 (m, 4H), 6.40 (br, 1H), 6.75 (d, J=8.0 Hz, 1H), 6.93 (d, J=8.5 Hz, 1H), 7.02 (dd, J=2.0, 8.0 Hz, 1H), 7.08 (d, J=2.0 Hz, 1H), 7.18 (d, J=8.5 Hz, 1H), 7.45 (m, 1H), 7.50 (s, 1H). MS (ESI) m/z 473 (M+H)⁺. Anal. calcd. for C₂₄H₂₅ClN₂O₄S•0.5 H₂O: C, 59.81; H, 5.44; N, 5.81. Found: C, 59.76; H, 5.80; N, 5.43.

Example 308

(Benzodioxan-6-yl)[2-chloro-4-(E-((3-carboethoxypiperidin-1-yl) carbonyl)ethenyl)phenyl] sulfide

The title compound was prepared by the procedures described in Example 300 substituting ethyl isonipecotate with ethyl nipecotate. ¹H NMR (CDCl₃, 300 MHz) δ 1.25 (t, J=7.0 Hz, 3H), 1.60-1.90 (br, 2H), 2.10 (br, 1H), 2.52 (br, 1H), 3.00-3.50 (br, 2H), 3.80 (br, 1H), 4.10-4.20 (m, 4H), 4.28-4.35 (m, 4H), 6.74 (d, J=8.0 Hz, 1H), 6.92 (d, J=8.0 Hz, 1H), 7.02 (dd, J=2.0, 8.0 Hz, 1H), 7.08 (d, J=2.0 Hz, 1H), 7.18 (m, 1H), 7.50-7.03 (m, 3H). MS (ESI) m/z 488 (M+H)⁺. Anal. calcd. for C₂₅H₂₆ClNO₅SNa•0.5 H₂O: C, 60.42; H, 5.48; N, 2.82. Found: C, 60.61; H, 5.51; N, 2.42.

Example 309

(Benzodioxan-6-yl)[2-chloro-4-(*E*-(2-carboethoxypiperidin-1-yl) carbonyl)ethenyl
phenyl] sulfide

The title compound was prepared by the same procedure described in Example

- 5 300 substituting ethyl isonipecotate with ethyl pipecolate. ¹H NMR (CDCl₃, 300 MHz) δ 1.30 (t, J=7.0 Hz, 3H), 1.30-1.50 (br, 3H), 1.55-1.85 (br, 3H), 2.30 (m, 1H), 4.00 (m, 1H), 4.20 (m, 2H), 4.30 (m, 4H), 5.44 (br, 1H), 6.85 (d, J=8.0 Hz, 1H), 6.90 (d, J=8.0 Hz, 1H), 7.00 (dd, J=2.0, 8.0 Hz, 1H), 7.07 (d, J=2.0 Hz, 1H), 7.10-7.20 (m, 2H), 7.22 (m, 1H), 7.50 (s, 1H). MS (ESI) m/z 488 (M+H)⁺. Anal. calcd. for
- 10 C₂₅H₂₆ClNO₅S: C, 61.53; H, 5.37; N, 2.87. Found: C, 61.86; H, 5.63; N, 2.56.

Example 310

(2-Methoxyphenyl)-[2,3-dichloro-4(*E*-[(morpholin-1-yl)carbonyl]ethenyl)phenyl]
sulfide

15

Example 310A2,3-Dichloro-4-trifluoromethanesulfonyloxy-benzaldehyde

- 2,3-Dichloro-4-hydroxy-benzaldehyde (9.10 g, J. Med. Chem. 19 (4), 534, 1994) was dissolved in 45 ml. pyridine at room temperature. The solution was placed
- 20 in an ice bath and immediately, 15.63 g. of trifluoromethanesulfonic anhydride was added slowly. [Note: If the pyridine solution is cooled to zero before addition of triflic anhydride the aldehyde crystallizes out and the mixture cannot be stirred.]

- After the addition is complete the dark mixture was stirred for 1 hour at room temperature. It was then poured into a stirred mixture of ice water, 100 ml. of concentrated HCl and ether. [Note: Not everything is soluble in this mixture] The ether layer was separated, dried over sodium sulfate, and the solvent removed. Warm
- 5 heptane was added to this residue, and any insoluble material was filtered. The solution was concentrated to give 8.74 g. (57% yield) of product as an orange oil which solidified in the refrigerator.

Example 310B

10 2,3-Dichloro-4-(2-methoxyphenylthio)-benzaldehyde

- 2,3-Dichloro-4-trifluoromethanesulfonyloxy-benzaldehyde (2.50 g.) was dissolved in 6 ml. acetonitrile. 2-Methoxybenzenethiol (2.55 g. of 70% pure material, 50% excess) was added. With cooling 2.50 g. diisopropylethylamine was added slowly. The solution was removed from the ice bath, whereon a solid formed. The
- 15 solution was warmed in a 50C waterbath for 5 minutes. More acetonitrile (5 ml.) was added and the mixture was cooled in ice, and then filtered to get 2.047 g. of product, m.p. 137-139C.

Example 310C

20 2,3-Dichloro-4-(2-methoxyphenylthio)-cinnamic acid

- A mixture of 2,3-dichloro-4-(2-methoxyphenylthio)-benzaldehyde (2.03 g.), 1.44 g. malonic acid, 5 ml. pyridine, and 0.100 g piperidine was heated to 115 degrees

for 1.5 hours. The mixture was cooled, and ice and HCl were added. The resulting solid was filtered, washed with water and dissolved in tetrahydrofuran. This solution was dried over sodium sulfate, the solvent removed and ether added to give 1.733 g of product, m.p. 187-188C.

5

Example 310D

(2-Methoxyphenyl)-[2,3-dichloro-4(E-[(morpholin-1-yl)carbonyl]ethenyl)phenyl]

sulfide

The title compound was prepared according to the procedure of Example 1, substituting the cinnamic acid of Example 310C, giving a white solid, m.p. 161-162C. ¹H-NMR (CDCl₃ 300 MHz) δ 3.83 (s, 3H), 6.55 (d, J=9Hz, 1H), 6.70 (broad d, J=15 Hz, 1H), 6.99-7.05 (m, 2H), 7.26 (d, J=9 Hz, 1H), 7.43-7.50 (m, 2H), 8.07 (broad d, J=15 Hz, 1H) Anal. Calcd. for C₂₀H₁₉Cl₂NO₃S: C, 56.61; H, 4.51; N, 3.30. Found: C, 56.75; H, 4.57; N, 2.61.

15

Example 311

(2-Methoxyphenyl)-[2,3-dimethyl-4(E-[(morpholin-1-yl)carbonyl]ethenyl)phenyl]

sulfide

The title compound was prepared according to the procedures of Example 310. ¹H-NMR (CDCl₃ 300 MHz) δ 2.39 (s, 3H), 2.42 (s, 3H), 3.60-3.80 (m, 8H), 3.90 (s, 3H), 6.69 (d, J=15 Hz, 1H), 6.82-6.94 (m, 3H), 7.05 (d, J=9Hz, 1H), 7.20-7.30 (m,

(2H), 8.06 (d, $J=15$ Hz, 1H). Anal. Calcd. for $C_{22}H_{23}NO_3S$: C, 68.91; H, 6.57; N, 3.65.
 Found: C, 68.75; H, 6.67; N, 3.24.

Example 312

5 (2-Isopropylphenyl)[2-nitro-4-(*E*-(indol-5-ylamino)carbonyl)ethenyl] phenyl] sulfide

The title compound was prepared according to the procedures of Example 1.

1H NMR (DMSO- d_6 , 300 MHz) δ 11.04 (s, 1H), 10.10 (s, 1H), 8.52 (d, 1H, $J = 1.5$ Hz), 8.02 (s, 1H), 7.81 (dd, 1H, $J = 1.8, 8.5$ Hz), 7.53-6.63 (m, 4H), 7.39 (m, 1H), 7.25-7.35 (m, 3H), 6.94 (d, 1H, $J = 15.8$ Hz), 7.72 (d, 1H, $J = 8.5$ Hz), 6.40 (m, 1H), 3.33 (m, 1H), 1.16 (d, 6H, $J = 6.6$ Hz). MS (ESI) m/z 458, 480, 915. Anal. Calcd for $C_{26}H_{23}N_3O_3S \cdot 0.22 H_2O$: C, 67.67; H, 5.12; N, 9.10. Found: C, 67.68; H, 5.19; N, 9.08.

Example 313

15 (Benzodioxan-6-yl)[2-chloro-4-(*E*-((3-carboxypiperidin-1-yl) carbonyl)ethenyl) phenyl] sulfide

The title compound was prepared by hydrolysis of the compound of Example 308 under basic condition (aq. NaOH/EtOH). 1H NMR (DMSO- d_6 , 300 MHz) δ 1.10-1.40 (m, 2H), 1.60 (m, 1H), 1.76-1.96 (m, 3H), 2.88 (m, 1H), 3.98 (m, 1H), 3.98 (m, 1H), 4.30 (m, 4H), 6.72 (d, $J=8.0$ Hz, 1H), 7.02 (m, 3H), 7.30 (m, 2H), 7.48 (m, 1H), 7.92 (m, 1H). MS (ESI) m/z 458 (M+H) $^+$. Anal. calcd. for $C_{23}H_{21}ClNO_3SNa$: C, 55.76; H, 4.58; N, 2.83. Found: C, 55.76; H, 4.78; N, 2.63.

phenyl] sulfide

15

(Benzodioxan-6-yl)[2-chloro-4-(*E*-((4-(*tert*-butoxycarbonyl)piperazin-1-yl)

15 carboxyl)ethenyl] phenyl] sulfide

The title compound was prepared by the procedures described in Example 300 substituting ethyl isonipecotate with 1-Boc-piperazine. ¹H NMR (CDCl₃, 300 MHz) δ 1.50 (s, 9H), 3.50 (br, s 4H), 3.70 (br, 4H), 4.28-4.35 (m, 4H), 6.74 (d, J=8.0 Hz, 1H), 6.82 (m, 1H), 6.92 (d, J=8.0 Hz, 1H), 7.02 (dd, J=2.0, 8.0 Hz, 1H), 7.17 (d, J=2.0 Hz, 1H), 7.28 (d, J=8.0 Hz, 1H), 7.50 (s, 2H), 7.58 (m, 1H). MS (ESI) m/z 517 (M+H)⁺.

20 Anal. calcd. for C₂₆H₂₉ClN₂O₅S•0.1 H₂O: C, 60.19; H, 5.67; N, 5.40. Found: C, 60.20; H, 5.97; N, 5.11.

Example 316

(Benzodioxan-6-yl)[2-chloro-4-(*E*-(2-carboxypiperidin-1-yl) carbonyl)ethenyl
phenyl] sulfide

The title compound was prepared by hydrolysis of the compound of Example

5 309 under basic conditions (aq. NaOH/EtOH). ¹H NMR (DMSO-*d*₆, 300 MHz) δ
 1.10-1.40 (m, 3H), 1.45-1.60 (m, 2H), 2.25-2.45 (m, 2H), 2.55-2.80 (m, 1H), 4.30 (m,
 4H), 4.50 (m, 1H), 6.70 (d, J=8.0 Hz, 1H), 7.00 (m, 3H), 7.10 (m, 1H), 7.25 (d, J=16.0
 Hz, 1H), 7.48 (d, J=8.0 15.5 Hz, 1H), 7.90 (d, J=15.5 Hz, 1H). MS (ESI) m/z 458
 (M+H)⁺. Anal. calcd. for C₂₃H₂₁ClNO₅SNa•1.3 H₂O: C, 54.69; H, 4.73; N, 2.45.

10 Found: C, 54.67; H, 4.71; N, 2.77.

Example 317

(Benzodioxan-6-yl)[2-chloro-4-(*E*-(3-(tetrazol-5-yl)morpholin-1-yl)
carbonyl)ethenyl] phenyl] sulfide

15 The title compound was prepared by the procedures described in Example 262.

¹H NMR (CDCl₃, 300 MHz) δ 1.50-1.70 (m, 2H), 3.15 (br, 1H), 3.70-3.90 (m, 2H),
 4.25-4.35 (m, 4H), 4.55 (m, 1H), 5.04 (br, 1H), 6.72 (d, J=8.0 Hz, 1H), 6.93 (d, J=8.0
 Hz, 1H), 7.03 (dd, J=2.0, 8.0 Hz, 1H), 7.07 (d, J=2.0 Hz, 1H), 7.20-7.30 (m, 2H), 7.50
 (m, 1H), 7.65 (m, 1H). MS (ESI) m/z 486 (M+H)⁺. Anal. calcd. for C₂₂H₂₀ClN₅O₄S

20 •H₂O: C, 52.43; H, 4.40; N, 13.90. Found: C, 52.34; H, 4.35; 13.62.

Example 318

(Benzodioxan-6-yl)[2-chloro-4-(*E*-[(4-(methylaminocarbonyl)piperazin-1-yl)carbonyl]ethenyl) phenyl] sulfide

The title compound was prepared by deprotection of the of Example 315 compound using anhydrous TFA in dichloromethane, followed by treatment with

- 5 methyl isocyanate. ¹H NMR (CDCl₃, 300 MHz) δ 2.88 (s, 3H), 3.50 (br, 4H), 3.72 (br, 4H), 4.30 (m, 4H), 6.74 (d, J=8.0 Hz, 1H), 6.82 (d, J=15.0 Hz, 1H), 6.92 (d, J=8.0 Hz, 1H), 7.03 (dd, J=2.0, 8.0 Hz, 1H), 7.08 (d, J=2.0 Hz, 1H), 7.20 (d, J=8.0 Hz, 1H), 7.50 (s, 1H), 7.60 (m, 1H). MS (ESI) m/z 474 (M+H)⁺. Anal. calcd. for C₂₃H₂₄ClN₃O₄S: C, 57.63; H, 5.17; N, 8.77. Found: C, 57.53; H, 5.02; N, 8.58.

10

Example 319

(2-Methoxyphenyl)-[2,3-dichloro-4(*E*-[(4-carboxypiperidin-1-yl)carbonyl]ethenyl) phenyl] sulfide

The title compound was prepared according to the procedures of Example 310.

- 15 ¹H-NMR (CDCl₃, 300 MHz) δ 1.66-1.83 (m, 2H), 1.95-2.09 (m, 2H), 2.57-2.69 (m, 1H), 2.94-3.08 (m, 1), 3.15-3.31 (m, 1H), 3.72 (s, 3H), 3.90-4.05 (m, 1H), 4.41-4.55 (m, 1H), 6.55 (d, J=9Hz, 1H), 6.73 (d, J=15Hz, 1H), 7.00-7.05 (m, 2H), 7.27 (d, J=8Hz, 1H), 7.44-7.50 (m, 2H), 7.92 (d, J=15Hz, 1H). Anal. Calcd. for C₂₂H₂₁Cl₂NO₄S: C, 56.66; H, 4.54; N, 3.00. Found: C, 56.89; H, 4.84; N, 2.64.

20

Example 320

(Benzodioxan-6-yl)[2-chloro-4-(*E*-((4-(tetrazol-5-yl)piperidin-1-yl) carbonyl)ethenyl) phenyl] sulfide

The title compound was prepared by the procedures described in Example 314 substituting 3-(tetrazol-5-yl)piperidine with 4-(tetrazol-5-yl)piperidine. The crude reaction product was purified by reversed-phase HPLC. ¹H NMR (DMSO-d₆, 300 MHz) δ 1.22 (m, 1H), 1.55-1.75 (m, 2H), 2.06 (m, 1H), 2.45 (m, 1H), 4.22 (m, 4H), 4.30 (m, 4H), 6.70 (m, 1H), 7.00 (dd, J=2.0, 8.0 Hz, 2H), 7.25-7.40 (m, 4H), 7.50 (m, 1H). MS (ESI) m/z 484 (M+H)⁺.

Example 321

(2-Methoxyphenyl)-[3-chloro-4(*E*-[(morpholin-1-yl)carbonyl]ethenyl)phenyl]sulfide

The title compound was prepared according to the procedures of Example 1, giving a white solid, m.p. 124-125°C. ¹H-NMR (CDCl₃, 300 MHz) δ 3.60-3.80 (m, 8H), 3.85 (s, 3H), 6.80 (d, J=15 Hz, 1H), 6.95-7.01 (m, 2H), 7.05 (dd, J=9Hz, 2 Hz, 1H), 7.15 (d, J=2Hz, 1H), 7.35-7.48 (m, 3H), 7.75 (d, J=15 Hz, 1H). Anal. Calcd. for C₂₀H₂₀ClNO₃S: C, 61.61; H, 5.17; N, 3.59. Found: C, 61.43; H, 5.30; N, 3.73.

Example 322

(2-Isopropylphenyl)[2-nitro-4-(*E*-((4-oxopiperidin-1-yl)carbonyl)ethenyl) phenyl] sulfide

The title compound was prepared according to the procedures of Example 1. ¹H NMR (CDCl₃, 300 MHz) δ 8.45 (s, 1H), 7.50-7.57 (m, 3H), 7.42(br d, 1H, J = 8.1

Hz), 7.30 (m, 1H), 7.02 (br, 1H), 6.72 (d, 1H, $J = 8.4$ Hz), 4.01 (br s, 4H), 3.44 (quintet, 1H, $J = 6.8$ Hz), 2.56 (br m, 4H), 1.18 (d, 6H, $J = 7.1$ Hz). MS (ESI) m/z 425, 457. Anal. Calcd for $C_{23}H_{24}N_2O_4S$: C, 65.07; H, 5.70; N, 6.60. Found: C, 64.92; H, 5.67; N, 6.62.

5

Example 323

(Benzodioxan-6-yl)[2-trifluoromethyl-4-(*E*-((3-*R*-carboethoxypiperidin-1-yl)carbonyl)ethenyl) phenyl] sulfide

The title compound was prepared by the procedures described in Example 248, substituting ethyl (±)nipecotate with ethyl nipecotate tartrate, giving a white solid. 1H NMR ($CDCl_3$, 300 MHz) δ 1.26 (t, $J = 7.4$ Hz, 3H), 1.46-1.67 (m, 1H), 1.67-1.98 (m, 2H), 1.98-2.23 (m, 1H), 2.46-2.63 (m, 1H), 3.10-3.42 (m, 1H), 3.53-4.13 (m, 2H), 4.16 (q, $J = 7.4$ Hz, 2H), 4.25-4.40 (m, 4H), 4.60-4.88 (m, 1H), 6.91 (d, $J = 8.4$ Hz, 1H), 6.93 (d, $J = 15.3$ Hz, 1H), 6.97-7.05 (m, 2H), 7.07 (d, $J = 2.7$ Hz, 1H), 7.42 (d, $J = 8.4$ Hz, 1H), 7.59 (d, $J = 15.3$ Hz, 1H), 7.77 (s, 1H). MS (ESI⁺) (M+H)⁺ at m/z 522.

15

Example 324

(Benzodioxan-6-yl)[2-trifluoromethyl-4-(*E*-((3-*R*-carboxypiperidin-1-yl)carbonyl)ethenyl) phenyl] sulfide

The title compound was prepared by the procedures described in Example 251, substituting the ethyl ester from Example 248 with ethyl ester from Example 323, giving a white solid. 1H NMR ($CDCl_3$, 300 MHz) δ 1.48-1.71 (m, 1H), 1.71-2.01 (m,

20

2H), 2.01-2.20 (m, 1H), 2.53-2.70 (m, 1H), 3.18-3.54 (m, 1H), 3.86-4.20 (m, 2H), 4.20-4.33 (m, 4H), 4.45-4.75 (m, 1H), 6.90 (d, $J = 8.7$ Hz, 1H), 6.95-7.04 (m, 3H), 7.06 (d, $J = 2.4$ Hz, 1H), 7.35-7.45 (br m, 1H), 7.60 (d, $J = 15.3$ Hz, 1H), 7.75 (s, 1H). MS (ESI⁺) (M+H)⁺ at m/z 494.

5

Example 325

(Benzodioxan-6-yl)[2,3-dichloro-4-(E-((3-(pyrrolidin-2-on-1-yl)prop-1-ylamino)carbonyl) ethenyl)phenyl]sulfide

- The title compound was prepared by the procedures described in Example 240, substituting 4-fluoro-3-trifluoromethylbenzaldehyde with 2,3-dichloro-4-trifluoromethanesulfoxybenzaldehyde, giving a white solid. ¹H NMR (CDCl₃, 300 MHz) δ 1.71-1.82 (m, 2H), 2.08 (p, $J = 7.5$ Hz, 2H), 2.46 (t, $J = 7.5$ Hz, 2H), 3.2603.50 (m, 6H), 4.23-4.36 (m, 4H), 6.36 (t, $J = 15.6$ Hz, 1H), 6.60 (d, $J = 8.7$ Hz, 1H), 6.44 (d, $J = 8.7$ Hz, 1H), 7.03 (dd, $J = 2.4, 8.7$ Hz, 1H), 7.09 (d, $J = 2.4$ Hz, 1H), 7.31 (d, $J = 8.7$ Hz, 1H), 7.94 (d, $J = 15.6$ Hz, 1H). MS (ESI⁺) (M+H)⁺ at m/z 507, 509, 511. Anal. Calcd for C₂₄H₂₄Cl₂N₂O₄S · 1.87 H₂O: C, 53.27; H, 5.17; N, 5.18. Found: C, 53.30; H, 5.17; N, 4.83.

- (Benzodioxan-6-yl)[2,3-dichloro-4-(E-((4-acetyl)piperazin-1-yl) carbonyl)ethenyl) phenyl] sulfide

The title compound was prepared by the procedures described in Example 325,

substituting aminopropyl pyrrolidinone with 1-acetylpiperazine. white solid; ^1H NMR (CDCl_3 , 300 MHz) δ 2.17 (s, 3H), 3.50-3.94 (m, 8H), 4.26-4.40 (m, 4H), 6.61 (d, J = 8.7 Hz, 1H), 6.71 (d, J = 15.6 Hz, 1H), 6.95 (d, J = 8.4 Hz, 1H), 7.04 (dd, J = 2.4, 8.4 Hz, 1H), 7.09 (d, J = 2.4 Hz, 1H), 7.30 (d, J = 8.7 Hz, 1H), 7.99 (d, J = 15.6 Hz, 1H).

- 5 MS (ESI $^+$) ($\text{M}+\text{Na}$) $^+$ at m/z 515, 517, 519. Anal. Calcd for $\text{C}_{23}\text{H}_{22}\text{Cl}_2\text{N}_2\text{O}_4\text{S} \cdot 0.52 \text{CH}_2\text{Cl}_2$: C, 52.55; H, 4.32; N, 5.21. Found: C, 52.63; H, 4.16; N, 4.82.

Example 327

(Benzodioxan-6-yl)[2,3-dichloro-4-(*E*-(3-carboethoxypiperidin-1-yl)
carbonyl)ethenyl] phenyl] sulfide

10

The title compound was prepared by the procedures described in Example 325, substituting aminopropyl pyrrolidinone with ethyl nipecotate, giving a white solid. ^1H NMR (CDCl_3 , 300 MHz) δ 1.26 (t, J = 7.0 Hz, 3H), 1.66-1.96 (m, 2H), 1.96-2.21 (m, 1H), 2.44-2.60 (m, 1H), 2.85-3.40 (m, 2H), 3.50-3.70 (m, 1H), 3.80-4.10 (m, 2H),
 15 4.15 (q, J = 7.0 Hz, 2H), 4.26-4.40 (m, 4H), 6.66 (d, J = 8.7 Hz, 1H), 6.74 (d, J = 15.3 Hz, 1H), 6.95 (d, J = 8.4 Hz, 1H), 7.03 (dd, J = 2.4, 8.4 Hz, 1H), 7.09 (d, J = 2.4 Hz, 1H), 7.25-7.38 (m, 1H), 7.93 (d, J = 15.3 Hz, 1H). MS (ESI $^+$) ($\text{M}+\text{Na}$) $^+$ at m/z 544, 546, 548.

20

Example 328

(Benzodioxan-6-yl)[2,3-dichloro-4-(*E*-((4-carboethoxypiperidin-1-yl)
carbonyl)ethenyl] phenyl] sulfide

The title compound was prepared by the procedures described in Example 325, substituting aminopropyl pyrrolidinone with ethyl isonipecotate, giving a white solid.

¹H NMR (CDCl₃, 300 MHz) δ 1.26 (t, *J* = 7.2 Hz, 3H), 1.69 (td, *J* = 3.9, 10.8 Hz, 1H), 1.74 (td, *J* = 3.9, 10.8 Hz, 1H), 1.82-2.05 (m, 2H), 2.50-2.63 (m, 1H), 2.84-3.31 (m, 2H), 3.81-4.06 (m, 1H), 4.15 (q, *J* = 7.2 Hz, 2H), 4.24-4.34 (m, 4H), 4.34-4.59 (m, 1H), 6.61 (d, *J* = 8.7 Hz, 1H), 6.74 (d, *J* = 15.6 Hz, 1H), 6.94 (d, *J* = 8.7 Hz, 1H), 7.03 (dd, *J* = 2.7, 8.7 Hz, 1H), 7.08 (d, *J* = 2.7 Hz, 1H), 7.29 (d, *J* = 8.7 Hz, 1H), 7.90 (d, *J* = 15.6 Hz, 1H). MS (ESI⁺) (M+H)⁺ at *m/z* 522, 524, 526. Anal. Calcd for C₂₅H₂₅Cl₂NO₅S: C, 57.48; H, 4.82; N, 2.68. Found: C, 57.82; H, 4.96; N, 2.28.

Example 329

(Benzodioxan-6-yl)[2,3-dichloro-4-(*E*-((3-carboxypiperidin-1-yl) carbonyl)ethenyl)phenyl] sulfide

The title compound was prepared by the procedures described in Example 155, substituting the ethyl ester from Example 137 with the ethyl ester from Example 327,

and KOH with NaOH, providing a white solid. ¹H NMR (CDCl₃, 300 MHz) δ 1.70-2.0 (m, 2H), 2.0-2.20 (m, 1H), 2.54-2.68 (m, 1H), 3.03-3.46 (m, 2H), 3.80-4.11 (m, 2H), 4.27-4.40 (m, 4H), 4.50-4.70 (m, 1H), 6.60 (d, *J* = 8.9 Hz, 1H), 6.79 (d, *J* = 15.3 Hz, 1H), 6.94 (d, *J* = 8.5 Hz, 1H), 7.03 (dd, *J* = 2.1, 8.5 Hz, 1H), 7.08 (d, *J* = 2.1 Hz, 1H), 7.30 (d, *J* = 8.9 Hz, 1H), 7.93 (d, *J* = 15.3 Hz, 1H). MS (ESI⁺) (M-2H)⁺ at *m/z* 492, 494, 496. Anal. Calcd for C₂₃H₂₁Cl₂NO₅S · 0.73 H₂O: C, 54.43; H, 4.46; N, 2.76. Found: C, 54.43; H, 4.39; N, 2.49.

Example 330

(Benzodioxan-6-yl)[2,3-dichloro-4-(*E*-((4-carboxypiperidin-1-yl) carbonyl)ethenyl) phenyl] sulfide

- 5 The title compound was prepared by the procedures described in Example 155, substituting the ethyl ester from Example 137 with the ethyl ester from Example 328, and KOH with NaOH, to produce a white solid. ¹H NMR (d⁶-DMSO, 300 MHz) δ 1.33-1.55 (m, 2H), 1.62-1.78 (m, 2H), 1.93-2.07 (m, 1H), 2.90 (brt, *J* = 10.5 Hz, 1H), 3.16 (brt, *J* = 10.5 Hz, 1H), 3.96 (br d, *J* = 13.5 Hz, 1H), 4.09 (br d, *J* = 13.5 Hz, 1H), 10 4.26-4.42 (m, 4H), 6.60 (d, *J* = 9.0 Hz, 1H), 7.04-7.08 (m, 2H), 7.13 (d, *J* = 1.5 Hz, 1H), 7.22 (d, *J* = 15.3 Hz, 1H), 7.70 (d, *J* = 15.3 Hz, 1H), 7.86 (d, *J* = 9.0 Hz, 1H). MS (ESI⁺) (M+H)⁺ at *m/z* 516, 518, 520. Anal. Calcd for C₂₃H₂₀Cl₂N₁NaO₅S · 0.36 Et₂O: C, 54.06; H, 4.38; N, 2.58. Found: C, 53.99; H, 4.37; N, 2.22.

15

Example 331

(2-Isopropylphenyl)[2,3-dichloro-4-(*E*-((3-(1-pyrrolidin-2-onyl)propylamino) carbonyl)ethenyl) phenyl] sulfide

- The title compound was prepared by the procedures described in Example 325, substituting 6-mercaptobenzodioxane with 2-isopropylbenzenethiol, to give a white 20 solid. ¹H NMR (CDCl₃, 300 MHz) δ 1.19 (d, *J* = 7.2 Hz, 6H), 1.76 (p, *J* = 5.8 Hz, 2H), 2.08 (p, *J* = 7.65 Hz, 2H), 2.46 (t, *J* = 7.65 Hz, 2H), 3.32 (q, *J* = 5.8 Hz, 2H), 3.36-3.51 (m, 5H), 6.35 (d, *J* = 15.3 Hz, 1H), 6.40 (d, *J* = 8.7 Hz, 1H), 7.10 (brt, *J* =

7.5 Hz, 1H), 7.20-7.30 (m, 2H), 7.42-7.53 (m, 2H), 7.94 (d, $J = 15.3$ Hz, 1H). MS (ESI⁺) (M+H)⁺ at m/z 491, 493, 495. Anal. Calcd for C₂₅H₂₈Cl₂N₂O₂S · 0.7 CH₂Cl₂: C, 56.03; H, 5.38; N, 5.08. Found: C, 56.06; H, 5.22; N, 5.01.

5

Example 332

(2-Isopropylphenyl)[2,3-dichloro-4-(*E*-((4-acetylpiperazin-1-yl) carbonyl)ethenyl) phenyl] sulfide

The title compound was prepared by the procedures described in Example 326, substituting 6-mercaptobenzodioxane with 2-isopropylbenzenethiol, providing a white
 10 solid. ¹H NMR (CDCl₃, 300 MHz) δ 1.19 (d, $J = 7.2$ Hz, 6H), 2.17 (s, 3H), 3.46 (septet, $J = 7.2$ Hz, 1H), 3.50-3.90 (m, 8H), 6.41 (d, $J = 8.7$ Hz, 1H), 6.71 (d, $J = 15.3$ Hz, 1H), 7.21-7.35 (m, 2H), 7.44-7.57 (m, 3H), 7.99 (d, $J = 15.3$ Hz, 1H). MS (ESI⁺) (M+H)⁺ at m/z 477, 479, 481. Anal. Calcd for C₂₄H₂₆Cl₂N₂O₂S · 0.32 CH₂Cl₂: C, 57.89; H, 5.32; N, 5.55. Found: C, 57.85; H, 5.25; N, 5.74.

15

Example 333

(2-Isopropylphenyl)[2,3-dichloro-4-(*E*-((3-carboethoxypiperidin-1-yl) carbonyl)ethenyl) phenyl] sulfide

The title compound was prepared by the procedures described in Example 327,
 20 substituting 6-mercaptobenzodioxane with 2-isopropylbenzenethiol, giving a white solid. ¹H NMR (CDCl₃, 300 MHz) δ 1.20 (d, $J = 7.2$ Hz, 6H), 1.20-1.35 (m, 5H), 1.65-1.93 (m, 1H), 1.93-2.16 (m, 1H), 2.43-2.58 (m, 1H), 3.06-3.35 (m, 1H), 3.47

(septet, $J = 7.2$ Hz, 1H), 3.77-4.23 (m, 4H), 4.50-4.77 (m, 1H), 6.41 (d, $J = 8.4$ Hz, 1H), 6.80 (d, $J = 15.3$ Hz, 1H), 7.18-7.32 (m, 2H), 7.40-7.55 (m, 2H), 7.93 (d, $J = 15.3$ Hz, 1H). MS (ESI⁺) (M+H)⁺ at m/z 506, 508, 510.

5

Example 334

(2-Isopropylphenyl)[2,3-dichloro-4-(*E*-((4-carboethoxypiperidin-1-yl) carbonyl)ethenyl) phenyl] sulfide

The title compound was prepared by the procedures described in Example 328, substituting 6-mercaptobenzodioxane with 2-isopropylbenzenethiol, to give a white solid. ¹H NMR (CDCl₃, 300 MHz) δ 1.19 (d, $J = 7.2$ Hz, 6H), 1.26 (t, $J = 7.05$ Hz, 3H), 1.69 (td, $J = 3.9, 10.8$ Hz, 1H), 1.74 (td, $J = 3.9, 10.8$ Hz, 1H), 1.88-2.06 (m, 2H), 2.50-2.63 (m, 1H), 2.84-3.08 (m, 1H), 3.08-3.32 (m, 1H), 3.47 (septet, $J = 7.2$ Hz, 1H), 3.86-4.06 (m, 1H), 4.15 (q, $J = 7.05$ Hz, 2H), 4.37-4.61 (m, 1H), 6.40 (d, $J = 8.7$ Hz, 1H), 6.73 (d, $J = 15.6$ Hz, 1H), 7.22-7.35 (m, 2H), 7.44-7.57 (m, 3H), 7.92 (d, $J = 15.6$ Hz, 1H). MS (ESI⁺) (M+H)⁺ at m/z 506, 508, 510. Anal. Calcd for C₂₆H₂₉Cl₂NO₃S · 0.01 H₂O: C, 61.64; H, 5.77; N, 2.76. Found: C, 61.64; H, 5.90; N, 2.70.

20

Example 335

(2-Isopropylphenyl)[2,3-dichloro-4-(*E*-((3-carboxypiperidin-1-yl) carbonyl)ethenyl) phenyl] sulfide

The title compound was prepared by the procedures described in Example 329,

substituting 6-mercaptobenzodioxane with 2-isopropylbenzenethiol, giving a white solid. ^1H NMR (CDCl_3 , 300 MHz) δ 1.19 (d, $J = 7.2$ Hz, 1H), 1.43-1.67 (m, 1H), 1.67-1.97 (m, 2H), 1.97-2.19 (m, 1H), 2.52-2.64 (m, 1H), 3.04-3.38 (m, 1H), 3.47 (septet, $J = 7.2$ Hz, 1H), 3.75-4.10 (m, 2H), 4.44-4.70 (m, 1H), 6.40 (d, $J = 8.4$ Hz, 1H), 6.79 (d, $J = 15.3$ Hz, 1H), 7.18-7.29 (m, 2H), 7.41-7.53 (m, 3H), 7.93 (d, $J = 15.3$ Hz, 1H). MS (ESI^+) ($\text{M}+\text{H}$) $^+$ at m/z 478, 480, 482. Anal. Calcd for $\text{C}_{24}\text{H}_{25}\text{Cl}_2\text{NO}_3\text{S} \cdot 0.05 \text{ H}_2\text{O} \cdot 0.01 \text{ EtOH}$: C, 60.13; H, 5.29; N, 2.92. Found: C, 60.14; H, 5.11; N, 2.52.

10

Example 336

(2-Isopropylphenyl)[2,3-dichloro-4-(*E*-((4-carboxypiperidin-1-yl) carbonyl)ethenyl)phenyl] sulfide

The title compound was prepared by the procedures described in Example 330, substituting 6-mercaptobenzodioxane with 2-isopropylbenzenethiol, giving a white solid. ^1H NMR ($\text{d}^6\text{-DMSO}$, 300 MHz) δ 1.16 (d, $J = 7.2$ Hz, 6H), 1.33-1.53 (m, 2H), 1.64-1.78 (m, 2H), 1.97-2.10 (m, 1H), 2.88 (brt, $J = 10.5$ Hz, 1H), 3.15 (brt, $J = 10.5$ Hz, 1H), 3.97 (br d, $J = 13.2$ Hz, 1H), 4.11 (br d, $J = 13.2$ Hz, 1H), 6.41 (d, $J = 9.0$ Hz, 1H), 7.22 (d, $J = 15.6$ Hz, 1H), 7.31-7.42 (m, 1H), 7.53 (d, $J = 7.8$ Hz, 1H), 7.56-7.64 (m, 2H), 7.71 (d, $J = 15.6$ Hz, 1H), 7.85 (d, $J = 9.0$ Hz, 1H). MS (ESI^+) ($\text{M}+\text{H}$) $^+$ at m/z 478, 480, 482. Anal. Calcd for $\text{C}_{24}\text{H}_{24}\text{Cl}_2\text{NNaO}_3\text{S} \cdot 0.95 \text{ H}_2\text{O}$: C, 55.70; H, 5.04; N, 2.71. Found: C, 55.69; H, 4.90; N, 2.57.

Example 337

(1-Methylindol-5-yl)[2,3-dichloro-4-(*E*-((3-carboethoxypiperidin-1-yl) carbonyl)ethenyl) phenyl] sulfide

The title compound was prepared by the procedures described in Example 283,

- 5 substituting 4-fluoro-3-chlorobenzaldehyde with 2,3-dichloro-4-trifluoromethanesulfoxybenzaldehyde, giving a white solid. ¹H NMR (CDCl₃, 300 MHz) δ 1.23 (t, *J* = 7.5 Hz, 3H), 1.46-1.67 (m, 1H), 1.67-1.95 (m, 2H), 1.95-2.17 (m, 1H), 2.43-2.60 (m, 1H), 3.02-3.42 (m, 1H), 3.67-3.92 (m, 2H), 3.86 (s, 3H), 4.13 (q, *J* = 7.5 Hz, 2H), 4.59-4.80 (m, 1H), 6.46 (d, *J* = 8.7 Hz, 1H), 6.54 (d, *J* = 3.0 Hz, 1H),
- 10 6.77 (d, *J* = 15.3 Hz, 1H), 7.15 (d, *J* = 3.0 Hz, 1H), 7.19 (d, *J* = 8.7 Hz, 1H), 7.37 (d, *J* = 8.7 Hz, 1H), 7.42 (d, *J* = 8.7 Hz, 1H), 7.89 (s, 1H), 7.92 (d, *J* = 15.3 Hz, 1H). MS (ESI⁺) (M+H)⁺ at *m/z* 517, 519, 521.

Example 338

- 15 (1-Methylindol-5-yl)[2,3-dichloro-4-(*E*-((3-carboxypiperidin-1-yl) carbonyl)ethenyl) phenyl] sulfide

- The title compound was prepared by the procedures described in Example 155, substituting the ethyl ester from Example 137 with ethyl ester from Example 337, and KOH with NaOH, to give a white solid. ¹H NMR (d⁶-DMSO, 300 MHz) δ 1.29-1.45
- 20 (m, 1H), 1.45-1.78 (m, 2H), 1.78-2.02 (m, 1H), 2.20-2.40 (m, 1H), 2.82 (brt, *J* = 10.5 Hz, 1H), 3.08 (brt, *J* = 10.5 Hz, 1H), 3.80-4.07 (m, 2H), 3.86 (s, 3H), 4.38-4.50 (m, 1H), 6.42 (d, *J* = 8.4 Hz, 1H), 6.54 (d, *J* = 3.0 Hz, 1H), 7.19 (d, *J* = 15.3 Hz, 1H), 7.32

(dd, $J = 1.8, 8.7$ Hz, 1H), 7.48 (d, $J = 3.0$ Hz, 1H), 7.64 (d, $J = 8.7$ Hz, 1H), 7.67-7.77 (m, 2H), 7.87 (d, $J = 1.8$ Hz, 1H). MS (ESI⁺) (M+H)⁺ at m/z 489, 491, 493. Anal. Calcd for C₂₄H₂₂Cl₂N₂O₃S · 0.56 CH₂Cl₂: C, 54.94; H, 4.34; N, 5.22. Found: C, 54.89; H, 4.44; N, 5.32.

5

Example 339

(1-Methylindol-5-yl)[2,3-dichloro-4-(*E*-((4-carboethoxypiperidin-1-yl) carbonyl)ethenyl) phenyl] sulfide

The title compound was prepared by the procedures described in Example 285, substituting 4-fluoro-3-chlorobenzaldehyde with 2,3-dichloro-4-trifluoromethanesulfoxybenzaldehyde, providing a white solid. ¹H NMR (CDCl₃, 300 MHz) δ 1.25 (t, $J = 7.2$ Hz, 3H), 1.62-1.79 (m, 2H), 1.87-2.04 (m, 2H), 2.41-2.63 (m, 1H), 2.85-3.41 (m, 2H), 3.85 (s, 3H), 3.87-4.10 (m, 1H), 4.15 (q, $J = 7.2$ Hz, 2H), 4.32-4.60 (m, 1H), 6.46 (d, $J = 8.7$ Hz, 1H), 6.54 (d, $J = 3.0$ Hz, 1H), 6.71 (d, $J = 15.3$ Hz, 1H), 7.15 (d, $J = 3.0$ Hz, 1H), 7.17 (d, $J = 8.7$ Hz, 1H), 7.36 (dd, $J = 2.4, 8.4$ Hz, 1H), 7.42 (d, $J = 8.4$ Hz, 1H), 7.88 (d, $J = 2.4$ Hz, 1H), 7.90 (d, $J = 15.3$ Hz, 1H). MS (ESI⁺) (M+H)⁺ at m/z 517, 519, 521. Anal. Calcd for C₂₆H₂₆Cl₂N₂O₃S · 0.12 H₂O: C, 60.10; H, 5.09; N, 5.39. Found: C, 60.09; H, 5.21; N, 5.54.

20

Example 340

(1-Methylindol-5-yl)[2,3-dichloro-4-(*E*-((4-carboxypiperidin-1-yl) carbonyl)ethenyl) phenyl] sulfide

The title compound was prepared by the procedures described in Example 155, substituting the ethyl ester from Example 137 with ethyl ester from Example 339, and KOH with NaOH, to give a white solid. ^1H NMR (d^6 -DMSO, 300 MHz) δ 1.31-1.53 (m, 2H), 1.62-1.76 (m, 2H), 1.94-2.09 (m, 1H), 2.88 (brt, $J = 10.5$ Hz, 1H), 3.13 (brt, $J = 10.5$ Hz, 1H), 3.86 (s, 3H), 3.93 (br d, $J = 13.2$ Hz, 1H), 4.09 (br d, $J = 13.2$ Hz, 1H), 6.41 (d, $J = 8.7$ Hz, 1H), 6.53 (dd, $J = 0.9, 3.0$ Hz, 1H), 7.04 (d, $J = 15.3$ Hz, 1H), 7.32 (dd, $J = 2.1, 8.7$ Hz, 1H), 7.48 (d, $J = 3.0$ Hz, 1H), 7.64 (d, $J = 8.7$ Hz, 1H), 7.69 (d, $J = 15.3$ Hz, 1H), 7.73 (d, $J = 8.7$ Hz, 1H), 7.88 (d, $J = 2.1$ Hz, 1H). MS (ESI $^+$) (M+H) $^+$ at m/z 489, 491, 493. Anal. Calcd for $\text{C}_{24}\text{H}_{21}\text{Cl}_2\text{N}_2\text{NaO}_3\text{S} \cdot 0 \text{ H}_2\text{O}$: C, 56.37; H, 4.14; N, 5.48. Found: C, 56.44; H, 4.38; N, 5.20.

An alternative method for preparing Example 340 is given below.




Example 340A

1-Methyl-5-iodoindole

To a solution of 5-iodoindole (75 g, 0.31 mol) in dry THF (750mL), at -78°C was added sodium hydride (60% in mineral oil, 14.85 g, 0.37 mol) in one portion. The suspension was stirred at -78°C for 1 hour after which iodomethane (28.8 mL, 0.46 mol) was added. The reaction mixture was stirred overnight with a slow elevation on temperature to room temperature (no more dry ice was added). Ether (600mL) and hexane (1.2L) were added and the mixture was washed with brine (1.6L) and water (1.5L), dried over Na_2SO_4 and filtered. The solution was concentrated and the residual brown solid was recrystallized from hexane to give the title compound

(66 g). The impure fraction from the mother liquor was flash chromatographed (8% EtOAc in hexane) to give an additional quantity of desired product (12.5 g, combined yield of 99%). MS (DCI/NH₃) m/e 258 (M+H)⁺.



Example 340B

1-Methyl-S-triisopropylsilyl-5-indolethiol


Potassium hydride (35% in mineral oil, 12.03 g, 0.105 mol) was charged to a 250 mL RBF and was washed with dry THF (2x50mL). The resultant KH powder was then suspended in dry THF (75 mL), and cooled to 5 °C. Triisopropylsilylthiol (20.0 g, 0.105 mol) was slowly added via syringe over a period of 15 minutes. Vigorous escape of hydrogen gas was observed with addition of the thiol. The suspension was stirred at 5°C for 1 our and became homogenous. After another hour stirring at room temperature, this solution was cannulated to a THF solution (100mL) containing Example 340A (24.5 g, 95.5 mmol) and tetrakis(triphenylphosphine)palladium(0) (2.2 g, 1.91 mmol). The yellow suspension was stirred at 70°C for 1 hour. After cooled, ether and hexane were added, and the mixture was washed with brine, dried (Na₂SO₄) and concentrated. The residual oil was purified by flash chromatography (silica gel, 3% EtOAc in hexane) to give the title compound (26.7 g, 88%). MS (DCI/NH₃) m/e 320 (M+H)⁺.

290

Example 340C4-Bromo-2,3-dichlorophenol

To a solution of 2,3-dichlorophenol (200 g, 1.227 mol) in dichloromethane
 5 (800 mL), at 0 °C was added dropwise bromine (196.1g, 1.227 mol) from a dropping
 funnel within 1 hour. The red solution was stirred overnight (0°C - rt), and washed
 with 10% NaHSO₃. The organic phase was dried over Na₂SO₄, and concentrated. The
 residual white solid was recrystallized from hexane to give example 340C as white
 needles (207g, 70%). MS (DCI/NH₃) *m/e* 241 (M+H)⁺.


10

Example 340DMethyl 2,3-dichloro-4-(4-hydroxyphenyl)acrylate

A 1 L RBF was charged with Example 340C (48.4 g, 0.2 mol), Pd₂(dba)₃ (4.6
 15 g, 5 mmol), (Tol)₃P (4.66 g, 15.2 mmol), and purged with nitrogen. Dry DMF (300
 mL), methyl acrylate (51.66 g, 0.6 mol) and triethylamine (84 mL, 0.6 mol) were then
 added. The reaction mixture was purged with nitrogen and stirred at 100°C (oil bath)
 for 16 hours. After cooled to room temperature, a lot of white crystalline material
 formed. Ethyl acetate (500 mL) and brine (not saturated, 800 mL) were added, and
 20 stirred. The white crystalline material dissolved. A little insoluble black solid (Pd)

was filtered off. To the solution was then added, with stirring, saturated NaCl solution (2 L) and hexane (500 mL). The mixture was stirred for 1 hour. The formed yellowish solid was collected by filtration, washed with water (400 mL), acetonitrile (50 mL) and 1:1 ethyl acetate/hexane (500 mL), and dried to give pure desired


5 compound (44.99g, 91%). MS (DCI/NH₃) *m/e* 247 (M+H)⁺.



Example 340E

Methyl 2,3-dichloro-4-(trifluoromethanesulfonyl)phenylacrylate


10 To a suspension of Example 340D (18.62 g, 75.4 mmol) in pyridine (150 mL) at 5 °C was added trifluoromethylsulfonyl anhydride (25.53 g, 90 mmol) very slowly. The suspension was stirred at 5 °C for 1 hour and became homogeneous. The solution was kept at 5 °C for 2 hours and at room temperature for 20 minutes. Ether (700mL) was added and the mixture was washed 10%HCl (700 mL)/brine (300 mL), 10%HCl
15 (100 mL)/brine (900 mL), and brine (500 mL). The organic phase was dried (Na₂SO₄) and concentrated to give the title compound (24.86 g, 87%). MS (DCI/NH₃) *m/e* 379 (M+H)⁺.



Example 340F(1-Methylindol-5-yl)[2,3-dichloro-4-(E-(carboxyethenyl)phenyl] sulfide


To a solution of Example 340B (38.5 g, 0.12 mol) and Example 340E (30.3 g, 0.08 mol) in dry N-methylpyrrolidinone (300 mL) was added CsF (18.2 g, 0.12 mol) at 5°C under nitrogen atmosphere. After 1 hour stirring at the same temperature, the cooling bath was removed, and the mixture was stirred at room temperature for 0.5 hour. Ethyl acetate (800 mL) was added, and the mixture was washed with brine and water, and concentrated. The residual oil was separated by flash chromatography (20% EtOAc/hexane) to give a yellow solid (30 g).

This yellow solid was dissolved in THF (150 mL), and was added a solution of LiOH (4.0 g, 0.16 mol) in H₂O (50 mL). The mixture was stirred at room temperature for 1 hour and more water (100 mL) was added to form a transparent solution. After overnight stirring the solution was acidified with 10 % aq. HCl. The mixture was concentrated under reduced pressure to about 100 mL. The formed solid material was collected by filtration, washed with water (200 mL), acetonitrile (30 mL), 1:1 ether/hexane, and dried to give the title compound (22.3 g, overall 74%). MS (DCI/NH₃) *m/e* 378 (M+H)⁺.

Example 340G

(1-Methylindol-5-yl)[2,3-dichloro-4-(*E*-((4-carbomethoxypiperidin-1-yl)carbonyl)ethenyl)phenyl] sulfide

To a solution of Example 340F (9.5 g, 25.1 mmol) and methyl isonipicotate (7.19 g, 50.2 mmol) in DMF (70 mL) was added EDC (9.64 g, 50.2 mmol), HOBT (6.78 g, 50.2 mmol) and triethylamine (7.0 mL, 50.2 mmol). The reaction mixture was stirred at room temperature for 15 hours. Ethyl acetate (800 mL) was added, and the mixture was washed with brine, and concentrated. The residue was purified by flash chromatography (60% EtOAc in hexane) to give example 340G as white powder (10.86 g, 94%). MS (ESI⁺) *m/z* 503 (M+H)⁺.



Example 340

(1-Methylindol-5-yl)[2,3-dichloro-4-(*E*-((4-carboxypiperidin-1-yl)carbonyl)ethenyl)phenyl] sulfide, sodium salt

To a suspension of Example 340G (11.8 g, 23.6 mmol) in THF (150 mL) was added a solution of lithium hydroxide monohydrate (1.98 g, 47.2 mmol) in H₂O (30 mL). The mixture was stirred at room temperature overnight. Water (120 mL) was added and formed transparent solution was stirred for another hour before 10% HCl (30 mL) was added. The mixture was concentrated under reduced pressure to about 120 mL. The formed solid material was collected by filtration, washed with water, acetonitrile, and dried to give a white solid (11.0 g).

10.50 grams of the solid was suspended in methanol (60 mL), and was treated with a solution NaOH (0.859g) in methanol (20 mL). After all of the solid material went into solution, the solvent was removed under reduced pressure. The residual yellow oil was triturated with ether, and dried to give the title compound as yellow powder (11.33 g, 95%).

Example 341

(2-Ethoxyphenyl)-[2,3-dichloro-4(E-[4-carboxypiperidin-1-yl]carbonyl]ethenyl)phenyl] sulfide

10 The title compound was prepared according to the procedures of Example 310, substituting 2-ethoxybenzenethiol prepared according to the procedures of Example 97A. ¹H-NMR (CD₃OD, 300 MHz) Potassium salt δ 1.20 (t, J=7Hz, 3H), 1.55-1.72 (m, 2H), 1.88-1.98 (m, 2H), 2.32 (m, 1H), 2.88 (t, J=12Hz, 1H), 3.20 (t, J=12 Hz, 1H), 4.05 (q, J=7Hz, 2H), 4.14 (d, J=12 Hz, 1H), 4.48, (d, J= 12 Hz, 1H), 6.64 9d, J=9Hz, 15 1H), 7.00-7.15 (m, 3H), 7.44-7.50 (m, 2H), 7.56 (d, J=9Hz, 1H), 7.90 (d, J=15 Hz, 1H) Anal. Calcd. for C₂₃H₂₂KCl₂NO₄S 0.5 H₂O: C, 52.37, H, 4.39, N, 2.66. Found: C, 52.23; H, 4.56; N, 2.49.

Example 342

20 (2-Ethoxyphenyl)-[2,3-dichloro-4(E-[morpholin-1-yl]carbonyl]ethenyl)phenyl]

sulfide

The title compound was prepared according to the procedures of Example 310,

- substituting 2-ethoxybenzenethiol prepared according to the procedures of Example 97A. ¹H-NMR (CDCl₃, 300 MHz) δ 1.25 (t, J=7Hz, 3H), 3.55-3.80 (m, 8H), 4.05 (q, J=7Hz, 2H), 6.63 (d, J=9Hz, 1H), 6.71 (d, J=15 Hz, 1H), 6.95-7.03 (m, 2H), 7.26 (d, J=9Hz, 1H), 7.39-7.50 (m, 2H), 7.99 (d, J=15 Hz, 1H) Anal. Calcd. for
- 5 C₂₁H₂₁Cl₂NO₃S: C, 57.54; H, 4.82; N, 3.20. Found: C, 57.55; H, 4.77; N, 3.14.

Example 343

(2-Ethoxyphenyl)-[2,3-dichloro-4(E-[(3-carboxypiperidin-1-yl)carbonyl]ethenyl)phenyl] sulfide

- 10 The title compound was prepared according to the procedures of Example 310, substituting 2-ethoxybenzenethiol prepared according to the procedures of Example 97A. ¹H-NMR (CD₃OD 300MHz) δ 1.20 (t, J=7Hz, 3H), broad peaks totaling 9 protons at 1.4-1.95, 2.0-2.14, 2.22-2.35, 2.75-3.134.10-4.34, 4.69-4.76, 4.05 (q, J=7Hz, 2H), 6.64 (d, J=9Hz, 1H), 7.03 (t, J=8Hz, 1H), 7.10 (d, J=9Hz, 1H), 7.22 (d, J=15 Hz, 1H), 7.45-7.50 (m, 2H), 7.62 (d, J=9Hz, 1H), 7.80 (d, J=15 Hz, 1H). The
- 15 acid (303 mg, 0.63 mmol) was dissolved in 3 mL of methanol. A solution of KOH (0.60 mmol) in 1 mL of methanol was added. The resultant solution was stirred for 5 min and concentrated in vacuo. Ether (5 mL) was added, and the mixture was stirred for 1 hr. The resultant powder was collected by filtration and dried under vacuum at
- 20 60C to give 307 mg of a solid, water-soluble product. Anal. Calcd. for C₂₃H₂₂KC₆N₄O₄S 0.5 H₂O: C, 52.37; H, 4.39; N, 2.66. Found: C, 52.20; H, 4.65; N, 3.04.

Example 344

(2-Isopropylphenyl)[2-nitro-4-(*E*-((3-carboethoxypyrrolidin-1-yl)carbonyl)ethenyl)
phenyl] sulfide

5 Prepared according to the procedures of Example 71, giving a yellow solid.

¹H NMR (DMSO-d₆, 300MHz) δ 1.14 (d, J = 7.0 Hz, 6H); 1.20 (t, J= 7.0 Hz, 3H);
 1.92-2.30 (m, 2H); 3.10-4.01 (m, 6H); 4.06-4.17 (m, 2H); 6.64 (d, J = 8.5 Hz, 1H);
 7.06-7.17 (m, 1H), 7.34-7.62 (m, 5H); 7.88-7.96 (m, 1H); 8.62 (dd, J = 1.5, 8.5 Hz,
 1H). MS (APCI) (M+H)⁺ at m/z 469. Anal calcd for C₂₅H₂₈N₂S₁O₅: C, 64.08; H, 6.02;

10 N, 5.98. Found: C, 64.12; H, 5.98; N, 5.89.

Example 345

(2-Isopropylphenyl)[2-nitro-4-(*E*-((3-carboxypyrrolidin-1-yl)carbonyl)ethenyl)
phenyl] sulfide

15 Prepared according to the procedures of Example 71, giving a yellow solid.

¹H NMR (DMSO-d₆, 300MHz) δ 1.14 (d, J = 6.8 Hz, 6H); 1.92-2.24 (m, 2H); 3.01-
 3.92 (m, 6H); 6.64 (dd, J = 1.7, 8.5 Hz, 1H); 7.04-7.16 (m, 1H), 7.33-7.61 (m, 5H);
 7.87-7.95 (m, 1H); 8.61 (dd, J = 1.7, 8.5 Hz, 1H). MS (APCI) (M+H)⁺ at m/z 441.

Anal calcd for C₂₃H₂₄N₂S₁O₅: C, 62.71; H, 5.49; N, 6.36. Found: C, 62.47; H, 5.39; N,

20 6.09.

Example 346

(2-Isopropylphenyl)[2,3-difluoro-4-(*E*-((3-carboethoxypiperidin-1-yl)carbonyl)ethenyl) phenyl] sulfide

Prepared according to the procedures of Example 71, giving a yellow solid.

¹H NMR (DMSO-d₆, 300MHz) δ 1.18 (d, J = 7.0 Hz, 6H); 1.10-1.22 (m, 3H); 1.30-2.07 (br m, 4H); 2.50-3.45 (br m, 3H); 3.55-4.47 (br m, 5H); 6.62-6.72 (m, 1H); 7.23-7.73 (m, 7H). MS (APCI) (M+H)⁺ at m/z 474.

Example 347

(2-Isopropylphenyl)[2,3-difluoro-4-(*E*-((3-carboxypiperidin-1-yl)carbonyl)ethenyl) phenyl] sulfide

Prepared according to the procedures of Example 71, giving a yellow solid.

¹H NMR (DMSO-d₆, 300MHz) δ 1.18 (d, J = 7.0 Hz, 6H); 1.30-2.03 (br m, 4H); 2.25-3.50 (br m, 4H); 3.87-4.51 (br m, 2H); 6.62-6.72 (m, 1H); 7.23-7.73 (m, 7H). MS (APCI) (M+H)⁺ at m/z 446.

Example 348

(2-Isopropylphenyl)[2,3-difluoro-4-(*E*-((4-carboxypiperidin-1-yl)carbonyl)ethenyl) phenyl] sulfide

Prepared according to the procedures of Example 71, giving a yellow solid.

¹H NMR (DMSO-d₆, 300MHz) δ 1.18 (d, J = 6.8Hz, 6H); 1.30-1.91 (br m, 4H); 2.50-3.50 (br m, 4H); 4.02-4.34 (br m, 2H); 6.62-6.72 (m, 1H); 7.23-7.73 (m, 7H). MS (APCI) (M+H)⁺ at m/z 446.

Example 349

(Benzodioxan-6-yl)[2-trifluoromethyl-4-(E-((3-ethoxycarbonylpyrrolidin-1-yl)carbonyl)ethenyl) phenyl] sulfide

5 The title compound was prepared according to the procedures of Example 1.

¹H NMR (CDCl₃, 300 MHz) δ 7.77 (s, 1H), 7.62 (d, 1H, J = 15.4 Hz) 7.42 (d, 1H, J = 8.5 Hz), 7.06 (d, 1H, J = 2.1 Hz), 6.98-7.04 (m, 2H), 6.91 (d, 1H, J = 8.1 Hz), 6.68 (dd, 1H, J = 3.3, 15.3 Hz), 4.30 (m, 4H), 4.19 (q, 2H, J = 7.0 Hz), 3.56-3.92 (m, 4H), 3.06-3.24 (m, 1H), 2.10-2.35 (m, 2H), 1.28 and 1.29 (two t, 3H, J = 7.2 Hz). MS (ESI)

10 *m/z* 508, 1015.

Example 350

(Benzodioxan-6-yl)[2-trifluoromethyl-4-(E-((3-carboxypyrrolidin-1-yl)carbonyl)ethenyl) phenyl] sulfide

15 The title compound was prepared by hydrolysis of the compound of Example 349 according to standard procedures. ¹H NMR (DMSO-d₆, 300 MHz) δ 8.10 (d, 1H, J = 9.9 Hz), 7.84 (t, 1H, J = 7.8 Hz), 7.46 (d, 1H, J = 15.3 Hz), 7.10 (d, 1H, J = 15.3 Hz), 6.97-7.06 (m, 4H), 4.30 (m, 4H), 3.50 (br, overlapped with water residue peak), 3.00 (m, 1H), 2.10 (m, 1H), 2.00 (m, 1H). MS (ESI) *m/z* -478, -957.

20

Example 351

(2-Methoxyphenyl)[2-chloro-3-trifluoromethyl-4-(E-((4-carboethoxypiperidin-1-yl)carbonyl)ethenyl) phenyl] sulfide

Example 351A

5 3-Chloro-4-hydroxy-2-(trifluoromethyl)benzaldehyde

Chloroform (6.7g, 2.0 eq.) was added dropwise to a stirred mixture of Ca(OH)₂ (8.95g, 120 mmol.), K₂CO₃ (13.5g, 98 mmol.), 2-chloro-3-(trifluoromethyl)phenol (5.0g, 22 mmol.), and H₂O (50 mL) at 60°-70° over 2 h. The reaction mixture was cooled, and acidified with conc. HCl. The product was extracted into EtOAc and dried
10 over Na₂SO₄. Solvent was evaporated, the crude product was separated and purified through a silica column, eluting with hexane and EtOAc (3:2) to give 580 mg (10%) of the title compound.

Example 351B

15 (2-Methoxyphenyl)[2-chloro-3-trifluoromethyl-4-(E-carboxyethenyl) phenyl] sulfide

The title compound was prepared according to the procedures described in Example 310, substituting the compound of Example 351A for 4-hydroxy-2,3-dichlorobenzaldehyde.

20 Example 351C

(2-Methoxyphenyl)[2-chloro-3-trifluoromethyl-4-(E-((4-carboethoxypiperidin-1-yl)carbonyl)ethenyl) phenyl] sulfide

- To the acyl chloride (37 mg, 0.1 mmol) prepared from the compound of Example 351B, as a solution in CH_2Cl_2 was added 1.2 eq. of ethyl isonipicotate and 1.2 eq. of Hunig's base. The mixture was stirred at room temperature for 20 min., ~90% of the solvent was removed in vacuo, and the resultant solution was loaded on a
- 5 silica column to elute with hexane and EtOAc (3:2) to give 51mg (98%) of the title compound. $^1\text{H-NMR}$ (CDCl_3 , 300MHz) δ 1.25 (t, $J=7.5\text{Hz}$, 3H), 1.65-1.78 (m, 2H), 1.92-2.02 (br, 2H), 2.51-2.60 (m, 1H), 2.93-3.24 (br, 2H), 3.82 (s, 3H), 3.88-3.96 (m, 1H), 4.15 (q, $J=7.5\text{Hz}$, 2H), 4.40-4.50 (br, 1H), 6.48 (d, $J=15\text{Hz}$, 1H), 6.72 (d, $J=9\text{Hz}$, 1H), 7.02 (d, $J=7.5\text{Hz}$, 2H), 7.12 (d, $J=9\text{Hz}$, 1H), 7.49 (t, $J=9\text{Hz}$, 2H), 7.86
- 10 (qq, $J=4.5\text{Hz}$, 1H). MS (DCI/NH_3) m/e 528 ($\text{M}+\text{H}$) $^+$.

Example 352

(2-Methoxyphenyl)[2-chloro-3-trifluoromethyl-4-(*E*-(4-carboethoxypiperidin-1-yl)carbonyl)ethenyl) phenyl] sulfide

- 15 The compound of Example 351 was hydrolyzed by aq. NaOH in EtOH at rt. to give 90% yield of the title compound. $^1\text{H NMR}$ (DMSO, 300MHz) δ 1.37-1.52 (br, 2H), 1.78-1.86 (br, 2H), 2.45-2.55 (m, 1H), 2.83 (t, $J=12\text{Hz}$, 1H), 3.17 (t, $J=13.5\text{ Hz}$, 1H), 3.80 (s, 3H), 4.07 (d, $J=12\text{Hz}$, 1H), 4.26 (d, $J=13.5\text{Hz}$, 1H), 6.75 (d, $J=9\text{Hz}$, 1H), 6.98 (d, $J=15\text{Hz}$, 1H), 7.11(t, $J=9\text{Hz}$, 1H), 7.26 (d, $J=9\text{Hz}$, 1H), 7.53 (d, $J=7.5\text{Hz}$, 1H),
- 20 7.62 (d, $J=9\text{Hz}$, 2H), 7.70 (qq, $J=4.5\text{Hz}$, 1H). MS (DCI/NH_3) m/e 500($\text{M}+\text{H}$) $^+$.

Example 353


(2-Methoxyphenyl)[2-chloro-3-trifluoromethyl-4-(*E*-((morpholin-1-yl)carbonyl)ethenyl) phenyl] sulfide

Prepared according to the procedures of Example 351, giving 50 mg (91%) of the title compound. ¹H-NMR (CDCl₃, 300MHz) δ 3.56-3.62 (br m, 2H), 3.67-3.77 (br m, 6H), 3.85 (s, 3H), 6.45 (d, J=15Hz, 1H), 6.73 (d, J=9Hz, 1H), 7.03 (d, J=9Hz, 2H), 7.09 (t, J=9Hz, 1H), 7.52 (d, J=9Hz, 2H), 2.93 (qq, J=6Hz, 1H). MS (DCI/NH₃) *m/z* 458 (M+H)⁺.

Example 354

(Benzodioxan-6-yl) [4-(*E*-((4-carboxypiperidin-1-yl) carbonyl)ethenyl)naphthyl] sulfide


The methods of Example 310 and 311 were used to convert 4-hydroxy-2-naphthaldehyde and 6-benzodioxanethiol to the desired product as a yellow solid. ¹H NMR (DMS-d₆, 300MHz) δ 1.50 (br s, 2H), 1.83-1.92 (m, 2H), 2.5-2.6 (m, 1H), 2.85-2.95 (m, 1H), 3.18-3.29 (m, 1H), 4.22 (br s, 5H), 4.30-4.38 (m, 1H), 6.87-6.92 (m, 3H), 7.38 (d, J=15Hz, 1H), 7.45 (d, J=7.5Hz, 1H), 7.64-7.70 (m, 2H), 7.93 (d, J=7.5Hz, 1H), 8.20-8.45 (m, J=3H). MS(ESI⁺) *m/z* 476 (M+H)⁺. Anal calcd for C₂₇H₂₅NO₃S·0.67H₂O: C, 66.50; H, 5.44; N, 2.87. Found: C, 66.56; H, 5.81; N, 2.49.



Example 355

(2-Methoxyphenyl) [2,3-dichloro-4-(E-((4-(spiro-hydantoin-5-yl)-piperidin-1-yl)carbonyl)ethenyl)phenyl] sulfide


- The title compound was prepared from Example 310C, using the procedures described in Example 340 and substituting methyl isonipecotate with piperidine-4-spiro-5'-hydantoin, which was prepared according to a literature method (Wysong, C., et al, *J. Org. Chem.* **1996**, 7650). ¹H NMR (300 MHz, DMSO-d₆) δ 1.65 (m, 2H), 1.75 (m, 2H), 3.05 (m, 1H), 3.50 (m, 1H), 4.12 (m, 1H), 4.20 (m, 1H), 6.56 (d, J=6.5Hz, 1H), 7.10 (t, J=8.0Hz, 1H), 7.22 (d, J=8.0 Hz, 1H), 7.28 (d, J=15.6 Hz, 1H), 7.49 (dd, J=8.0, 1.7Hz, 1H), 7.56 (t, J=8.2Hz, 1H), 7.76 (d, J=15.6Hz, 1H), 7.84 (d, J=8.6Hz, 1H), 8.58 (s, 1H), 10.73(s, 1H). MS (ESI⁺) *m/z* 504 (M-H)⁺.

Example 356

- (2-Methoxyphenyl) [2,3-dichloro-4-(E-(4-(2-(2-hydroxyethoxy)ethyl)piperazin-1-yl)carbonyl)ethenyl)phenyl] sulfide

- The title compound was prepared from Example 310C by the procedures described in Example 340 and substituting methyl isonipecotate with N-[2-(2-hydroxyethoxy)ethyl]piperazine. ¹H NMR (300 MHz, DMSO-d₆) δ 3.10 (m, 2H), 3.50 (m, 4H), 4.50 (m, 2H), 4.70 (s, 1H), 6.57 (d, J=8.5Hz, 1H), 7.09 (t, J=8.0Hz, 1H), 7.23 (d, J=8.0Hz, 1H), 7.26 (d, J=15.5Hz, 1H), 7.49 (dd, J=7.8, 1.7Hz, 1H), 7.57 (t,

J=8.2Hz, 1H), 7.78 (d, J=15.6Hz, 1H), 7.80 (d, J=7.8Hz, 1H). MS (ESI⁺) *m/z* 545 (M-H)⁻.




5

Example 357

(2-Methoxyphenyl)[2,3-dichloro-4-(E-((4-ethylpiperazin-1-yl)carbonyl)ethenyl)phenyl] sulfide

The title compound was prepared from Example 310C by the procedures described in Example 340 and substituting methyl isonipicotate with 1-

10 ethylpiperazine. ¹H NMR (300 MHz, CDCl₃) δ 1.09 (t, J=7.1Hz, 3H), 2.42 (q, J=7.1Hz, 2H), 2.47 (m, 4H), 3.60 (m, 2H), 3.75 (m, 2H), 3.82 (s, 3H), 6.56 (d, J=8.5Hz, 1H), 6.74(d, J=15.3Hz, 1H), 7.02 (m, 2H), 7.26 (d, J=8.5Hz, 1H), 7.46 (m, 2H), 7.94 (d, J=15.5Hz, 1H). MS (ESI⁺) *m/z* 451 (M+H)⁺.



15

Example 358

(2-Isopropylphenyl)[2,3-dichloro-4-(E-((2-(2-hydroxyethoxy)ethyl)piperazin-1-yl)carbonyl)ethenyl] sulfide

The title compound was prepared from the cinnamide acid of Example 331, using the procedures described in Example 340 and substituting methyl isonipecotate with N-[2-(2-hydroxyethoxy)ethyl]piperazine. ¹H NMR (300 MHz, DMSO-d₆) δ 1.18 (d, 6H), 3.0 (m, 3H), 3.30 (m, 2H), 3.50 (m, 10H), 3.80 (m, 2H), 4.50 (t, 1H), 6.45 (d, 1H), 7.30 (d, 1H), 7.35 (dd, 1H), 7.55 (d, 1H), 7.60 (m, 2H), 7.75 (d, 1H), 7.80 (d, 1H). MS (ESI⁺) *m/z* 523 (M+H)⁺.

Example 359

(Benzodioxan-6-yl)[2,3-bis(trifluoromethyl)-4-(E-((4-carboxypiperidin-1-yl)carbonyl)ethenyl)phenyl]sulfide

Example 359A

1-Methyl-2,3-bis(trifluoromethyl)-7-oxabicyclo[2.2.1]hepta-2,5-diene

Hexafluoro-2-butyne (21.0 g, 0.13 mol) was transferred into a reaction bottle and added 2-methylfuran (12.86 g, 0.157 mol). This resulting mixture bottle was sealed and heated for 15 hr. at 120 °C. After cooling, the excess 2-methylfuran was rotoevaporated in vacuo at rt, to give crude title product (29 g, 92%), which was used directly.

Example 359B

4-Methyl-2,3-bis(trifluoromethyl)phenol

A mixture of Example 359A (12.0 g, 0.05 mol) and boron trifluoride-diethyl ether complex (150 ml) was stirred at room temp overnight, then neutralized carefully with 20% aqueous potassium carbonate, then the mixture was extracted with ether. The ether layer was dried over MgSO_4 and evaporated under reduced pressure to
5 afford 10.4g (85%) of the title compound.

Example 359C

4-[4-Bromobenzene sulfonyloxy-2,3-bis(trifluoromethyl)]benzylbromide

The phenol compound of Example 359B (10 g, 0.04 mol) was treated with 4-
10 bromobenzenesulfonyl chloride (11.0 g, 0.043 mol) and Hunig's base (5.56 g, 0.043 mol) in CH_2Cl_2 (150 ml). The solution was washed with water, brine and dried over MgSO_4 . After evaporating the solvent, N-bromosuccinimide (7.3 g, 0.04 mol) and benzoyl peroxide (200 mg) were added and the mixture was suspended in CCl_4 (100ml). The resulting mixture was refluxed for 13 hr. When the reaction was
15 cooled, the white solid was filtered and washed with CCl_4 to afford the crude title compound. This crude product was used for next step without further purification.

Example 359D

4-Hydroxy-2,3-bis(trifluoromethyl)benzaldehyde

20 The crude product of Example 359C was dissolved in 60 ml of DMSO and 20 ml of CH_2Cl_2 , and 12 g of trimethylamine N-oxide added. The resulting mixture was stirred at rt for 2.5 hr. The reaction mixture was poured into an ice cold 50% saturated

aqueous NaCl solution (200 ml) and extracted with ether (3X100 ml). The combined organic layer was washed with brine and dried over Na₂SO₄. After evaporation of solvent, the product was purified by column chromatography, eluted with hexane:EtOAc (3:2) to provide 3.0 g of the title compound, plus 4.0 g of recovered 4-

5 [4-bromobenzenesulfonyloxy-2,3-bis(trifluoromethyl)]toluene.

Example 359E

(Benzodioxan-6-yl)-[2,3-bis(trifluoromethyl)-4-(E-carboethenyl)phenyl]sulfide

The title compound was prepared according to the procedures described in

10 Example 330, substituting the compound of Example 359D for 4-hydrox-2,3-dichlorobenzaldehyde.

Example 359F


(Benzodioxan-6yl)[2,3-bis(trifluoromethyl)-4-(E-((4-carboxypiperidin-1-yl)carbonyl)ethenyl)phenyl]sulfide

15

The title compound was prepared from Example 359E by the procedures described in Example 330, giving a white solid. ¹H NMR (CD₃OD, 300MHz) δ 1.65(br s, 2H), 1.93-2.04 (m, 2H), 2.57-2.65 (m, 1H), 2.95-3.05 (m, 1H), 3.25 (m, 1H), 4.12 (m, 1H), 4.28 (m, 4H), 4.41 (m, 1H), 6.92-7.03 (m, 4H), 7.25 (d, J=9Hz, 1H), 7.72 (d, J=9Hz, 1H), 7.72-7.81 (m, 1H). MS (ESI) *m/e* 562 (M+H)⁺. Anal calcd for C₂₅H₂₁NO₃F₆S: C, 53.48; H, 3.77; N, 2.49. Found: C, 53.42; H, 3.69; N, 2.25.


20

307

Example 360

(2-Methoxyphenyl) [2,3-dichloro-4-(E-((4-(carboxymethylamino)carbonyl)-piperidin-1-yl)carbonyl)ethenyl]phenyl sulfide

5

Example 360A


(2-Methoxyphenyl)[2,3-dichloro-4-(E-((4-(methylaminomethylcarboxylate)carbonyl)-piperidin-1-yl)carbonyl)ethenyl]phenyl sulfide

10

The title compound was prepared by the procedure described in Example 363 using glycine methyl ester as the coupling substrate. HPLC (Supelco C-18 column, water:acetonitrile 50:90- 90:50, 9 minute elution, flow rate 1.5 mL/min, $t_r = 6.11$ min. MS (APCI) m/e 537 ($M+H$)⁺;

15

¹H NMR (300 MHz, DMSO- d_6) δ 1.46(m, 3H), 1.78(br d, 2H), 2.79(m, 1H), 3.15(m, 1H), 3.62(s, 3H), 3.80(s, 3H), 3.83(d, 2H), 4.20(m, 1H), 4.40(m, 1H), 6.58(d, 1H), 7.09(t, 1H), 7.22(d, 1H), 7.25(dd, 1H), 7.48(d, 1H), 7.56(t, 1H), 7.72(d, 1H), 7.81(d, 1H), 8.28(t, 1H). Anal calcd for $C_{25}H_{26}Cl_2N_2O_5S \cdot 1.3 H_2O$: C, 53.54; H, 5.14; N, 4.99. Found: C, 53.49; H, 4.88; N, 4.75.



Example 360B

(2-Methoxyphenyl) [2,3-dichloro-4-(E-((4-(carboxymethylamino)carbonyl)piperidin-1-yl)carbonyl)ethenyl]phenyl sulfide

- 5 The title compound was hydrolyzed as described in Example 340H. HPLC (Supelco C-18 column, water:acetonitrile 90:0- 0:90, 30 minute elution, flow rate 0.8 mL/min) rt 26.14 min. ¹H NMR (300 MHz, DMSO-d₆) δ 1.46 (m, 2H), 1.75 (m, 2H), 2.73 (m, 1H), 3.12 (m, 1H), 3.70 (m, 2H), 3.79 (s, 3H), 4.02 (m, 1H), 4.20 (m, 1H), 4.41 (m, 1H), 6.65 (d, 1H), 7.09 (dt, 1H), 7.22 (d, 1H), 7.25 (dd, 1H), 7.48 (dd, 1H), 7.58 (m, 1H), 7.72 (d, 1H), 7.82 (d, 1H), 8.11 (m, 1H). MS (APCI) *m/e* 523 (M+H)⁺.

Example 361

(2-Methoxyphenyl) [2,3-bis(trifluoromethyl)-4-(E-((4-carboxymethyl)piperazin-1-yl)carbonyl)ethenyl]phenyl sulfide

- 15 The title compound was prepared according to the procedures of Example 22, employing the compound of Example 359D as starting material, to give a white solid. ¹H NMR (CD₃OD, 300 MHz) δ 3.07-3.12 (m, 4H), 3.48 (s, 2H), 3.74 (s, 3H), 3.89 (br s, 4H), 6.99-7.18 (m, 4H), 7.53 (d, J=9Hz, 2H), 7.72 (d, J=9Hz, 1H), 7.78-7.88 (m, 1H). MS (ESI) *m/z* 549 (M+H)⁺. Anal calcd for C₂₆H₂₆F₆N₂O₄S·0.9HAc: C, 51.43, H, 4.28, N, 4.65. Found: C, 51.48, H, 4.12, N, 4.45.

Example 362

(2-Methoxyphenyl) [2,3-bis(trifluoromethyl)-4-(E-((4-N-(2-hydroxyethyl)piperazin-1-yl)carbonyl)ethenyl)phenyl]sulfide

The title compound was prepared by the procedures described in Example 356,

- 5 employing the compound of Example 359D as starting material to give an oil. ¹H NMR (CDCl₃, 300 MHz) δ 2.68 (br s, 6H), 3.71 (br s, 4H), 3.80 (br s, 5H), 6.55 (d, J=15Hz, 1H), 6.93-7.02 (m, 2H), 7.10 (d, J=9Hz, 1H), 7.35 (d, J=9Hz, 1H), 7.41-7.50 (m, 2H), 7.82 (qq, J=15Hz, 1H). MS (ESI) *m/z* 535 (M+H)⁺. Anal calcd for C₂₄H₂₄F₆N₂O₃SHCl: C, 50.49; H, 4.41; N, 4.91. Found: C, 50.72; H, 4.70; N, 4.55.

10

Example 363


(1-Methylindol-5-yl) [2,3-dichloro-4-(E-((4-(carbo-2,3-dihydroxypropylamino)piperidin-1-yl)carbonyl)ethenyl)phenyl] sulfide

To a solution of Example 340H (100mg, 0.2 mmol) and 3-amino-1,2-

- 15 propanediol (37.4 mg, 0.41 mmol) in DMF (3 mL) was added EDC (78 mg, 0.41 mmol), HOBT (55 mg, 0.41 mmol) and triethylamine (0.057 mL, 0.41 mmol). The reaction mixture was stirred at room temperature for 15 hours. Ethyl acetate (60 mL) was added, and the mixture was washed with brine. The aqueous phase was extracted with 10% MeOH in methylene chloride. The combined organic phases were
- 20 concentrated to dry. The residual material was triturated with water, filtered, washed with water, acetonitrile and ethyl acetate, and dried to give example 363 (92 mg, 80%). ¹H NMR (300 MHz, DMSO-*d*₆) δ 1.44 (m, 1H), 1.72 (m, 1H), 2.41 (m, 1H),

2.70 (t, 1H), 3.00 (m, 2H), 3.20 (m, 2H), 3.27 (m, 2H), 3.50 (m, 2H), 3.90 (s, 3H), 4.18 (br d, 1H), 4.40 (br d, 1H), 4.50 (t, 1H), 4.77 (d, 1H), 6.40 (d, 1H), 6.58 (d, 1H), 7.19 (d, 1H), 7.35 (d, 1H), 7.50 (d, 1H), 7.66 (d, 1H), 7.70 (m, 2H), 7.80 (t, 1H), 7.88 (s, 1H). MS (ESI⁺) m/z 562 (M+H)⁺. Anal. calcd for C₂₇H₂₉Cl₂N₃SO₄·0.25H₂O:


5 C, 57.19; H, 5.24; N, 7.41. Found: C, 57.07; H, 5.22; N, 7.13.



Example 364

10

(2-Methoxyphenyl) [2,3-dichloro-4-(E-(2,3-dihydroxypropionyl)piperazin-1-yl)carbonyl]ethenyl]phenyl] sulfide




Example 364A

15

(2-Methoxyphenyl) [2,3-dichloro-4-(E-((piperazin-1-yl)carbonyl)ethenyl)phenyl] sulfide


The title compound was prepared by the procedures described in Example 340G substituting methyl isonipecotate with piperazine. MS (DCI/NH₃) m/e 423 (M+H)⁺.

Example 364B

(2-Methoxyphenyl) [2,3-dichloro-4-(E-4-(2,3-dihydroxypropionyl)piperazin-1-yl)carbonyl]ethenyl]phenyl] sulfide


5 The title compound was prepared by the procedures described in Example 340, substituting methyl isonipecotate with Example 364A and substituting Example 340G with DL-glycric acid Ca salt. ^1H NMR (300 MHz, DMSO- d_6) δ 3.2-3.8 (m, 12H), 4.38 (t, 1H), 6.58 (d, 1H), 7.10 (t, 1H), 7.27 (d, 1H), 7.28 (d, 1H), 7.50 (d, 1H), 7.60 (t, 1H), 7.79 (d, 1H), 7.83 (d, 1H). MS (ESI $^+$) m/z 511 (M+H) $^+$.

10

Example 365


(2-Methoxyphenyl) [2,3-dichloro-4-(E-4-(2,3-dihydroxy-3-carboxypropionyl)piperazin-1-yl)carbonyl]ethenyl]phenyl] sulfide

15 The title compound was prepared by the procedures described in Example 340 substituting methyl isonipecotate with Example 364A and substituting Example 340G with meso-tartaric acid. ^1H NMR (300 MHz, CDCl $_3$) δ 3.70 (m, 8H), 4.33 (br s, 1H), 4.72 (br s, 1H), 6.58 (d, 1H), 6.77 (d, 1H), 7.03 (m, 2H), 7.25(d, 1H), 7.50 (d, 1H), 7.52 (d, 1H), 8.00 (d, 1H). MS (ESI $^+$) m/z 555 (M+H) $^+$.

**Example 366**


(1-Methylindol-5-yl) [2,3-dichloro-4-(E-((4-(carboxymethylamino)carbonyl)-
piperidin-1-yl)carbonyl)ethenyl]phenyl] sulfide

The title compound was prepared by the procedures described in Example 363 substituting 3-amino-1,2-propanediol with glycine methyl ester hydrochloride followed by hydrolysis. ¹H NMR (300 MHz, DMSO-d₆) δ 1.42 (m, 2H), 1.75 (m, 2H), 2.45 (m, 1H), 2.78 (m, 1H), 3.10 (m, 1H), 3.72 (d, 2H), 3.90 (s, 3H), 4.18 (br d, 1H), 4.40 (br d, 1H), 6.42 (d, 1H), 6.57 (d, 1H), 7.18 (d, 1H), 7.32 (d, 1H), 7.50 (d, 1H), 7.65 (d, 1H), 7.67 (d, 1H), 7.70 (m, 1H), 7.88 (s, 1H), 8.18 (t, 1H). MS (ESI⁺) *m/z* 546 (M+H)⁺. Anal. calcd for C₂₆H₂₅N₃Cl₂SO₄: C, 57.15; H, 4.61; N, 7.69. Found: C, 57.17; H, 4.64; N, 7.39.

**Example 367**

(1-Methylindol-5-yl) [2,3-dichloro-4-(E-((4-sulfopiperidin-1-
yl)carbonyl)ethenyl]phenyl] sulfide


The title compound was prepared from Example 340F, by the procedures described in Example 340G, substituting methyl isonipecotate with piperidine-4-sulfonic acid. ^1H NMR (300 MHz, DMSO- d_6) δ 1.40 (m, 2H), 1.90 (m, 2H), 3.03 (m, 1H), 4.10 (m, 3H), 4.42 (br d, 1H), 6.40 (d, $J=8.8\text{Hz}$, 1H), 6.53 (d, $J=3.1\text{Hz}$, 1H), 7.15 (d, $J=15.3\text{Hz}$, 1H), 7.33 (dd, $J=8.5$, 1.7Hz, 1H), 7.48 (d, $J=3.1\text{Hz}$, 1H), 7.65 (d, $J=8.5\text{Hz}$, 1H), 7.67 (d, $J=15.2\text{Hz}$, 1H), 7.74 (d, $J=8.8\text{Hz}$, 1H), 7.87 (d, $J=1.5\text{Hz}$). MS (ESI $^+$) m/z 525 (M+H) $^+$. Anal. calcd for C₂₃H₂₂N₂Cl₂S₂O₄·0.8 TFA: C, 47.91; H, 3.73; N, 4.54. Found: C, 47.71; H, 3.84; N, 4.73.



Example 368

(1-Methylindol-5-yl) [2,3-dichloro-4-(E-(4-methylhomopiperazin-1-ylcarbonyl)ethenyl)phenyl] sulfide

The title compound was prepared by the procedures described in Example 340G substituting methyl isonipecotate with N-methyl homopiperazine. ^1H NMR (300 MHz, DMSO- d_6) δ 2.06 (m, 2H), 2.81 (m, 2H), 3.17 (m, 2H), 3.55 (m 3H), 3.70 (s, 3H), 3.86 (s, 3H), 4.05 (m, 1H), 6.42 (dd, $J=8.4$, 3.3Hz, 1H), 6.54 (d, $J=3.0\text{Hz}$, 1H), 7.08 (dd, $J=15.4$, 7.5Hz, 1H), 7.35 (dd, $J=8.8$, 2.0Hz, 1H), 7.49 (d, $J=3.0\text{Hz}$, 1H), 7.65 (d, $J=8.5\text{Hz}$, 1H), 7.73 (d, $J=8.8\text{Hz}$, 1H), 7.80 (d, $J=15.2\text{Hz}$, 1H), 7.88 (d, $J=2.0\text{Hz}$, 1H). MS (ESI $^+$) m/z 474 (M+H) $^+$. Anal. calcd for C₂₆H₂₆N₃Cl₂SF₃O₃·0.75 TFA: C, 49.01; H, 4.00; N, 6.23. Found: C, 48.71; H, 4.09; N, 6.13.

**Example 369**

(1-Methylindol-5-yl) [2,3-dichloro-4-(E-(4-tetrahydrofuroyl)piperazin-1-yl)carbonyl)ethenyl)phenyl] sulfide

5

The title compound was prepared by the procedures described in Example

340G substituting methyl isonipecotate with 1-tetrahydrofuroylpiperazine. ¹H NMR


(300 MHz, DMSO-d₆) δ 1.80 (m, 2H), 2.00 (m, 2H), 3.50 (m, 8H), 3.75 (m, 2H), 3.88

(s, 3H), 4.68 (t, 1H), 6.42 (d, 1H), 6.57 (d, 1H), 7.19 (d, 1H), 7.32 (d, 1H), 7.48 (d,

10 1H), 7.65 (d, 1H), 7.70 (d, 1H), 7.75 (d, 1H), 7.87 (s, 1H). MS (ESI⁺) *m/z* 544

(M+H)⁺. Anal calcd for C₂₇H₂₇N₃Cl₂SO₃: C, 59.56; H, 4.99; N, 7.71. Found: C,

59.40; H, 4.94; N, 7.61.



15

Example 370

(Benzodioxan-6-yl) [2-(benzodioxan-6-thioxy)-4-(E-((4-morpholino)carbonyl)ethenyl)phenyl] sulfide

Example 370A(E)-Morpholino 2,4-difluorocinnamide


The title compound was processed as reported in Example 1C substituting morpholine (1.04 mL, 11.9 mmol) for the amine and *trans*-2,4-difluorocinnamic acid (1.00 g, 5.4 mmol) for the carboxylic acid. The title compound was obtained as an off-white foam (1.4 g, 100%). ¹H NMR (DMSO-d₆, 300 MHz) δ 8.04 (dd, J=15.26, 8.82 Hz, 1H), 7.53 (d, J=14.91 Hz, 1H), 7.38-7.30 (m, 1H), 3.61-3.48 (m, 8H). MS (APCI) *m/z* 254 (M+H)⁺.

10

Example 370BMorpholinyl-(E)-2,4-bis(1,4-benzodioxane-6-mercaptan)cinnamic amide


Example 370A (233 mg, 1.00 mmol) was combined with cesium carbonate (652 mg, 2.00 mmol), 1,4-benzodioxane-6-thiol (370 mg, 2.20 mmol), and DMF (5 mL). The mixture was processed as reported in Example 1A to provide the title compound (220 mg, 40%) as a white foam. ¹H NMR (DMSO-d₆, 300 MHz) δ 7.83 (d, J=15.20 Hz, 1H), 7.80 (d, J=8.20 Hz, 1H), 7.17 (d, J=15.3 Hz, 1H), 7.02 (dd, J=8.5, 2.0 Hz, 1H), 6.87-6.75 (m, 6H), 6.48 (s, 1H), 4.33-4.25 (m, 8H), 3.61-3.48 (m, 8H). MS (APCI) *m/z* 550 (M+H)⁺.

20

Example 371

(2-Methoxyphenyl) [2,3-dichloro-4-(E-((4-amino-4-carboxypiperidin-1-yl)(carbonyl)ethenyl)phenyl)] sulfide

To a suspension of Example 355 (700 mg, 1.4 mmol) in DME (10 mL) was added a solution of (BOC)₂O (1.51 g, 6.9 mmol) in DME (5 mL), triethylamine (0.23 mL, 1.7 mmol) and DMAP (9 mg, 0.07 mmol). The reaction mixture was stirred at room temperature overnight. Additional triethylamine (0.23 mL) and DMAP (30 mg) were added, and the mixture was heated at 60 °C for 6 hours. After aqueous work up, the crude product was suspended in DME (5 mL) and water (5 mL) containing 200 mg of NaOH. The suspension was stirred for 5 hours at room temperature, and separated by HPLC to give the title compound (300 mg, 45%). ¹H NMR (300 MHz, DMSO-d₆) δ 1.78 (m, 2H), 2.10 (m, 2H), 3.60 (m, 2H), 3.80 (s, 3H), 3.86 (m, 2H), 6.58 (d, 1H), 7.10 (d, 1H), 7.25 (d, 1H), 7.28 (d, 1H), 7.50 (d, 1H), 7.58 (t, 1H), 7.77 (d, 1H), 7.80 (d, 1H), 8.50 (br s, 2H). MS (ESI⁺) *m/z* 481 (M+H)⁺. Anal calcd for C₂₂H₂₂N₂Cl₂SO₄·0.75 H₂O: C, 47.34; H, 4.06; N, 4.60. Found: C, 47.31; H, 4.05; N, 4.43.




Example 372

(2-Methoxyphenyl)[2,3-dichloro-4-((4-furoyl)piperazin-1-yl)(carbonyl)ethenyl]phenyl] sulfide

To a solution of Example 364A (100mg, 0.24 mmol) and 2-furfural (30mg, 0.24 mmol) in dichloroethane (2 mL) was added $\text{NaBH}(\text{OAc})_3$ (142 mg, 0.67 mmol) under nitrogen atmosphere. The mixture was stirred for 16 hours at room temperature.

Dichloromethane (20 mL) was added and the mixture was washed with 5% NaHCO_3 ,

- 5 then with brine, and the organic phase was separated and concentrated. The residual solid was chromatographed by flash chromatography (5% $\text{MeOH}/\text{CH}_2\text{Cl}_2$) and desired fractions were combined, concentrated and dried to afford the title compound as an off-white solid (84 mg, 69%). HPLC (Supelco C-18 column, water:acetonitrile 100:0- 0:100, 15 minute elution, flow rate 1.5 mL/min) rt 11.90 min. ^1H NMR (300
- 10 MHz, $\text{DMSO}-d_6$) δ 2.39 (m, 4H), 3.52 (s, 2H), 3.55 (m, 2H), 3.63 (m, 2H), 3.79 (s, 3H), 6.29 (d, 1H), 6.40 (m, 1H), 6.57 (d, 1H), 7.08 (dt, 1H), 7.21 (d, 1H), 7.23 (dd, 1H), 7.48 (dd, 1H), 7.57 (m, 2H), 7.72 (d, 1H), 7.80 (d, 1H). MS (ESI) m/e 503 ($\text{M}+\text{H}$) $^+$.



15

Example 373


(1-Methylindol-5-yl) [2,3-dichloro-4-(E-(4-(carbo-3-sulfopropylamino)piperidin-1-yl)carbonyl)ethenyl]phenyl sulfide

- The title compound was prepared from Example 340H by the procedures
- 20 described in Example 363 substituting 3-amino-1,2-propanediol with 3-amino-1-propanesulfonic acid. ^1H NMR (300 MHz, $\text{DMSO}-d_6$) δ 1.40 (m, 2H), 1.70 (m, 4H),

2.38 (m, 1H), 2.42 (m, 2H), 2.70 (m, 1H), 3.05 (m, 3H), 3.86 (s, 3H), 4.18 (br d, 1H),
 4.40 (br d, 1H), 6.40 (d, 1H), 6.55 (d, 1H), 7.20 (d, 1H), 7.35 (d, 1H), 7.50 (d, 1H),
 7.65 (d, 1H), 7.70 (d, 1H), 7.77 (d, 1H), 7.87 (d, 1H). MS (ESI⁺) m/z 610 (M+H)⁺.

Anal calcd for C₂₇H₂₉N₃Cl₂S₂O₅·1.5 TFA: C, 46.10; H, 3.93; N, 5.38. Found: C,

5 46.52; H, 4.03; N, 5.66.



Example 374

(2-Methoxyphenyl)[2,3-dichloro-4-(E-(4-acetylamino-4-carboxypiperidin-1-yl)carbonyl)ethenyl]phenyl sulfide

10

To a suspension of Example 371 (90 mg, 0.187 mmol) and triethylamine (0.08 mL, 0.57 mmol) in DMF (3 mL) was added acetyl chloride (0.1 mL) at room temperature. The mixture was stirred for 3 hours. Ethyl acetate (60 mL) was added, and the mixture was washed with brine. The organic phase was dried, filtered and concentrated. The residue was separated by HPLC (C-18, CH₃CN/H₂O) to give example 374 (56 mg, 57%).

15

¹H NMR (300 MHz, DMSO-d₆) δ 1.78 (m, 2H), 1.82 (s, 3H), 1.98 (m, 2H), 3.05 (t, 1H), 3.38 (t, 1H), 3.80 (s, 3H), 4.00 (br d, 1H), 4.12 (br d, 1H), 6.58 (d, 1H), 7.08 (t, 1H), 7.23 (d, 1H), 7.25 (d, 1H), 7.50 (d, 1H), 7.58 (t, 1H), 7.78 (d, 1H), 7.80 (d, 1H),
 20 8.18 (s, 1H). MS (ESI⁺) m/z 523 (M+H)⁺. Anal calcd for C₂₄H₂₄N₂Cl₂SO₅·0.35TFA:

C, 52.80; H, 4.40; N, 5.05. Found: C, 52.74; H, 4.42; N, 5.11.

Example 375

(2-Methoxyphenyl) [2,3-bis(trifluoromethyl)-4-(E-((4-carboxypiperidin-1-yl)carbonyl)ethenyl)phenyl]sulfide

5 The title compound was prepared by the procedures described in Example 352, employing the compound of Example 359D to give a white solid. ¹H NMR(CD₃OD, 300 MHz) δ 1.65 (br s, 2H), 1.94-2.03 (m, 2H), 2.57-2.67 (m, 1H), 2.95-3.05 (m, 1H), 3.23-3.32 (m, 1H), 3.75 (s, 3H), 4.12 (br s, 1H), 4.40 (br s, 1H), 7.00 (d, J=15Hz, 1H), 7.03-7.20 (m, 3H), 7.47-7.53 (m, 2H), 7.68 (d, J=9Hz, 1H), 7.77 (qq, J=15Hz, 1H). MS (ESI) *m/z* 534 (M+H)⁺. Anal calcd for C₂₄H₂₁NF₆O₄S: C, 54.03; H, 3.97; N, 2.63. Found: C, 54.11; H, 4.04; N, 1.76.

Example 376

(2-Methoxyphenyl) 5-[8-(E-((4-(aminocarbonyl)piperidin-1-yl)carbonyl)ethenyl)quinolinyl]sulfide

15

Example 376A

5-Chloro-8-(trifluoromethanesulfonyloxy)quinoline

5-Chloro-8-hydroxyquinoline was treated as described in Example 340E to provide the title compound. ¹H NMR (DMSO-d₆, 300 MHz) δ 7.59 (7.5Hz, 1H), 7.65-7.69 (m, 2H), 8.63 (dd, J₁=9Hz, J₂=1.5Hz, 1H), 9.21 (dd, J₁=6Hz, J₂=1.5Hz, 1H). MS (APCI-NH₃) *m/e* 312, 314 (M+H)⁺.

Example 376B5-Chloro-8-[E-(methoxycarbony)ethenyl]quinoline

The method of Example 340D was used, substituting the product from Example 376A for Example 340C. Thus, Example 376A (6.23 g, 20.0 mmol) was converted to the title compound (2.22 g, 45%). ¹H NMR (DMSO-d₆, 300 MHz) δ 3.78 (s, 3H), 6.98 (d, J=16.5Hz, 1H), 7.78-7.83 (m, 1H), 7.88 (d, J=9Hz, 1H), 8.32 (d, J=9Hz, 1H), 8.65 (dd, J₁=9Hz, J₂=1.5Hz, 1H), 8.85 (d, J=16.5 Hz, 1H), 9.12 (dd, J₁=4.5Hz, J₂=1.5Hz, 1H). MS (APCI-NH₃) *m/e* 248, 250 (M+H)⁺.

10

Example 376C(2-Methoxyphenyl) 5-[8-(E-(methoxycarbonyl)ethenyl)quinolinyl]sulfide

The method of Example 340F was used, substituting the product from Example 376B for Example 340E. Thus, Example 376B (2.19 g, 8.84 mmol) was converted to the title compound (1.07 g, 36%). ¹H NMR (DMSO-d₆, 300 MHz) δ 3.83 (s, 3H), 6.80 (d, J=16.5Hz, 1H), 6.86-6.99 (m, 2H), 7.16 (d, J=6Hz, 1H), 7.33-7.38 (m, 1H), 7.44 (d, J=7.5Hz, 1H), 7.67-7.72 (m, 1H), 8.22 (d, J=7.5Hz, 1H), 8.63 (dd, J₁=9Hz, J₂=1.5Hz, 1H), 8.82 (d, J=16.5Hz, 1H), 9.07 (dd, J₁=6Hz, J₂=1.5Hz), 12.48 (s, 1H). MS (APCI-NH₃) *m/e* 338 (M+H)⁺.

20

Example 376C(2-Methoxyphenyl) 5-[8-(E-((4-aminocarbonyl)piperidin-1-yl)carbonyl)ethenyl]quinolinyl]sulfide

The method of Example 340G was used, substituting the product from Example 376B for Example 340F, and substituting 4-piperidinecarboxamide for methyl


isonipecotate. ^1H NMR ($\text{DMSO}-d_6$, 300 MHz) δ 1.71-2.82 (m, 2H), 2.96-2.03 (m, 2H), 2.44-2.52 (m, 1H), 2.81-2.94 and 3.16-3.30 (m, 1H), 3.37-3.54 (m, 2H), 3.88 (s,

- 5 3H), 4.17-4.34 and 4.60-4.80 (m, 1H), 5.72 (s, 2H), 6.82 (t, 4.5Hz, 1H), 6.90 (dd, $J_1=4.5\text{Hz}$, $J_2=0.75\text{Hz}$, 1H), 6.93 (d, 6Hz, 1H), 7.23-7.28 (m, 1H), 7.40 (d, $J=9\text{Hz}$, 1H), 7.47-7.50 (m, 1H), 7.51 (d, $J=6\text{Hz}$, 1H), 7.82 (d, $J=4.5\text{ Hz}$, 1H), 8.57 (d, $J=9\text{Hz}$, 1H), 8.74 (dd, $J_1=4.5\text{Hz}$, $J_2=0.75\text{Hz}$, 1H), 9.00 (m, 1H).

10

Example 377

(2-Methoxyphenyl) [2-trifluoromethyl-4-(E-((4-carboxypiperidin-1-yl)carbonyl)ethenyl)phenyl]sulfide



15

Example 377A

2-Trifluoromethyl-4-(thiobenzodioxan-6-yl)cinnamic acid


- A solution of commercially available 4-fluoro-2-(trifluoromethyl)cinnamic acid (5 g, 21.4 mmol) in ethyl acetate (200 mL) under nitrogen at ambient temperature was treated with a solution of diazomethane in diethyl ether to a persistent yellow
- 20 color, stirred an additional ten minutes, then quenched by dropwise addition of glacial acetic acid. The resultant clear solution was washed with saturated NaHCO_3 ,

brine, dried (MgSO₄), filtered through a plug of silica, rinsed with ethyl acetate and concentrated to give 5.4 grams of a yellow oil. A solution of this methyl ester (2.5 g, 10 mmol) and 6-mercaptobenzodioxane (1.9 g, 11 mmol) in 40 mL of dimethylformamide was treated with cesium carbonate (3.9 g, 12 mmol), and stirred

5 at room temperature for 20 hours. The resultant orange heterogeneous solution was diluted with diethyl ether and water, washed with 1 M NaOH, distilled water, brine, dried (MgSO₄), filtered through a plug of silica, concentrated and then flash chromatographed with 20% ethyl acetate/hexane followed by 33% ethyl acetate/hexane to give 2.8 g of a light yellow syrup. A solution of this diaryl sulfide


10 ester (2.8 g, 7.1 mmol) in THF (21 mL) and distilled water (7 mL) was treated with lithium hydroxide hydrate (450 mg, 10.7 mmol) and stirred 67 hours at ambient temperature. The resultant solution was diluted with distilled water, washed with diethyl ether, acidified to pH 1-2 with 3 M H₂SO₄, extracted with diethyl ether, washed with brine, dried (MgSO₄) and concentrated to give 2.7 g (7.1 mmol) of the

15 title compound as an off-white powder (71%). ¹H NMR (300 MHz, d₆-DMSO) δ 7.97 (d, 1H), 7.72 (dq, 1H), 7.47 (d, 1H), 7.31 (dd, 1H), 7.05 (m, 3H), 6.58 (d, 1H), 4.3 (m, 4H). MS (APCI-NH₃) *m/e* 383 (M+H)⁺, 400 (M+NH₄)⁺.

Example 377B

(Benzodioxan-6-yl)[3-(trifluoromethyl-4-(E-((2-carboxypiperidin-1-yl)carbonyl)ethenyl)phenyl]sulfide

Example 377A (382 mg, 1 mmol) was coupled with (d,l)-ethyl picipolate according to the procedure of Example 340G. The derived ethyl ester was hydrolyzed using the method of Example 340H to give 280 mg of the title compound as a light yellow foam (84%). Analytical HPLC: 4.6X250 mm C18 column, 0.8 mL/min, 254 nm, CH₃CN:H₂O with 0.1% TFA, 0:100 (0 min), ramp to 90:10 (0-10 min), 90:10 (10-18 min), ramp to 0:100 (18-20 min), rt 11.29 min (98.2 area%). ¹H NMR (300 MHz, d₆-DMSO) δ 8.07 (t, 1H), 7.65 (dq, 1H), 7.38 (m, 3H), 7.03 (m, 3H), 5.15 (m, 1H), 4.4 (m, 1H), 4.29 (m, 4H), 4.1 (m, 1H), 3.2 (m, 1H), 2.2 (m, 1H), 1.68 (m, 2H), 1.3 (m, 2H). MS (APCI-NH₃) *m/e* 494 (M+H)⁺, 511 (M+NH₄)⁺.




Example 378

(1-Methylindol-5-yl) [2,3-dichloro-4-(E-(((1S,4S)-5-tert-butyloxycarbonyl-2,5-diazabicyclo(2,2,1)heptan-2-yl)carbonyl)ethenyl)phenyl] sulfide

The title compound was prepared by the procedures described in Example 340 substituting methyl isonipicotate with t-butyl (1S,4S)-(-)-2,5-diazabicyclo(2,2,1)heptane-2-carboxylate. ¹H NMR (300 MHz, DMSO-d₆) δ 1.40 (s, 9H), 1.82 (m, 2H), 3.17 (m, 1H), 3.30 (m, 2H), 3.58 (m, 1H), 3.82 (s, 3H), 4.05 (m,

1H), 4.40 (m, 1H), 4.75 (br s, 1H), 4.92 (br s, 1H), 6.42 (dd, 1H), 6.58 (d, 1H), 6.75 (d, 1H), 7.05 (d, 1H), 7.35 (d, 1H), 7.50 (d, 1H), 7.65 (d, 1H), 7.68 (d, 1H), 7.78 (t, 1H), 7.77 (s, 1H). MS (ESI⁺) *m/z* 558 (M+H)⁺. Anal calcd for C₂₈H₂₉N₃Cl₂SO₃: C, 60.21; H, 5.23; N, 7.52. Found: C, 60.23; H, 5.36; N, 7.41.


5



Example 379

(1-Methylindol-5-yl) [2,3-dichloro-4-(E/Z-((1S,4S)-2,5-diazabicyclo(2.2.1)heptan-2-ylcarbonyl)ethenyl)-2,3-dichlorophenyl] sulfide

- 10 To a solution of Example 378 (820 mg, 1.47 mmol) in CH₂Cl₂ (20 mL) was added trifluoroacetic acid (2mL) at 0°C. The yellow solution was stirred at the same temperature for 2 hours. More CH₂Cl₂ (50 mL) was added and the solution was poured into water (100 mL) containing NaHCO₃ (4.5 g). The insoluble material was collected by filtration, washed with water and methanol. The CH₂Cl₂ solution was
- 15 concentrated, and the residual solid was filtered, washed with water, methanol and CH₂Cl₂. The combined solid was dried to give the title compound (650 mg, 95%).
- ¹H NMR (300 MHz, DMSO-d₆) δ 1.70 (m, 2H), 2.90 (m, 1H), 3.50 (m, 4H), 3.88 (s, 3H), 4.85 (m, 1H), 6.45 (d, 1H), 6.60 (dd, 1H), 6.77 (d, 1H), 7.05 (dd, 1H), 7.25 (s, 1H), 7.35 (dd, 1H), 7.65 (d, 1H), 7.70 (d, 1H), 7.80 (d, 1H). MS (ESI⁺) *m/z*
- 20 458(M+H)⁺.

Example 380

(1-Methylindol-5-yl) [2,3-dichloro-4-(E-(4-hydroxy-3-carboxypiperidin-1-ylcarbonyl)ethenyl)phenyl] sulfide


5 To a suspension of Example 340G (300 mg, 0.794 mmol) and methyl 4-oxo-3-piperidine carboxylate hydrochloride (307 mg, 1.59 mmol) in DMF (10 mL) was added EDC (305 mg, 1.59 mmol), HOBt (215 mg, 1.59 mmol) and triethylamine (0.443 mL, 1.59 mmol). The suspension was stirred at room temperature overnight. Ethyl acetate (100 mL) was added and the mixture was washed with brine, water and

10 was concentrated. The residual oil was separated by flash chromatography (60% EtOAc in hexane) to give a white solid (220 mg).

180 mg of this solid was dissolved in THF (10 mL). A solution of lithium hydroxide monohydrate (29 mg, 0.68 mmol) in water (10 mL) was added. The mixture was stirred at room temperature 2 hours, NaBH₄ (50 mg) was then added.

15 After 4 hours stirring, the solution was acidified and concentrated to 5 mL. The formed white solid was collected by filtration, washed with water, acetonitrile, and dried to give the title compound (92 mg). ¹H NMR (300 MHz, DMSO-d₆) δ 1.60 (m, 2H), 3.00 (m, 1H), 3.40 (m, 1H), 3.85(1H, 4.05 (m, 1H), 4.20 (m, 1H), 4.35 (m, 1H), 5.00 (m, 1H), 6.42 (d, 1H), 6.58 (d, 1H), 7.20 (dd, 1H), 7.35 (d, 1H), 7.50 (d, 1H),

20 7.6-7.8 (m, 3H), 7.90 (s, 1H). MS (ESI⁺) *m/z* 505 (M+H)⁺. Anal calcd for C₂₄H₂₂N₂Cl₂SO₄: C, 57.03; H, 4.38; N, 5.54. Found: C, 56.77; H, 4.17; N, 5.34.

Example 381


(1-Methylindol-5-yl) [2,3-dichloro-4-(E-(S-oxothiomorpholin-1-

5

ylcarbonyl)ethenyl)phenyl] sulfide

The title compound was prepared by the procedures described in Example 340 substituting methyl isonipecotate with thiomorpholine S-oxide. ¹H NMR (300 MHz, CDCl₃) δ 2.70 (m, 2H), 2.85 (m, 2H), 3.85 (s, 3H), 3.90 (m, 2H), 4.20 (m, 1H), 4.60 (m, 1H), 6.45 (d, 1H), 6.55 (d, 1H), 6.70 (d, 1H), 7.18 (d, 1H), 7.20 (d, 1H), 7.38 (d, 1H), 7.41 (d, 1H), 7.77 (s, 1H), 7.98 (d, 1H). MS (ESI⁺) *m/z* 479 (M+H)⁺.

10

Example 382

(2-Methoxyphenyl) [2,3-dichloro-4-(E-(4-

15


sulfophenylamino)carbonyl)ethenyl)phenyl] sulfide

The title compound was prepared by the procedures described in Example 1C substituting Example 1B with (2-methoxy) [2,3-dichloro-4-(E-(2-carboxyethenyl)phenyl] sulfide and substituting 6-amino-1-hexanol with sulfanilic acid. ¹H NMR (300 MHz, DMSO-d₆) δ 3.82 (s, 3H), 6.65 (d, 1H), 6.82 (d, 1H), 7.12

(t, 1H), 7.25 (d, 1H), 7.5-7.7 (m, 7H), 7.85 (d, 1H), 10.40 (s, 1H). MS (ESI⁺) *m/z* 510

(M+H)⁺. Anal calcd for C₂₂H₁₇Cl₂NS₂O₅·0.65TFA: C, 50.80; H, 3.25; N, 2.55.

Found: C, 50.75; H, 3.43; N, 2.65.




Example 383

(2-Methoxyphenyl) [2,3-dichloro-4-(E-((4-carboxyphenylamino)carbonyl)ethenyl)phenyl] sulfide

The title compound was prepared by the procedures described in Example 1C


substituting Example 1B with (2-methoxy) [2,3-dichloro-4-(E-(2-carboxyethenyl)phenyl] sulfide and substituting 6-amino-1-hexanol with 4-aminobenzoic acid. ¹H NMR (300 MHz, DMSO-d₆) δ 3.82 (s, 3H), 6.65 (d, 1H), 6.82 (d, 1H), 7.10 (t, 1H), 7.30 (d, 1H), 7.60 (m, 3H), 7.82 (t, 3H), 7.90 (d, 1H), 7.92 (d, 1H), 10.65 (s, 1H), 12.75 (s, 1H). MS (ESI⁺) *m/z* 474 (M+H)⁺.

328

Example 384

[3-(4-Morpholino)phenyl] [2,3-dichloro-4-(E-((4-carboxypiperidin-1-yl)carbonyl)ethenyl)phenyl] sulfide

5


Example 384A

10 (3-Bromophenyl) [2,3-dichloro-4-(E-[methoxycarbonyl]ethenyl)phenyl] sulfide

To a solution of the resultant compound from Example 340E (12.0 g, 31.7 mmol) in N-methylpyrrolidinone (63 mL) at 0 °C (under dry N₂) was added 3-bromothiophenol (4.0 mL, 7.3 g, 38.8 mmol) and a solution of lithium *tert*-butoxide (3.1 g, 38.8 mmol), and the resulting solution was stirred for 3 h at 0 °C. The reaction

15 was diluted with 500 mL EtOAc and extracted sequentially with 100 mL water, 3 x 60 mL of 1 N aq. NaOH, then 2 x 100 mL brine. The organic phase was dried over Na₂SO₄, filtered, and concentrated in vacuo to produce the crude title compound (9.2 g). ¹H NMR (DMSO-d₆, 300 MHz) δ 3.75 (s, 3H), 6.67 (d, J=15 Hz, 1H), 6.83 (d, J=9 Hz, 1H), 7.46-7.59 (m, 2H), 7.72-7.76 (m, 2H), 7.80 (t, J=2.5 Hz, 1H), 7.85 (d, J=9 Hz, 1H), 7.88 (d, J=15 Hz, 1H); MS (APCI) *m/e* 419 (M+H)⁺.


20



0054795-03400


Example 384B(3-Bromophenyl) [2,3-dichloro-4-(*E*-carboxyethenyl)phenyl] sulfide

Using the procedure for Example 340H, Example 348A was hydrolyzed to the title compound. ¹H NMR (DMSO-*d*₆, 300 MHz) δ 6.56 (d, J=16.5 Hz, 1H), 6.84 (d, J=9 Hz, 1H), 7.45-7.58 (m, 2H), 7.72 (m, 1H), 7.77-7.86 (m, 4H), 12.75 (br s, 1H); (ESI) *m/e* 401, 403 (M-H)⁺.

Example 384C

10 (3-bromophenyl) [2,3-dichloro-4-(*E*-[4-ethoxycarbonylpiperidin-1-yl]carbonyl]ethenyl)phenyl] sulfide

The title compound (750 mg, 58%) was prepared from Example 384B (1.0 g, 2.48 mmol), using the procedures described in Example 340G substituting methyl isonipecotate with ethyl isonipecotate. ¹H NMR (300 MHz, DMSO-*d*₆) δ 1.18 (t, J=7.5 Hz, 3H), 1.38-1.56 (m, 2H), 1.82-1.92 (m, 2H), 2.50-2.69 (m, 1H), 2.80-2.93 (m, 1H), 3.14-3.27 (m, 1H), 4.07 (t, J=7.5 Hz, 2H), 4.10-4.35 (m, 2H), 6.92 (d, J=9 Hz, 1H), 7.30 (d, J=15 Hz, 1H), 7.43-7.52 (m, 2H), 7.67-7.77 (m, 3H), 7.92 (d, J=9 Hz, 1H).




Example 384D

[3-(4-Morpholino)phenyl] [2,3-dichloro-4-(E-[(4-ethoxycarbonylpiperidin-1-yl)carbonyl]ethenyl)phenyl] sulfide


- 5 The procedure of D. W.; Wolfe, J. P.; Buchwald, S. L. *J. Am. Chem. Soc.* **1998**, *120*, 9722-9723, was adapted. To a stirred solution of Example 384D (180 mg, 0.331 mmol) in ethylene glycol dimethyl ether (1 mL) containing 1-(N,N-dimethylamino)-1'-(dicyclohexylphosphino)biphenyl (7 mg, 5 mol%), Pd₂(dba)₃ (8 mg, 2.5 mol%), and morpholine (0.058 mL, 0.663 mmol) was added powdered K₃PO₄
- 10 (141 mg, 0.663 mmol). The reaction mixture was bubbled with N₂ for 5 min and heated at 90 °C in sealed tube for 18 h. Then the solvent was removed under reduced pressure and residue was diluted with methylene chloride (1 mL). The title compound (90 mg, 50%) was isolated by flash chromatography on silica gel eluting with 20% acetone-hexane. ¹H NMR (DMSO-d₆, 300 MHz) δ 1.18 (t, J=7.5 Hz, 3H), 1.35-1.55
- 15 (m, 2H), 1.79-1.91 (m, 2H), 2.58-2.69 (m, 1H), 2.70-2.94 (m, 2H), 3.16 (t, J=4.5 Hz, 2H), 3.15 (t, J=5 Hz, 4H), 3.73 (t, J=4.5 Hz, 4H), 3.78 (t, J=5 Hz, 2H), 4.08 (q, J=7.5 Hz, 2H), 4.11-4.36 (m, 2H), 6.70 (d, J=8.25 Hz, 1H), 6.97 (m, 1H), 7.10-7.27 (m, 2H), 7.24 (d, J=15 Hz, 1H), 7.39 (m, 1H), 7.73 (d, J=15 Hz, 1H), 7.86 (d, J=8.25 Hz, 1H); MS (ESI) *m/e* 549, 551 (M+H)⁺.

331

Example 384E

[3-(4-Morpholino)phenyl] [2,3-dichloro-4-(E-[(4-carboxypiperidin-1-yl)carbonyl]ethenyl)phenyl] sulfide

- 5 The title compound (38 mg, 67%) was prepared from Example 384D (60 mg, 0.11mmol) using the procedures described in Example 340H. ¹H NMR (DMSO-*d*₆, 400 MHz) δ 1.37-1.55 (m, 2H), 1.81-1.90 (m, 2H), 2.52-2.58 (m, 1H), 2.80-2.94 (m, 1H), 3.10-3.15 (m, 3H), 3.67-3.75 (m, 3H), 3.76-3.99(m, 3H), 4.04-4.16 (M, 1H), 4.22-4.33 (m, 1H), 6.71 (d, J=8Hz, 1H), 6.96 (d, J=7Hz, 1H), 7.07 (m, 1H), 7.12 (s, 1H), 7.24 (d, J=15Hz, 1H), 7.38 (t, j=7Hz, 1H), 7.73 (d, J=15Hz, 1H), 7.85 (d, J=8Hz, 1H). MS (ESI) *m/e* 521, 523 (M+H)⁺, 519, 521 (M-H)⁻.

Example 385


- 15 (2-Methoxyphenyl)[2,3-bis(trifluoromethyl)-4-(E-((4-phenylcarboxypiperidin-1-yl)carbonyl]ethenyl)phenyl]sulfide



Example 385A4-Phenylpiperidine-4-carboxylic acid

4-Cyano-4-phenylpiperidine hydrochloride (2.0g, 0.11 mol) was dissolved in 8 mL of conc. H_2SO_4 and 4mL of H_2O , then the solution was heated at reflux for 4 h.

- 5 The solution was cooled and then NaOH was added to precipitate a white solid. The solid was collected, then dissolved in methanol, and the solution was filtered and concentrated to obtain a white solid. This dried solid was used for without purification for Example 385B.




10

Example 385BMethyl 4-phenylpiperidine-4-carboxylate hydrochloride

Dissolved the 4-phenylpiperidinecarboxylic acid in 10 mL of methanol and added 2 mL of thionyl chloride dropwise at room temperature. The resulting mixture

- 15 was stirred overnight at room temperature. Evaporated solvent in vacuo, added toluene and evaporated excess thionyl chloride in vacuo. This white powder salt was used for next step without further purification.




20

Example 385C

(2-Methoxyphenyl)[2,3-bis(trifluoromethyl)-4-(E-((4-phenylcarboxypiperidin-1-yl)carbonyl)ethenyl)phenyl]sulfide

The methyl ester of the title compound was prepared by the procedures described in Example 356, employing the compound of Example 359D as starting

- 5 material, to give an oil. The resultant methyl ester was hydrolyzed with aq. NaOH in methanol at 60 °C for 4 h to give a white solid. ¹H NMR (CD₃OD, 300 MHz) δ 1.88 (br t, J=13.5 Hz, 2H), 2.59(br d, J=13.5 Hz, 2H), 3.13(br t, J=13.5 Hz, 1H), 3.75 (s, 3H), 3.44 (br t, J=13.5 Hz, 1H), 4.12 (br d, J=13.5Hz, 1H), 4.42(br d, J=13.5 Hz, 1H), 6.35 (d, J=15 Hz, 1H), 7.0-7.46 (m, 7H), 7.43-7.55 (m, 3H), 7.62-7.85 (m, 2H);
- 10 MS(ESI) *m/z* 610(M+H)⁺. Anal calcd for C₃₀H₂₅F₆NO₄S H₂O: C, 57.49; H, 4.13; N, 2.20. Found: C, 57.12; H, 3.93; N, 1.77.



Example 386


- 15 (2-Methoxyphenyl) [2,3-dichloro-4-(E-(((4-hydroxylaminocarbonyl)piperidin-1-yl)carbonyl)ethenyl)phenyl] sulfide

To a suspension of Example 319 (300 mg, 0.64 mmol) in CH₂Cl₂ (10 mL) was added oxalyl chloride (67 μL) and 2 drops of DMF. The yellow suspension was stirred at room temperature for 2 h to give an orange solution which was then

20 concentrated under reduced pressure, and dried under vacuum. An aliquot of the resulting acid chloride solution (2 mL) was added to a solution containing o-

trimethylsilyloxyamine (101 mg, 0.96 mmol), Hunig's base (122 μ L, 0.7 mmol) and DMAP (2 mg) in CH_2Cl_2 (3 mL). After the solution was stirred at room temperature for 1 h, TBAF (1.0 M solution in THF, 1.5 mL) was then added. The brown solution was stirred at room temperature for another h, then it was purified by HPLC (Zorbax,

- 5 C-18) to give the title compound as white solid (71 mg). ^1H NMR (300 MHz, DMSO- d_6) δ 1.50 (m, 2H), 1.70 (m, 2H), 2.28 (m, 1H), 2.70 (m, 1H), 3.09 (m, 1H), 3.79 (s, 3H), 4.23 (m, 1H), 4.45 (m, 1H), 6.55 (d, J = 8.8 Hz, 1H), 7.08 (t, J = 7.4 Hz, 1H), 7.25 (m, 2H), 7.48 (d, J = 7.2 Hz, 1H), 7.54 (t, J = 8.2 Hz, 1H), 7.73 (d, J = 15.3 Hz, 1H), 7.82 (d, J = 8.8 Hz, 1H), 8.55 (br s, 1H), 10.46 (s, 1H). MS (ESI $^+$) m/z 481
- 10 (M+H) $^+$.



Example 387

(2-Methoxyphenyl) [2,3-dichloro-4-(E-((N-carboxymethyl)-N-phenylamino)carbonyl)ethenyl]phenyl sulfide


15

The title compound was prepared by the procedures described in Example 1C substituting Example 1B with (2-methoxy) [2,3-dichloro-4-(E-(2-carboxyethenyl)phenyl] sulfide and substituting 6-amino-1-hexanol with N-phenylglycine ethyl ester following by hydrolysis. ^1H NMR (300 MHz, DMSO- d_6) δ

20 3.76 (s, 3H), 4.40 (s, 2H), 6.35 (d, J = 15.5 Hz, 1H), 6.46 (d, J = 8.4 Hz, 1H), 7.05 (t, J


= 7.3 Hz, 1H), 7.22 (m, 2H), 7.35 (t, $J = 7.5$ Hz, 3H), 7.44 (t, $J = 7.2$ Hz, 3H), 7.55 (t, $J = 7.4$ Hz, 1H), 7.76 (d, $J = 15.4$ Hz, 1H); MS (ESI⁺) m/z 488,490 (M+H)⁺. Anal. calcd for C₂₄H₁₉NCl₂O₄S: C, 59.02; H, 3.92; N, 2.87. Found: C, 58.71; H, 4.10; N, 2.58.

5

Example 388

(2-Methoxyphenyl) [3-chloro-6-hydroxy-4-(E-((3-carboxypiperidin-1-yl)carbonyl)ethenyl)phenyl]sulfide


10

Example 388A

(2-Methoxyphenyl) ((3-chloro-6-hydroxy-4-formyl)phenyl)sulfide

2-Methoxythiophenol (3.5 mL, 28.9 mmol) and 2,4-dichoro-6-


15 hydroxybenzaldehyde (5.00g, 26.3 mmol) were processed as described in Example 1 to provide the title disulfide (6.71 g, 87%) as a pale yellow solid. ¹H NMR (DMSO-d₆, 300 MHz) δ 10.18 (s, 1H), 7.61 (dd, $J=7.4$ Hz, $J=1.7$ Hz, 1H), 7.56 (dd, $J=7.7$ Hz, $J=1.9$ Hz, 1H), 7.25 (d, $J=7.3$ Hz, 1H), 7.11 (dt, $J=7.7$ Hz, $J=1.5$ Hz, 1H), 6.69 (d, $J=1.8$ Hz, 1H), 6.38 (d, $J=1.5$ Hz, 1H), 3.80 (s, 3H); MS (APCI) m/z 294 (M+H)⁺.



Example 388B

(2-Methoxyphenyl)(3-chloro-6-allyloxy-4-benzaldehyde)sulfide

- 5 Allyl bromide (2.0 mL, 22.8 mmol) was added to a stirred solution of Example 388A (6.71 g, 22.8 mmol), cesium carbonate (14.86 g, 45.6 mmol), and DMF (45 mL). After 21 h, the pale yellow solution was diluted with 1 N aqueous HCl (100 mL) and extracted with Et₂O (2x75 mL). The ether extracts were combined, dried (MgSO₄), filtered, and concentrated to a yellow solid (7.20 g, 94%). ¹H NMR (DMSO-d₆, 300 MHz) δ 10.28 (s, 1H), 7.58 (dd, *J*=8.4 Hz, *J*=1.7 Hz, 1H), 7.52 (dd, *J*=7.8 Hz, *J*=1.7 Hz, 1H), 7.23 (d, *J*=8.1 Hz, *J*=1.0 Hz, 1H), 7.08 (dt, *J*=7.8 Hz, *J*=1.4 Hz, 1H), 6.82 (d, *J*=1.7 Hz, 1H), 6.52 (d, *J*=1.7 Hz, 1H), 5.97 (m, 1H), 5.33 (d, *J*=17.3 Hz, 1H), 5.28 (d, *J*=10.8 Hz, 1H), 4.61 (m, 2H), 3.80 (s, 3H); MS (APCI) *m/z* 335 (M+H)⁺.
- 10
- 15


Example 388C

(2-Methoxyphenyl)[3-chloro-6-allyloxy-4-((carboxy)ethenyl)phenyl]sulfide

Example 388B was processed as detailed in Example 1B. ¹H NMR (DMSO-

5 d₆, 300 MHz) δ 7.77 (d, *J*=16.3 Hz, 1H), 7.51 (dd, *J*=7.4 Hz, *J*=1.7 Hz, 1H), 7.43 (dd, *J*=7.4 Hz, *J*=1.7 Hz, 1H), 7.19 (dd, *J*=8.1 Hz, *J*=1.0 Hz, 1H), 7.05 (dt, *J*=7.4 Hz, *J*=1.4 Hz, 1H), 6.82 (d, *J*=1.3 Hz, 1H), 6.72 (d, *J*=15.9 Hz, 1H), 6.66 (d, *J*=1.7 Hz, 1H), 6.00 (m, 1H), 5.30 (d, *J*=9.8 Hz, 1H), 5.26 (d, *J*=3.1 Hz, 1H), 4.63 (m, 2H), 3.80 (s, 3H). MS (APCI) *m/z* 377 (M+H⁺) 394 (M+NH₄⁺).

10


Example 388D

(2-Methoxyphenyl)[3-chloro-6-hydroxy-4-(E-((3-carboxypiperidin-1-yl)carbonyl)ethenyl)phenyl]sulfide

15

The allyl group of Example 388C was removed as reported in the literature (Honda, M.; Morita, H.; Nagakura, I. J. Org. Chem. **1997**, *62*, 8932.) and the carboxylic acid was converted to the amide as reported in Example 165 to provide the title compound as a white solid. ¹H NMR (DMSO-d₆, 300 MHz) δ 7.73 (d, *J*=16.3 Hz, 1H), 7.51 (d, *J*=7.4 Hz, 1H), 7.43 (d, *J*=7.4 Hz, 1H), 7.19 (d, *J*=7.9 Hz, 1H), 7.05


(dt, $J=7.8$ Hz, $J=1.1$ Hz, 1H), 6.70 (d, $J=1.8$ Hz, 1H), 6.59 (d, $J=6.59$ Hz, 1H), 4.30 (m, 1H), 3.95 (m, 2H), 3.80 (s, 3H), 2.85 (m, 2H), 1.87 (m, 2H), 1.45 (m, 2H). MS (APCI) m/z 448 ($M+H^+$).



Example 389

(2-Methoxyphenyl) [2,3-dichloro-4-(E-(4-((1-(2-phenyl-1-carboxyethyl)amino)carbonyl)piperidin-1-yl)carbonyl)ethenyl]phenyl]sulfide

The methyl ester of the title compound was prepared by the procedure described in Example 363 using L-phenylalanine methyl ester as the coupling substrate. The methyl ester was then hydrolyzed as described in Example 340 to provide the title compound. HPLC (Supelco C-18 column, water:acetonitrile 100:0-0:100, 20 minute elution, flow rate 1.5 mL/min, RT = 13.97 min; 1H NMR (300 MHz, DMSO- d_6) δ 1.45 (m, 2H), 1.56 (m, 1H), 1.68 (m, 1H), 2.41 (m, 1H), 2.71 (m, 1H), 2.83 (m, 2H), 3.08 (m, 2H), 3.79 (s, 3H), 4.12 (m, 1H), 4.30 (m, 1H), 4.41 (m, 1H), 6.55 (d, 1H), 7.09 (t, 1H), 7.22 (m, 6H), 7.48 (dd, 1H), 7.57 (m, 1H), 7.72 (d, 1H), 7.81 (d, 1H), 8.11 (m, 1H), 12.64 (br s, 1H); MS (ESI) m/e 613 ($M+H^+$).




Example 390

(2-Methoxyphenyl) [2,3-dichloro-4-(E-(4-((1-(2-hydroxy-1-carboxyethyl)amino)carbonyl)piperidin-1-yl)carbonyl)ethenyl)phenyl]sulfide

The methyl ester of the title compound was prepared by the procedure


- 5 described in Example 363 using L-serine methyl ester as the coupling substrate. The methyl ester was then hydrolyzed as described in Example 340 to give the title compound. HPLC (Supelco C-18 column, water:acetonitrile 100:0- 0:100, 20 minute elution, flow rate 1.5 mL/min, RT = 11.79 min; ¹H NMR (300 MHz, DMSO-d₆) δ 1.48 (m, 2H), 1.72 (m, 2H), 2.55 (m, 2H), 2.71 (m, 1H), 3.10 (m, 1H), 3.62 (m, 2H),
 10 3.79 (s, 3H), 4.22 (m, 2H), 4.41 (m, 1H), 6.55 (d, 1H), 7.09 (t, 1H), 7.34 (m, 2H), 7.48 (m, 1H), 7.57 (m, 1H), 7.71 (d, 1H), 7.81 (d, 1H), 7.96 (br d, 1H); MS (ESI) *m/e* 553 (M+H)⁺.



15

Example 391

(3-(1-(3-Carboxypiperidinyl)phenyl)[2,3-dichloro-4-(E-(1,2,5,6-tetrahydropyridin-1-yl)carbonyl)ethenyl)phenyl]sulfide




Example 391A

(3-bromophenyl) [2,3-dichloro-4-(E-[(1,2,3,6-tetrahydropyridin-1-yl)carbonyl]ethenyl)phenyl]sulfide

The title compound (1.2 g, 103%) was prepared from Example 384B (1.00 g,

- 5 2.48 mmol), using the procedures described in Example 340G substituting methyl isonipecotate with 1,2,3,6-tetrahydropyridine. ¹H NMR (300 MHz, DMSO-d₆) δ 2.09-2.2 (m, 2H), 3.61-3.68 (m, 1H), 3.70-3.77 (m, 1H), 4.03 (m, 1H), 4.18 (m, 1H), 5.69-5.78 (M, 1H), 5.80-5.93 (m, 1H), 6.93 (d, J=9 Hz, 1H), 7.20-7.37 (m, 1H), 7.43-7.56 (m, 3H), 7.67-7.79 (m, 2H), 7.88-7.97 (m, 1H); MS (ESI) *m/e* 470, 472 (M+H)⁺.

10


Example 391B

[3-(3-ethoxycarbonylpiperidine)] [2,3-dichloro-4-(E-[(1,2,3,6-tetrahydropyridin-1-yl)carbonyl]ethenyl)phenyl] sulfide

- 15 The title compound (50 mg, 46%) was prepared by the procedures described in Example 384D, substituting morpholine with ethyl nipecotate. ¹H NMR (300 MHz, DMSO-d₆) δ 1.17 (t, J=6.8 Hz, 3H), 1.5-1.76 (m, 3H), 1.82-1.95 (m, 1H), 2.06-2.19 (m, 2H), 2.56-2.67 (m, 1H), 2.84-2.96 (m, 1H), 3.06-3.13 (m, 1H), 3.43-3.52 (m, 1H), 3.61-3.74 (m, 2H), 3.99-4.18 (m, 4H), 5.66-5.91 (m, 2H), 6.73 (d, J=9 Hz, 1H),

341

6.92 (d, $J=7.5$ Hz, 1H), 7.06-7.12 (m, 2H), 7.31-7.39 (m, 2H), 7.75 (d, $J=15$ Hz, 1H), 7.80-7.91 (m, 1H); MS (ESI) m/e 545, 547 (M+H)⁺.



5

Example 391C


[3-(3-carboxypiperidine)] [2,3-dichloro-4-(E-[(1,2,3,6-tetrahydropyridine)-1-yl]carbonyl]ethenyl]phenyl] sulfide

The title compound (20 mg, 49%) was prepared from Example 391B (43 mg, 0.08 mmol) using the procedures described in Example 340H. ¹H NMR (DMSO- d_6 , 10 500 MHz) δ 1.51-1.64 (m, 2H), 1.68-1.73 (m, 1H), 1.87-1.94 (m, 1H), 2.07-2.19 (m, 2H), 2.51-2.57 (m, 1H), 2.83-2.89 (m, 1H), 2.99-3.04 (m, 1H), 3.61-3.73 (m, 4H), 4.02 (br s, 1H), 4.15 (br s, 1H), 5.67-5.76 (m, 1H), 5.79-5.90 (m, 1H), 6.72 (d, $J=7.5$ Hz, 1H), 6.92 ($J=6.25$ Hz, 1H), 7.10-7.13 (m, 2H), 7.14-7.30 (m, 1H), 7.36 (t, $J=7.5$ Hz, 1H), 7.74 (d, $J=15$ Hz, 1H), 7.80-7.90 (m, 1H); MS (ESI) m/e 517, 519 (M+H)⁺.

15


00541735-033100

342

Example 392

(3-(4-(3-(2-pyrrolidin-1-yl)propylamino)carbonyl)ethenyl)phenyl]sulfide


5

Example 392A

(3-bromophenyl) [2,3-dichloro-4-(E-(((3-(2-pyrrolidin-1-yl)propylamino)carbonyl)ethenyl)phenyl)]sulfide

10

The title compound (1.25 g, 95%) was prepared from Example 384B (1.00 g, 2.475 mmol), using the procedures described in Example 340G substituting methyl isonipecotate with 3-aminopropylpyrrolidine. MS (ESI) *m/e* 529, (M+H)⁺, 527, (M-H).

Example 392B


15

00120-5674560

(3-(4-Pyrrolidin-1-yl)piperidin-1-yl)phenyl [2,3-dichloro-4-(E-(((3-(2-pyrrolidinon-1-yl)propylamino)carbonyl)ethenyl)phenyl)sulfide


The title compound (32 mg, 27%) was prepared from Example 392A as described in Example 384D, substituting morpholine with 4-(1-

- pyrrolidinyl)piperidine. ^1H NMR (500 MHz, DMSO- d_6) δ 1.60-1.67 (m, 4H), 1.84-2.90 (m, 5H), 2.91-2.03 (m, 1H), 2.04-2.11 (m, 3H), 2.20 (t, $J=7.5$ Hz, 2H), 2.75 (br t, $J=12.5$ Hz, 2H), 3.00-3.16 (m, XH), 3.21 (t, $J=7.5$ Hz, XH), 3.33(m, 1H), 3.46-3.64 (m, 1H), 3.87 (br d, $J=10$ Hz, 2H), 6.59 (d, $J=15$ Hz, 1H), 6.80 (d, $J=8.75$ Hz, 1H), 6.94 (d, $J=7.5$ Hz, 1H), 7.12-7.18 (m, 2H), 7.33 (t, $J=7.5$ Hz, 1H), 7.57 (d, $J=8.75$ Hz, 1H), 7.68 (d, $J=15$ Hz, 1H), 8.24 (t, $J=5$ Hz, 1H); MS (ESI) m/e 601, 603, $(M+H)^+$; 599, 601 $(M-H)^-$.



Example 393


- [3-(4-(Spiro-2,2-dioxolanyl)piperidin-1-yl)phenyl] [2,3-dichloro-4-(E-(4-morpholinyl)carbonyl)ethenyl)phenyl)sulfide



Example 393A

(3-bromophenyl) [2,3-dichloro-4-(E-((4-morpholino)carbonyl)ethenyl)phenyl] sulfide

The title compound (980 mg, 84%) was prepared from Example 384B (1.00 g, 2.48 mmol), using the procedures described in Example 340G substituting methyl isonipecotate with morpholine. ¹H NMR (400 MHz, DMSO-d₆) 3.53-3.63 (m, 6H),
 5 3.68 (br s, 2H), 6.93 (d, J=8 Hz, 1H), 7.27 (d, J=15 Hz, 1H), 7.44-7.52 (m, 2H), 7.67-7.74 (m, 2H), 7.78 (d, J=15 Hz, 1H), 7.80 (d, J=8 Hz, 1H); MS (ESI) *m/e* 474 (M+H)⁺.




10

Example 393B


[3-(4-(Spiro-2,2-dioxolanyl)piperidin-1-yl)phenyl] [2,3-dichloro-4-(E-((4-morpholinyl)carbonyl)ethenyl)phenyl]sulfide

The title compound (32 mg, 27%) was prepared from Example 393A as described in Example 384D, substituting morpholine with 1,4-dioxo-8-
 15 azaspiro[4,5]decane. ¹H NMR (500 MHz, DMSO-d₆) δ 1.68 (t, J=5 Hz, 4H), 3.52-3.60 (m, 7H), 3.66 (br s, 2H), 3.91 (s, 4H), 6.71 (d, J=8.75 Hz, 1H), 6.91 (m, 1H), 7.11-7.13 (m, 2H), 7.22 (d, J=15 Hz, 1H), 7.35 (m, 1H), 7.76 (d, J=15 Hz, 1H), 7.85 (d, J=8.75 Hz, 1H); MS (ESI) *m/e* 535, 537 (M+H)⁺.


**Example 394**

[3-(3-Carboxypiperidin-1-yl)phenyl] [2,3-dichloro-4-(E-[(4-carboxypiperidin-1-yl)carbonyl]ethenyl)phenyl] sulfide

- 5 The title compound (51 mg, 41%) was prepared from Example 384C as described in Example 384D, substituting morpholine with ethyl nipecotate followed by hydrolysis with LiOH as described in Example 340H. ¹H NMR (400 MHz, DMSO-d₆) δ 1.39-1.60 (m, 4H), 1.67-1.76 (m, 1H), 1.82-1.96 (m, 3H), 2.52-2.59 (m, 3H), 2.81-2.93 (m, 2H), 2.99-3.07 (m, 1H), 3.14-3.25 (m, 1H), 3.47-3.54 (m, 1H),
- 10 3.69 (dd, J₁=4 Hz, J₂=12 Hz, 1H), 4.05-4.17 (m, 1H), 4.24-4.34 (m, 1H), 6.72 (d, J=8 Hz, 1H), 6.92 (d, J=8 Hz, 1H), 7.11 (m, 2H), 7.23 (d, J=15 Hz, 1H), 7.34-7.40 (m, 1H), 7.73 (d, J=15 Hz, 1H), 7.85 (d, J=8 Hz, 1H); MS (ESI) m/e 563,565 (M+H)⁺.

**Example 395**

(2-(2-Carboxyethenyl)phenyl) [2,3-dichloro-4-(E-[(4-morpholinyl)carbonyl]ethenyl)phenyl]sulfide




Example 395A

(2-(2-Tert-butyloxycarbonyl)ethenyl)phenyl [2,3-dichloro-4-(E-(4-morpholinyl)carbonyl)ethenyl]phenyl sulfide


- 5 A solution of Example (2-bromophenyl) (50 mg, 0.11 mmol), tris(benzylideneacetone)dipalladium[0] (5.1 mg, 0.0056 mmol), and tri-*o*-tolylphosphine (11 mg, 0.035 mmol) in 0.2 mL DMF was degassed with nitrogen gas for 10 min, then triethylamine (50 μ L, 36 mg, 0.36 mmol) and tert-butyl acrylate (50 μ L, 44 mg, 0.34 mmol) were added to the solution, and the vessel was sealed under
- 10 nitrogen and heated in a 100 °C oil bath for 17 h. The reaction was concentrated under hi-vacuum, and the residue was partially purified by preparative TLC eluting with 10% acetone-CH₂Cl₂ to provide 42 mg (0.080 mmol, 73%) of the title compound as a crude material. The compound was further purified by preparative HPLC (30-100% MeCN in 0.1% aqueous TFA, 40 min elution, C-18 reverse-phase Sorbax 10
- 15 mm column, producing 26 mg (0.051 mmol, 47%) of the title compound as a glass.
- ¹H NMR (300 MHz, CDCl₃) δ 1.47 (s, 9H), 2.3-2.7 (v br s, 5H), 3.54-3.90 (2 br m, 8H), 6.32 (d, J=16 Hz, 1H), 6.46 (d, J=8 Hz, 1H), 6.69 (br d, J=15 Hz, 1H), 7.24 (br d, partially overlapped with CHCl₃, approx. 1H), 7.40-7.54 (m, 2H), 7.59 (dd, J=2,8 Hz, 1H), 7.75 (dd, J=2,8 Hz, 1H), 7.94 (br d, J=15 Hz, 1H), 7.98 (d, J=16 Hz, 1H); MS
- 20 (ESI) *m/e* 520, 522 (M+H)⁺.

347

Example 395B

(2-(2-Carboxy)ethenyl)phenyl) [2,3-dichloro-4-(E-((4-morpholinyl)carbonyl)ethenyl)phenyl] sulfide


- 5 Example 395A (26 mg, 0.050 mmol) was dissolved in 1 mL chloroform and 1 mL TFA and the solution was stirred at ambient temperature for 1 h. The solvent was concentrated under reduced pressure to provide 25 mg (109%) of the title compound as an 85:15 mixture of *E*- and *Z*-cinnamide isomers. Data for major isomer: ¹H NMR (300 MHz, CDCl₃) δ 3.55-3.85 (2 br m, 9H), 6.42 (d, J=16 Hz, 1H), 6.47 (d, J=8 Hz, 1H), 6.69 (d, J=15 Hz, 1H), 7.24 (d, partially overlapped with CHCl₃, approx. 1H), 7.43-7.56 (m, 2H), 7.78 (dd, J=2,8 Hz, 2H), 7.93 (d, J=15 Hz, 1H), 8.23 (d, J=16 Hz, 1H); MS (ESI) *m/e* 464, 466 (M+H)⁺.
- 10

Example 396

[3-(4-Carboxypiperidin-1-yl)phenyl] [2,3-dichloro-4-(E-((1,2,3,6-tetrahydropyridine)-1-yl)carbonyl)ethenyl] sulfide

The title compound (22 mg, 58%) was prepared from Example 391A as described in Example 384D, substituting morpholine with ethyl isonipecotate


followed by hydrolysis with LiOH as described in Example 340H. ^1H NMR (400 MHz, DMSO- d_6) δ 1.59-1.70 (m, 2H), 1.87-1.93 (m, 2H), 2.07-2.19 (m, 2H), 2.39-2.47 (m, 1H), 2.80-2.90 (m, 2H), 4.03 (br s, 1H), 4.16 (br s, 1H), 5.68-5.76 (m, 1H), 5.79-5.90 (m, 1H), 6.72 (d, $J=8$ Hz, 1H), 6.93 (d, $J=7$ Hz, 1H), 7.13 (m, 2H), 7.17-7.3 (m, 1H), 7.36 (t, $J=7$ Hz, 1H), 7.75 (d, $J=15$ Hz, 1H), 7.80-7.90 (m, 1H); MS (ESI) m/e 517, 519 ($M+H$) $^+$.



Example 397


[3-(4-Carboxypiperidinyl)phenyl] [2,3-dichloro-4-(E)-[4-morpholinyl]carbonyl]ethenyl]phenyl] sulfide

The title compound (39 mg, 79%) was prepared from Example 393A as described in Example 384D, substituting morpholine with ethyl isonipecotate followed by hydrolysis with LiOH as described in Example 340H. ^1H NMR (400 MHz, DMSO- d_6) δ 1.56-1.68 (m, 2H), 1.86-1.92 (m, 2H), 2.38-2.46 (m, 1H), 2.77-2.86 (m, 2H), 3.52-3.61 (m, 6H), 3.65-3.72 (m, 4H), 6.71 (d, $J=8$ Hz, 1H), 6.91 (d, $J=7$ Hz, 1H), 7.10 (m, 2H), 7.21 (d, $J=15$ Hz, 1H), 7.34 (t, $J=8$ Hz, 1H), 7.76 (d, $J=15$ Hz, 1H), 7.83 (d, $J=8$ Hz, 1H); MS (ESI) m/e 521, 523 ($M+H$) $^+$.

**Example 398**


[2-(4-Acetyl piperazin-1-yl)phenyl] [2,3-dichloro-4-(E-[(4-carboxypiperidin-1-yl)carbonyl]ethenyl)phenyl] sulfide

- 5 The title compound (19 mg, 83%) was prepared from Example 384C as described in Example 384D, substituting morpholine with 4-acetyl piperazine, followed by hydrolysis with LiOH as described in Example 340H. ¹H NMR (300 MHz, DMSO-d₆) δ 1.38-1.54 (m, 2H), 1.82-11.92 (m, 2H), 2.00 (s, 3H), 2.51-2.60 (m, 1H), 2.87-3.00 (m, 5H), 3.13-3.27 (m, 1H), 3.36-3.46 (m, 4H), 4.06-4.18 (m, 1H),
- 10 4.22-4.36 (m, 1H), 6.91 (d, J=7.5 Hz, 1H), 7.10-7.17 (m, 1H), 7.20-7.25 (m, 2H), 7.28 (d, J=15 Hz, 1H), 7.39-7.45 (m, 1H), 7.77 (d, J=15 Hz, 1H), 7.89 (d, J=7.5 Hz, 1H); MS (ESI) *m/e* 562, 564 (M+H)⁺.

**Example 399**

[3-(3-Carboxypiperidin-1-yl)phenyl] [2,3-dichloro-4-(E-[(4-morpholinyl)carbonyl]ethenyl)phenyl] sulfide

The title compound (30 mg, 60%) was prepared from Example 393A as described in Example 384D, substituting morpholine with ethyl nipecotate followed by hydrolysis with LiOH as described in Example 340H. ^1H NMR (400 MHz, DMSO- d_6) δ 1.51-1.60 (m, 1H), 1.66-1.72 (m, 1H), 1.87-1.94 (m, 1H), 2.79-2.87 (m, 1H), 2.96-3.02 (m, 1H), 3.44-3.72 (m, 12H), 6.71 (d, $J=8$ Hz, 1H), 6.90 (d, $J=7$ Hz, 1H), 7.09 (m, 2H), 7.21 (d, $J=15$ Hz, 1H), 7.32-7.38 (m, 1H), 7.76 (d, $J=15$ Hz, 1H), 7.84 (d, $J=8$ Hz, 1H); MS (ESI) m/e 521, 523 ($M+H$) $^+$.



Example 400

[3-(4-Carboxypiperidin-1-yl)phenyl] [2,3-dichloro-4-(E-[(4-(dimethylaminosulfamoyl)piperazin-1-yl)carbonyl]ethenyl)phenyl] sulfide




Example 400A

N,N-Dimethyl piperazinylsulfamide

To a solution of *tert*-butyl 1-piperazinecarboxylate (2.5 g, 13.42 mmol) in tetrahydrofuran (21.5 ml, 0.25 M) at 0 °C was added triethylamine (2.25 mL, 16.11 mmol) followed by dimethylsulfamoyl chloride (1.73 mL, 16.11 mmol). The reaction

mixture was stirred at 0 °C for 1 h, diluted with ethyl acetate (100 mL) and washed with saturated NaHCO₃ solution (2x30 mL), followed by brine (2x30 mL). The dried (Na₂SO₄) organic layer was evaporated to dryness under reduced pressure and the residue obtained was treated with 10% trifluoroacetic acid in methylene chloride (20 mL) at ambient temperature. After 48 h, methylene chloride was evaporated in vacuo to obtain a colorless syrup. This crude material was made basic (1 N NaOH, 50 mL), and the mixture was extracted sequentially with ethyl acetate (2x20 mL) and methylene chloride (2x30 mL). The combined organic layers were dried (Na₂SO₄) and evaporated to dryness under reduced pressure to obtain the title compound in quantitative yield. ¹H NMR (300 MHz, DMSO-d₆) δ 2.77 (s, 3H), 2.79 (s, 3H), 3.12-3.20 (m, 7H), 3.3 (m, 1H), 8.86 (br s, 1H); MS (ESI) *m/e* 194 (M+H)⁺.




Example 400B

[3-(4-Carboxypiperidin-1-yl)phenyl] [2,3-dichloro-4-(4-(dimethylaminosulfamoyl)piperazin-1-yl)carbonyl]ethenyl]phenyl] sulfide

The title compound was prepared from Example 384B as described in Example 340G, substituting methyl isonipecotate with Example 400A, followed by amination with ethyl isonipecotate as described in Example 396. ¹H NMR (500 MHz, MeOH-d₄) δ 2.79-2.88 (m, 2H), 2.01-2.08 (m, 2H), 2.48-2.53 (m, 1H), 2.84 (s,


6H), 2.91-2.99 (m, 2H), 3.24-3.29 (m, 2H), 3.66-3.73 (m, 2H), 3.77 (m, 6H), 6.80 (d, $J=7.5$ Hz, 1H), 7.05 (d, $J=7.5$ Hz, 1H), 7.11 (d, $J=15$ Hz, 1H), 7.16-7.22 (m, 2H), 7.39 (t, $J=7.5$ Hz, 1H), 7.62 (d, $J=7.5$ Hz, 1H), 7.95 (d, $J=15$ Hz, 1H); MS (ESI) m/e 625, 627 (M-H)⁺.

5

Example 401

(2-Methoxyphenyl)[2,3-bis(trifluoromethyl)-4-(E-((3-carboxypiperidin-1-yl)carbonyl)ethenyl)phenyl]sulfide


10

Example 401A

Ethyl 2-Furylacrylate

Ethyl iodide (64 mL, 0.796 mol) was added to furylacrylic acid (100 g, 0.724 mol), diisopropylethyl amine (140 mL, 0.796 mmol), in acetonitrile (1100 mL), and the mixture was heated to 60 °C. After 18 h, the dark solution was cooled to room temperature and concentrated in vacuo. The resulting brown sludge was diluted with Et₂O (500 mL), washed with 1 N aqueous HCl (2x250 mL), washed with 0.2 N aqueous NaOH (2x250 mL), washed with saturated aqueous NaHCO₃ (1x250 mL), dried (MgSO₄), filtered, and concentrated to a black oil (114 g, 95%). ¹H NMR

(DMSO- d_6 , 300 MHz) δ 7.84 (d, $J=1.7$ Hz, 1H), 7.46 (d, $J=15.6$ Hz, 1H), 6.97 (d, $J=3.4$ Hz, 1H), 6.33 (dd, $J=3.4$ Hz, $J=1.7$ Hz, 1H), 6.22 (d, $J=15.9$ Hz, 1H), 4.17 (q, $J=7.1$ Hz, 2H), 1.24 (t, $J=7.1$ Hz, 3H); MS (APCI) m/z 167 (M+H) $^+$.



5

Example 401B


Ethyl E-2,3-bis(trifluoromethyl)-4-hydroxycinnamate

A solution of Example 401A (20 g, 0.12 mol) in tetrahydrofuran (40 mL) at -50 °C in a 600 mL Parr stirred reactor was treated with hexafluoroacetylene (24.4 g, 0.15 mol), the reactor sealed and heated to 110 °C for 22 hours, allowed to slowly cool to room temperature, and then concentrated to a brown oil (36 g). This oil was then treated with boron trifluoride etherate (33 mL, 0.275 mol) at room temperature for 17 hours, additional boron trifluoride etherate (16 mL, 0.135 mol) added, stirred six hours, cooled to 0 °C, diethyl ether (200 mL) added, followed by slow addition of 15 mL of 2M potassium carbonate (vigorous gas evolution). This mixture was diluted with additional diethyl ether, layers separated, organic layer washed with brine, dried (MgSO₄) and concentrated to give 39 grams of a brown semi-solid. This semi-solid was diluted with 75 mL of dichloromethane and then flash chromatographed on silica gel with 10-50% ethyl acetate/hexane to provide the title compound (22.8 g, 58%). mp 138-140 °C; ^1H NMR (300 MHz, d_6 DMSO) δ 11.64 (bs, 1H), 7.95 (d, 1H), 7.78 (dq, 1H), 7.33 (d, 1H), 6.47 (d, 1H), 4.21 (q, 2H), 1.26 (t,

20

3H); MS (APCI-NH₃) *m/e* 329 (M+H)⁺, 346 (M+NH₄)⁺, 327 (M-H)⁻. Analytical HPLC: 4.6X250 mm Zorbax C18 column, 1.5 mL/min, 254 nm, CH₃CN:H₂O with 0.1% TFA, 0:100 ramp to 90:10 (0-10 min), 90:10 (10-18 min), ramp to 0:100 (18-20 min), Rt = 10.6 min (98.3 area%).


5




Example 401C

Ethyl E-4-(trifluoromethanesulfonyl)-2,3-bis(trifluoromethyl)cinnamate


- Triflic anhydride (670 μ L, 3.97 mmol) was added to a mixture of Example
- 10 401B (1.00 g, 3.05 mmol) and pyridine (6.5 mL). After 2 h, the dark solution was diluted with Et₂O (75 mL), washed with 1 N aqueous HCl (2x50 mL), washed with saturated aqueous NaHCO₃ (1x75 mL), dried (MgSO₄), filtered, and concentrated to a dark amber oil (1.35 g, 96%). ¹H NMR (DMSO-d₆, 300 MHz) δ 8.33 (d, *J*=8.8 Hz, 1H), 8.11 (d, *J*=8.8 Hz, 1H), 7.87-7.78 (m, 1H), 6.67 (d, *J*=16.0 Hz, 1H), 4.24 (q, *J*=7.1 Hz, 2H), 1.27 (t, *J*=7.1 Hz, 3H); MS (APCI) *m/z* 478 (M+NH₄)⁺, 495 (M+Cl)⁻.
- 15



**Example 401E**

(2-Methoxyphenyl) [2,3-bis(trifluoromethyl)-4-(E-((3-carboxypiperidin-1-yl)carbonyl)ethenyl)phenyl]sulfide

- 5 Example 401D was processed as detailed in Examples 137 and 155 to provide the title compound (168 mg, 86%) as a white solid. ^1H NMR (DMSO- d_6 , 300 MHz) δ 7.95 (d, 1H), 7.57 (m, 1H), 7.50 (t, 1H), 7.46 (d, 1H), 7.20 (d, 1H), 7.15 (d, 1H), 7.14 (d, 1H), 7.06 (t, 1H), 4.4 (m, 1H), 4.01 (m, 2H), 3.75 (s, 3H), 1.93 (m, 2H), 1.63 (m, 2H), 1.42 (m, 2H). MS (APCI) m/z 534 ($M+H^+$). Anal. calcd for
- 10 $C_{24}H_{21}F_6NO_4S+0.75M\ H_2O$: C, 52.70; H, 4.15; N, 2.56. Found: C, 53.01; H, 3.78; N, 2.17.


**Example 402**

- 15 (2-Methoxyphenyl) [2,3-bis(trifluoromethyl)-4-(E-((2-carboxypyrrolidin-1-yl)carbonyl)ethenyl)phenyl]sulfide

Example 401D was processed as reported in Example 401E, substituting L-proline methyl ester hydrochloride for the amine. ^1H NMR (DMSO- d_6 , 300 MHz) δ 7.98 (d, $J=8.2$ Hz, 1H), 7.64 (m, 1H), 7.53 (d, $J=8.2$ Hz, 1H), 7.50 (t, $J=7.4$ Hz, 1H), 7.21 (d, 1H), 7.19 (d, 1H), 7.07 (t, $J=7.8$ Hz, 1H), 6.95 (d, $J=15.0$ Hz, 1H), 4.34 (m,

20


¹H), 3.70 (m, 2H), 3.76 (s, 3H), 2.08 (m, 2H), 1.91 (m, 2H); MS (APCI) *m/z* 520 (M+H⁺). Anal. calcd for C₂₅H₁₉F₆NO₄S: C, 53.18; H, 3.69; N, 2.70. Found: C, 52.88; H, 3.86; N, 2.43.



5

Example 403

[3-(4-Carboxypiperidin-1-yl)phenyl] [2,3-dichloro-4-(E-((4-((trifluoromethylsulfonyl)piperazin-1-yl)carbonyl)ethenyl)phenyl] sulfide




10

Example 403A

Piperazine-1-trifluoromethylsulfonamide

The title compound (1.65 g, 72%) was prepared as described in Example 400A, substituting dimethylsulfamoyl chloride with trifluoromethanesulfonyl chloride

15 (1.26 ml, 11.81 mmol). MS (ESI) *m/e* 219 (M+H)⁺.




Example 403B

[3-(4-Carboxypiperidin-1-yl)phenyl] [2,3-dichloro-4-(E-((4-
((trifluoromethylsulfonyl)piperazin-1-yl)carbonyl)ethenyl)phenyl] sulfide

Example 403B (51 mg, 38%) was prepared from Example 384B as described in Example 340G, substituting methy isonipecotate with Example 403A followed by

- 5 amination with ethyl isonipecotate as described in Example 396. ¹H NMR (500 MHz, DMSO-d₆) δ 1.56-1.66 (m, 2H), 2.84-2.91 (m, 2H), 2.37-2.45 (m, 1H), 2.77-2.86 (m, 2H), 3.63-3.70 (m, 7H), 3.72-3.85 (m, 3H), 6.72 (d, J=8.75 Hz, 1H), 6.90 (d, J=7.5 Hz, 1H), 7.09 (m, 1H), 7.11 (s, 1H), 7.21 (d, J=15 Hz, 1H), 7.34 (t, J=7.5 Hz, 1H), 7.76 (d, J=15 Hz, 1H), 7.81 (d, J=8.75 Hz, 1H); MS (ESI) *m/e* 650, 652 (M-H).

10




Example 404

(2-Methoxyphenyl)[2,3-dichloro-4-(E-(piperidin-1-ylcarbonyl)ethenyl)phenyl] sulfide

- The title compound was prepared by the procedures described in Example 1C substituting Example 1B with (2-methoxy) [2,3-dichloro-4-(E-(2-
15 carboxyethenyl)phenyl] sulfide and substituting 6-amino-1-hexanol with piperidine. ¹H NMR (300 MHz, DMSO-d₆) δ 1.48 (m, 4H), 1.59 (m, 2H), 3.55 (m, 4H), 3.79 (s, 3H), 6.55 (d, J = 8.4 Hz, 1H), 7.08 (t, J = 7.4 Hz, 1H), 7.21 (d, J = 6.0 Hz, 1H), 7.25 (s, 1H), 7.48 (dd, J = 7.8, 1.7 Hz, 1H), 7.56 (m, 1H), 7.72 (d, J = 15.6 Hz, 1H), 7.82

(d, $J = 8.5$ Hz, 1H); MS (ESI⁺) m/z 422, 424 (M+H)⁺. Anal. calcd for C₂₁H₂₁NCl₂SO₂:


C, 59.72; H, 5.01; N, 3.32. Found: C, 59.52; H, 4.94; N, 3.05.



5

Example 405

(2-Hydroxyphenyl) [2,3-dichloro-4-(E-((4-morpholino)carbonyl)ethenyl)phenyl]sulfide



10


Example 405A

(2-Hydroxyphenyl) [2,3-dichloro-4-(E-(carboxy)ethenyl)phenyl]sulfide

Boron tribromide (84 mL of a 1.0M solution in CH₂Cl₂) was added to a suspension of Example 310C in CH₂Cl₂ (85 mL) at 0 °C. After addition was completed, the ice-water bath was removed, and the homogeneous dark solution was stirred for 2h before the mixture was poured into 1 N aqueous HCl (100 mL) and ice (100 g), and extracted with EtOAc (3x100 mL). The organic layers were combined, washed with brine (1x50 mL), dried (MgSO₄), filtered, and concentrated to a white solid (11.3 g). ¹H NMR (DMSO-d₆, 300 MHz) δ 10.26 (s, 1H), 7.82 (d, $J=15.6$, 1H), 7.74 (d, $J=8.5$ Hz, 1H), 7.44 (dt, $J=7.8$ Hz, $J=1.7$ Hz, 1H), 7.41 (dd, $J=7.4$ Hz, $J=1.7$

360

Hz, 1H), 7.05 (dd, $J=8.4$ Hz, $J=1.3$ Hz, 1H), 6.94 (dt, $J=7.8$ Hz, $J=1.4$ Hz, 1H), 6.52 (d, $J=8.2$ Hz, 1H), 6.50 (d, $J=16.0$ Hz, 1H); MS (APCI) m/z 339 (M-H)⁻, 375 (M+Cl)⁻.



5


Example 405B

(2-Hydroxyphenyl) [2,3-dichloro-4-(E-((4-morpholino)carbonyl)ethenyl)phenyl]sulfide

Example 405A (11.3 g) was processed as reported in Example 310D to provide the title product (8.47 g, 62%) as a white solid. ¹H NMR (DMSO-*d*₆, 300 MHz) δ 10.24 (s, 1H), 7.81 (d, $J=8.9$, 1H), 7.77 (d, $J=14.9$ Hz, 1H), 7.44 (dt, $J=6.4$ Hz, $J=1.7$ Hz, 1H), 7.39 (dd, $J=8.2$ Hz, $J=1.7$ Hz, 1H), 7.05 (dd, $J=8.1$ Hz, $J=1.0$ Hz, 1H), 6.94 (dt, $J=7.8$ Hz, $J=1.0$ Hz, 1H), 6.52 (d, $J=8.8$ Hz, 1H), 6.53 (d, $J=8.8$ Hz, 1H); MS (APCI) m/z 410 (M+H)⁺, 446 (M+Cl)⁺.


001620-56274560

361

Example 406

(2-Methoxyphenyl) [2,3-dichloro-4-(E-(((4-
carboxyphenyl)methyl)amino)carbonyl)ethenyl]phenyl]sulfide

- 5 The title compound was prepared by the procedures described in Example 1C substituting Example 1B with (2-methoxy) [2,3-dichloro-4-(E-(2-carboxyethenyl)phenyl] sulfide and substituting 6-amino-1-hexanol with methyl 4-(aminomethyl)benzoate hydrochloride following by hydrolysis. ¹H NMR (300 MHz, DMSO-d₆) δ 3.79 (s, 3H), 4.46 (s, 2H), 6.60 (d, J = 8.1 Hz, 1H), 6.66 (d, J = 15.6 Hz, 1H), 7.08 (t, J = 8.4 Hz, 1H), 7.25 (d, J = 8.5 Hz, 1H), 7.39 (d, J = 8.5 Hz, 2H), 7.51 (m, 3H), 7.75 (d, J = 15.6 Hz, 1H), 7.90 (d, J = 8.4 Hz, 2H), 8.83 (t, J = 5.7 Hz, 1H), 12.90 (brs, 1H); MS (ESI⁺) *m/z* 488,490 (M+H)⁺. Anal. calcd for C₂₄H₁₉NC₂O₄S: C, 59.02; H, 3.92; N, 2.87. Found: C, 58.97; H, 4.07; N, 2.71.
- 10



15


00160.567H560

Example 407

(2-Methoxyphenyl) [2,3-dichloro-4-(E-(((4-pyrrolidin-1-yl)piperidin-1-yl)carbonyl)ethenyl)phenyl]sulfide

The title compound was prepared by the procedures described in Example 1C

5 substituting Example 1B with (2-methoxy) [2,3-dichloro-4-(E-(2-carboxyethenyl)phenyl] sulfide and substituting 6-amino-1-hexanol with 4-(pyrrolidinyl)piperidine. ¹H NMR (300 MHz, DMSO-d₆) δ 1.48 (m, 2H), 1.84 (m, 2H), 2.00 (m, 2H), 2.10 (m, 2H), 2.65 (m, 1H), 3.10 (m, 3H), 3.35 (m, 1H), 3.50 (m, 1H), 3.80 (s, 3H), 4.38 (m, 2H), 4.52 (m, 1H), 6.56 (d, J = 8.5 Hz, 1H), 7.08 (t, J = 7.8 Hz, 1H), 7.22 (d, J = 7.8 Hz, 1H), 7.26 (d, J = 15.2 Hz, 1H), 7.48 (dd, J = 7.8, 1.7 Hz, 1H), 7.57 (t, J = 8.2 Hz, 1H), 7.76 (d, J = 15.3 Hz, 1H), 7.82 (d, J = 7.8 Hz, 1H); MS (ESI⁺) *m/z* 491, 493 (M+H)⁺. Anal. calcd for C₂₅H₂₈N₂Cl₂O₂S 1.8 TFA: C, 49.30; H, 4.31; N, 4.02. Found: C, 49.08; H, 4.31; N, 3.97.


Example 408

(2-Hydroxyphenyl)[2,3-dichloro-4-(E-((4-carboxypiperidin-1-yl)carbonyl)ethenyl)phenyl]sulfide

Example 405A (119 mg) was processed as detailed in Example 165 to provide

20 the title compound as a white solid (43 mg, 28%). ¹H NMR (DMSO-d₆, 300 MHz) δ 10.23 (s, 1H), 7.81 (d, J=8.8, 1H), 7.72 (d, J=15.2 Hz, 1H), 7.42 (dt, J=7.8 Hz, J=1.7

Hz, 1H), 7.39 (dd, $J=7.1$ Hz, $J=1.7$ Hz, 1H), 7.21 (d, $J=15.3$ Hz, 1H), 7.05 (dd, $J=8.2$ Hz, $J=1.0$ Hz, 1H), 6.93 (dt, $J=7.4$ Hz, $J=1.0$ Hz, 1H), 6.53 (d, $J=8.5$ Hz, 1H), 4.25 (m, 1H), 4.03 (m, 2H), 2.85 (m, 2H), 1.87 (m, 2H), 1.44 (m, 2H). MS (APCI) m/z 452 ($M+H^+$), 450 ($M-H^+$), 486 ($M+Cl^-$). Anal. calcd for $C_{21}H_{19}Cl_2NO_4S+0.25M H_2O$: C, 55.21; H, 4.30; N, 3.07. Found: C, 55.26; H, 4.29; N, 2.72.



Example 409

10 [3-(4-Carboxypiperidin-1-yl)phenyl] [2,3-dichloro-4-(E-((4-((methylsulfonyl)piperazin-1-yl)carbonyl)ethenyl)phenyl)] sulfide




Example 409A

Piperazine methylsulfonamide


15 The title compound (1.65 g, 72%) was prepared as described in Example 400A, substituting dimethylsulfamoyl chloride with methanesulfonyl chloride (1.26 mL, 11.81 mmol).

364

Example 409B

[3-(4-Carboxypiperidin-1-yl)phenyl] [2,3-dichloro-4-(E-((4-(methylsulfonyl)piperazin-1-yl)carbonyl)ethenyl)phenyl] sulfide


5 Example 409B (48 mg, 72%) was prepared from Example 384B as described in Example 340G, substituting methyl isonipecotate with Example 409A followed by amination with ethyl isonipecotate as described in Example 396. ¹H NMR (300 MHz, DMSO-d₆) δ 1.55-1.71 (m, 2H), 1.83-1.94 (m, 2H), 2.36-2.48 (m, 1H), 2.77-2.86 (m, 2H), 2.88 (s, 3H), 3.10-3.18 (m, 4H), 3.66-3.84 (m, 6H), 6.73 (d, J=8 Hz, 1H), 6.93 (d, J=7.5 Hz, 1H), 7.11 (m, 1H), 7.13 (s, 1H), 7.25 (d, J=15 Hz, 1H), 7.32-7.41 (m, 1H), 7.78 (d, J=15 Hz, 1H), 7.85 (d, J=8 Hz, 1H); MS (ESI) *m/e* 598, 600 (M+H)⁺; 596, 598 (M-H)⁻.

Example 410

(2-Aminophenyl) [2,3-dichloro-4-(E-((4-morpholinyl)carbonyl)ethenyl)phenyl] sulfide


00544795.03100 DATED: 5/6/2015

365

Example 410A*tert*-Butyl 2,3-dichloro-4-((trifluoromethyl)sulfonyloxy)cinnamate

The title compound was constructed according to the procedure for Example

- 5 340D and 340E, except using *tert*-butyl acrylate instead of methyl acrylate. ¹H NMR (300 MHz, DMSO-*d*₆) δ 8.11 (d, 1H), 7.78 (d, 1H), 7.72 (d, 1H), 6.72 (d, 1H), 1.5 (s, 9H); MS (APCI-NH₃) *m/e* 456 (M+Cl)⁻.


Example 410B*tert*-Butyl 2,3-dichloro-4-((triisopropylsilyl)thio)cinnamate

- Sodium hydride (3.05 g of 60% dispersion, 76 mmol) that had been rinsed with dry tetrahydrofuran (2x), was suspended in 128 mL of THF, cooled to -5 °C, and slowly treated with triisopropylsilyl thiol (12.2 mL, 57 mmol), maintaining an internal
- 15 temperature below 4 °C, stirred at 0 °C for 1.5 h, then added to a second flask containing Example 410A (20 g, 47.4 mmol) and tetrakis(triphenyl)phosphine palladium (4.4 g, 3.8 mmol) in 95 mL of THF. The reaction was heated at reflux for 8 h, then allowed to cool to ambient temperature and concentrated. The resultant slurry

was diluted with ethyl acetate, filtered through celite, washed with brine, dried (Na_2SO_4) and concentrated. The resultant black residue was flash chromatographed on silica gel with 2.5-5% acetone/hexane to provide the title compound (18.2 g, 83%).

^1H NMR (300 MHz, $\text{DMSO}-d_6$) δ 7.82 (d, 1H), 7.78 (d, 1H), 7.0 (d, 1H), 6.5 (d, 1H),

5 1.5 (s, 9H), 1.35 (m, 3H), 1.09 (d, 18H); MS (APCI- NH_3) m/e 462 ($\text{M}+\text{H}$) $^+$.




Example 410C


(2-Nitrophenyl) [2,3-dichloro-4-(E-((tert-butyloxycarbonyl)ethenyl)phenyl] sulfide

- 10 A solution of Example 410B in toluene (40 mL) was treated with cesium fluoride (600 mg, 4 mmol) followed by 2-fluoronitrobenzene (5.03 mL, 47.4 mmol), then heated at reflux for 2 h, then allowed to cool and the mixture was concentrated under reduced pressure. The resultant dark brown slurry was diluted with ethyl acetate, washed with water (2x), 1 M NaOH, (2x), water (2x), dried (Na_2SO_4) and
- 15 concentrated. The 21.2 grams of crude product was flash chromatographed on silica gel with 10% acetone/hexane to provide the title compound (8.92 g, 53%). ^1H NMR (300 MHz, $\text{DMSO}-d_6$) δ 8.17 (dd, 1H), 7.95 (d, 1H), 7.83 (d, 1H), 7.78 (m, 1H), 7.48 (m, 1H), 7.3 (dd, 1H), 7.17 (d, 1H), 6.6 (d, 1H), 1.5 (s, 3H); MS (APCI- NH_3) m/e 427 ($\text{M}+\text{H}$) $^+$.

367


Example 410D(2-Nitrophenyl) [2,3-dichloro-4-(E-((carboxy)ethenyl)phenyl] sulfide

A solution of Example 410C (3.2 g, 7.5 mmol) in dichloromethane (12 mL) at
 5 room temperature was treated with trifluoroacetic acid (4 mL), stirred 30 minutes, and
 concentrated to give the title compound (2.8 g, 100%) as an off-white solid. ¹H NMR
 (300 MHz, DMSO-d₆) δ 8.16 (dd, 1H), 7.94 (d, 1H), 7.86 (d, 1H), 7.76 (m, 1H), 7.48
 (m, 1H), 7.29 (dd, 1H), 7.11 (d, 1H), 6.61 (d, 1H); MS (APCI-NH₃) *m/e* 371 (M+H)⁺.

Example 410E(2-Nitrophenyl) [2,3-dichloro-4-(E-((4-morpholinyl)carbonyl)ethenyl)phenyl] sulfide

A solution of Example 410D (2.7 g, 7.29 mmol) in dimethylformamide (32
 mL) was treated with hydroxybenzotriazole hydrate (1.2 g, 8.0 mmol), morpholine
 15 (1.4 mL, 16 mmol) and then 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide
 hydrochloride (1.53 g, 8.0 mmol), stirred at room temperature for 64 hours. The
 heterogeneous mixture was filtered, the white solid washed with water, and then dried
 in a vacuum oven at 50 °C for 24 hours to provide 2.8 g (88%) of the title compound

as a white powder. mp 210-213 °C; ^1H NMR (300 MHz, d_6 DMSO) δ 8.15 (dd, 1H), 8.03(d, 1H), 7.82 (d, 1H), 7.74 (m, 1H), 7.45 (m, 1H), 7.32 (d, 1H), 7.2 (m, 2H), 3.7 (m, 2H), 3.6 (m, 6H); MS (APCI- NH_3) m/e 440 ($\text{M}+\text{H}$) $^+$.




Example 410F

(2-Aminophenyl) [2,3-dichloro-4-(E-((4-morpholinyl)carbonyl)ethenyl)phenyl] sulfide


- A solution of iron powder (1.3 g, 22.8 mmol) and ammonium chloride (292 mg, 5.46 mmol) in ethanol (9 mL) and distilled water (9 mL) at 105 °C was treated with example 410F (2 g, 4.55 mmol) in ethanol (20 mL), stirred for one hour and then allowed to cool to room temperature. The resultant heterogeneous black mixture was filtered through a plug of Celite, rinsed through with ethyl acetate (100 mL), the filtrate washed with 1 M potassium carbonate, brine, dried (Na_2SO_4) and concentrated to give 1.9 g (100%) of the title compound as an off-white powder. mp 230-240 °C (dec); ^1H NMR (300 MHz, d_6 DMSO) δ 7.9 (d, 1H), 7.8 (d, 1H), 7.2 (d, 1H), 6.95 (dt, 1H), 6.84 (m, 2H), 6.68 (d, 1H), 6.58 (dt, 1H), 5.05 (bs, 2H), 3.7 (m, 2H), 3.6 (m, 6H); MS (APCI- NH_3) m/e 410 ($\text{M}+\text{H}$) $^+$; Analytical HPLC: 4.6X250 mm Zorbax C18 column, 1.5 mL/min, 254 nm, $\text{CH}_3\text{CN}:\text{H}_2\text{O}$ with 0.1% TFA, 0:100 ramp to 90:10 (0-10 min), 90:10 (10-18 min), ramp to 0:100 (18-20 min), R_t = 9.2 min.

5



[3-(4-Ethoxycarbonyl)piperidin-1-yl]phenyl] [2,3-dichloro-4-(E-(thiomorpholin-
ylcarbonyl)ethenyl)phenyl] sulfide

The title compound was prepared by the procedures described in Example 397B substituting morpholine with thiomorpholine. MS (APCI⁺) *m/z* 565, 567 (M+H)⁺.




Example 411B

(3-(4-carboxypiperidin-1-yl)phenyl)[2,3-dichloro-4-(E-((S-oxothiomorpholin-1-yl)carbonyl)ethenyl)phenyl] sulfide


- 5 To a solution of Example 411A (107 mg, 0.189 mmol) in CH_2Cl_2 (6 mL) was added mCPBA (80%, 41 mg, 0.189 mmol) at 0 °C. After stirring at the same temperature for 2 h, THF (2 mL) was added. The solution was concentrated to 1 mL, and was diluted with THF to 3 mL. Lithium hydroxide monohydrate (24 mg) in water (1 mL) was then added. The mixture was stirred at room temperature for 3 hours.
- 10 The formed transparent solution was separated by HPLC (Zorbax C-18) to give the title compound (68 mg). ^1H NMR (300 MHz, $\text{DMSO}-d_6$) δ 1.64 (m, 2H), 1.90 (m, 2H), 2.41 (m, 1H), 2.86 (m, 4H), 3.62 (m, 2H), 3.95 (m, 1H), 4.18 (m, 1H), 4.3 (m, 4H), 6.71 (d, J = 8.4 Hz, 1H), 6.93 (d, J = 7.5 Hz, 1H), 7.12 (d, J = 7.5 Hz, 1H), 7.13 (s, 1H), 7.28 (d, J = 15.3 Hz, 1H), 7.36 (t, J = 8.8 Hz, 1H), 7.80 (d, J = 15.3 Hz, 1H),
- 15 7.88 (d, J = 8.8 Hz, 1H); MS (APCI $^+$) m/z 553, 555 ($\text{M}+\text{H}$) $^+$. Anal. calcd for $\text{C}_{25}\text{H}_{26}\text{N}_2\text{Cl}_2\text{S}_2\text{O}_4$ 2 TFA: C, 44.57; H, 3.61; N, 3.58. Found: C, 44.34; H, 3.76; N, 3.51.

371

Example 412

[3-(4-Carboxypiperidin-1-yl)phenyl] [2,3-dichloro-4-(E-((4-hydroxypiperidin-1-yl)carbonyl)ethenyl)phenyl] sulfide


5

Example 412A

(3-Bromophenyl) [2,3-dichloro-4-(E-((tert-butyloxycarbonyl)ethenyl)phenyl] sulfide

- To a solution of Example 384B (2.35 g, 5.82 mmol) in THF (23 mL) at 5 °C
- 10 was added *tert*-butyl trichloroacetimidate (2.6 mL, 14.54 mmol) and boron trifluoride-etherate (2.35 mL, 18.54 mmol). The solution was stirred at the same temperature for 10 minutes, and was then warmed up to room temperature for 5 h. The yellow solution was poured into aq. NaHCO₃ solution, and the mixture was extracted with ethyl acetate. The combined organic phases were washed with water, dried over
- 15 anhydrous MgSO₄, and concentrated. The residual white solid was dissolved in CH₂Cl₂ and was precipitated by adding hexane. The formed suspension was filtered through silica gel, and washed with 1:8 EtOAc/hexane. The solution was concentrated and was further purified by flash chromatography (silica gel, 1:20


EtOAc/hexane) to give the title compound (2.50 g, 94%). MS (APCI⁺) *m/z* 461 (M+H)⁺.



Example 412B


(3-(4-carboethoxypiperidin-1-yl)phenyl)[2,3-dichloro-4-(E-(carbo-t-butoxyethenyl)phenyl)] sulfide

A pressure tube was charged with Example 412A (589 mg, 1.28 mmol), Pd₂(dba)₃ (30 mg, 0.032 mmol), 2-dicyclohexylphosphanyl-2'-dimethylaminobiphenyl (26 mg, 0.064 mmol), and anhydrous K₃PO₄ (382 mg, 1.8 mmol), and was purged with nitrogen. DME (4 mL) and ethyl isonipecotate (242 mg, 1.54 mmol) were added via syringe, and the mixture was purged with nitrogen again. The red reaction mixture was stirred at room temperature for 0.5 h and at 95 °C for 15 h. After the reaction mixture was cooled, the it was diluted with ethyl acetate, and washed with brine. The aqueous phase was extracted with ethyl acetate. The combined ethyl acetate solution was concentrated and the residual oil was separated by flash chromatography (silica gel, 1:6 EtOAc/hexane) to give the title compound (523 mg, 76%). MS (APCI⁺) *m/z* 536 (M+H)⁺.

Example 412C

[3-(4-(Ethoxycarbonyl)piperidin-1-yl)phenyl] [2,3-dichloro-4-(E-
(carboxy)ethenyl)phenyl] sulfide


- 5 To a solution of Example 412B (510 mg, 0.95 mmol) in CH_2Cl_2 (8 mL) at 0 °C was added trifluoroacetic acid (1.6 mL). The yellow solution was stirred at 0 °C for 1 h, and was warmed to room temperature for 3 h. After diluting with CH_2Cl_2 , the solution was poured into aq. NaHCO_3 solution. The inorganic phase was acidified to pH 5, and was extracted with 10% MeOH in CH_2Cl_2 . The combined organic phases
- 10 were washed with water, concentrated under vacuum and dried to give the title compound (472 mg, 100%). MS (APCI⁺) m/z 480 (M+H)⁺.

Example 412D

- 15 [3-(4-carboxypiperidin-1-yl)phenyl] [2,3-dichloro-4-(E-((4-hydroxypiperidin-1-
yl)carbonyl)ethenyl)phenyl] sulfide

To a suspension of Example 412C (150 mg, 0.31 mmol) in DMF (3 mL) was added 4-hydroxypiperidine (63 mg, 0.62 mmol), 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide (120 mg, 0.62 mmol), HOBt (84 mg, 0.62 mmol) and triethylamine (87 μ L, 0.62 mmol) at room temperature. The mixture was stirred at the same


- 5 temperature for 15 h. Ethyl acetate was added, the mixture was washed with brine, water, and was concentrated. The residual oil was dissolved in THF (3 mL), and was added lithium hydroxide monohydrate (26 mg, 0.62 mmol) in water (1.5 mL). After stirring for 15 hours, the solution was separated by HPLC (Zorbax C-18) to give the title compound (132 mg, 55%). ^1H NMR (300 MHz, DMSO- d_6) δ 1.32 (m, 2H), 1.65
- 10 (m, 2H), 1.75 (m, 2H), 1.92 (m, 2H), 2.43 (m, 1H), 2.86 (t, J = 10.6 Hz, 2H), 3.15 (m, 1H), 3.32 (m, 1H), 3.71 (m, 3H), 3.95 (m, 2H), 6.73 (d, J = 8.5 Hz, 1H), 6.94 (d, J = 7.2 Hz, 1H), 7.13 (d, J = 7.8 Hz, 1H), 7.13 (s, 1H), 7.24 (d, J = 15.2 Hz, 1H), 7.37 (t, J = 8.1 Hz, 1H), 7.72 (d, J = 15.2 Hz, 1H), 7.85 (d, J = 8.5 Hz, 1H); MS (ESI $^+$) m/z 535, 537 (M+H) $^+$. Anal. calcd for $\text{C}_{26}\text{H}_{28}\text{N}_2\text{Cl}_2\text{SO}_4$ 0.25 TFA: C, 56.43; H, 5.05; N, 4.97.
- 15 Found: C, 56.37; H, 5.00; N, 4.91.

Example 413

(2-Glycoxyphenyl) [2,3-dichloro-4-(E-((4-morpholino)carbonyl)ethenyl)phenyl]sulfide

- 5 Diethyl azodicarboxylate (270 μ L, 1.47 mmol) was added to a suspension of Example 405 (400 mg, 0.95 mmol), triphenylphosphine (386 mg, 1.47 mmol), and THF (2.0 mL). After 16 h, the dark orange solution was diluted with EtOAc (40 mL), washed with 1 N aqueous HCl (1x20 mL), washed with 0.2 N aqueous NaOH (1x20 mL), washed with brine (1x20 mL), dried (MgSO_4), filtered, and concentrated. Flash
- 10 silica gel column chromatography (9:1 hexane:ethyl acetate) provided a mix of desired ester and triphenyl phosphine oxide. The mixture (200 mg) was combined with lithium hydroxide, monohydrate (34 mg, 0.81 mmol), THF (0.5 mL), and H_2O (0.5 mL). After 21 h, the cloudy solution was diluted with 0.2 N aqueous NaOH (30 mL), washed with CH_2Cl_2 (2x15 mL), combined with 1 N aqueous HCl until pH<2,
- 15 and extracted with EtOAc (2x20 mL). The EtOAc extracts were combined, washed with brine (1x20 mL), dried (MgSO_4), filtered, and concentrated to a white solid (87 mg, 47%). ^1H NMR ($\text{DMSO}-d_6$, 300 MHz) δ 7.80 (d, $J=7.8$, 1H), 7.77 (d, $J=15.3$ Hz, 1H), 7.51 (dt, $J=8.1$ Hz, $J=2.0$ Hz, 1H), 7.48 (d, $J=8.1$ Hz, 1H), 7.22 (d, $J=15.3$ Hz, 1H), 7.09 (d, $J=7.8$ Hz, 1H), 7.08 (dt, $J=7.1$ Hz, $J=1.0$ Hz, 1H), 6.71 (d, $J=8.9$ Hz,
- 20 1H), 4.77 (s, 2H), 3.66 (s, 2H), 3.58 (s, 6H); MS (APCI) m/z 468 ($\text{M}+\text{H}^+$); 466 ($\text{M}-\text{H}^-$)

, 502 (M+Cl)⁻. Anal. calcd for C₂₁H₁₉Cl₂NO₃S: C, 53.85; H, 4.09; N, 2.99. Found: C, 54.07; H, 4.28; N, 2.69.




5

Example 414


(2-(4-Butyroxyl)phenyl)[2,3-dichloro-4-(E-((4-morpholino)carbonyl)ethenyl)phenyl]sulfide

Ethyl 4-bromobutyrate was added to a mixture of Example 405 (300 mg, 0.731 mmol), cesium carbonate (358 mg, 1.10 mmol), and DMF (1.5 mL). After 16 h, the pale milky solution was diluted with EtOAc (30 mL), washed with 1 N aqueous HCl (2x25 mL), washed with brine (1x25 mL), dried (MgSO₄), filtered, and concentrated to a white solid (326 mg, 85%) as the ethyl ester. The ethyl ester (312 mg, 0.595 mmol), THF (1.5 mL), and H₂O (1.5 mL) were combined with lithium hydroxide, monohydrate (63 mg, 1.50 mmol). After 18 h, the clear solution was poured into 1 N aqueous HCl (25 mL) and extracted with EtOAc (2x25 mL). The organic layers were combined, dried (MgSO₄), filtered, and concentrated to a white solid (247 mg, 85%). ¹H NMR (DMSO-d₆, 300 MHz) δ 7.79 (d, *J*=8.5, 1H), 7.77 (d, *J*=15.6 Hz, 1H), 7.51 (dt, *J*=7.5 Hz, *J*=1.7 Hz, 1H), 7.48 (dd, *J*=7.5 Hz, *J*=1.0 Hz, 1H), 7.20 (d, *J*=14.9 Hz, 1H), 7.19 (d, *J*=9.5 Hz, 1H), 7.06 (t, *J*=7.5 Hz, 1H), 6.63 (d, *J*=8.5 Hz, 1H), 4.01 (t, *J*=6.1 Hz, 2H), 3.65 (s, 2H), 3.58 (s, 6H), 2.10 (t, *J*=7.4 Hz, 2H), 1.75 (m, 2H); MS (APCI) *m/z* 496 (M+H)⁺.

Example 415


[3-(4-Carboxypiperidin-1-yl)phenyl] [2,3-dichloro-4-(E-((4-hydroxyethyl)piperazin-1-yl)carbonyl)ethenyl]phenyl sulfide

The title compound was prepared by the procedures described in Example 412 substituting 4-hydroxypiperidine with 1-hydroxyethylpiperazine. ¹H NMR (300 MHz, DMSO-d₆) δ 1.70 (m, 2H), 1.94 (m, 2H), 2.98 (m, 2H), 3.05 (m, 2H), 3.18 (m, 2H), 3.54 (m, 2H), 3.65 (m, 3H), 3.78 (m, 2H), 6.77 (d, J = 8.8 Hz, 1H), 7.03 (d, J = 6.8 Hz, 1H), 7.28 (d, J = 14.9 Hz, 1H), 7.29 (m, 2H), 7.42 (t, J = 7.8 Hz, 1H), 7.78 (d, J = 15.3 Hz, 1H), 7.86 (d, J = 8.8 Hz, 1H); MS (ESI⁺) *m/z* 564, 566 (M+H)⁺.

Example 416

[3-(4-Carboxypiperidin-1-yl)phenyl] [2,3-dichloro-4-(E-((4-furoyl)piperazin-1-yl)carbonyl)ethenyl]phenyl sulfide

The title compound was prepared by the procedures described in Example 412 substituting 4-hydroxypiperidine with 1-furopyperazine. ¹H NMR (300 MHz, DMSO-d₆) δ 1.64 (m, 2H), 1.91 (m, 2H), 2.43 (m, 1H), 2.87 (m, 2H), 3.70 (m, 10H), 6.43 (m, 1H), 6.72 (d, J = 8.5 Hz, 1H), 6.94 (d, J = 7.8 Hz, 1H), 7.03 (d, J = 3.3 Hz, 1H), 7.13 (d, J = 7.8 Hz, 1H), 7.14 (s, 1H), 7.26 (d, J = 15.2 Hz, 1H), 7.36 (t, J = 7.5 Hz, 1H), 7.77 (d, J = 15.2 Hz, 1H), 7.86 (m, 2H); MS (ESI⁺) *m/z* 614, 616 (M+H)⁺. Anal. calcd for C₃₀H₂₉N₃Cl₂SO₃ 1.5 TFA: C, 50.45; H, 3.91; N, 5.35. Found: C, 50.53; H, 3.96; N, 5.35.




Example 417

[3-(4-Carboxypiperidin-1-yl)phenyl] [2,3-dichloro-4-(E-((pyrrolidin-1-yl)carbonyl)ethenyl)phenyl] sulfide

The title compound was prepared by the procedures described in Example 412 substituting 4-hydroxypiperidine with pyrrolidine. ¹H NMR (300 MHz, DMSO-d₆) δ 1.63 (m, 2H), 1.79 (m, 2H), 1.88 (m, 4H), 2.43 (m, 1H), 2.82 (m, 2H), 3.39 (t, J = 6.7 Hz, 2H), 3.59 (t, J = 6.8 Hz, 2H), 3.68 (m, 2H), 6.71 (d, J = 8.5 Hz, 1H), 6.92 (d, J = 7.5 Hz, 1H), 6.96 (d, J = 15.3 Hz, 1H), 7.12 (d, J = 6.8 Hz, 1H), 7.13 (s, 1H), 7.35 (t, J = 8.8 Hz, 1H), 7.72 (d, J = 15.3 Hz, 1H), 7.80 (d, J = 8.5 Hz); MS (ESI⁺) *m/z* 505, 507

(M+H)⁺. Anal. calcd for C₂₅H₂₆N₂Cl₂SO₃ 0.8 TFA: C, 53.54; H, 4.53; N, 4.69.

Found: C, 53.74; H, 4.40; N, 4.64.




Example 418

[3-(4-Carboxypiperidin-1-yl)phenyl] [2,3-dichloro-4-(E- ((diethylaminocarbonyl)ethenyl)phenyl)] sulfide

The title compound was prepared by the procedures described in Example 412 substituting 4-hydroxypiperidine with diethylamine. ¹H NMR (300 MHz, DMSO-d₆)

- 10 δ 1.06(t, J = 6.7 Hz, 3H), 1.11 (t, J = 6.7 Hz, 3H), 1.63 (m, 2H), 1.88 (m, 2H), 2.43
(m, 1H), 2.82 (m, 2H), 3.35 (q, J = 6.7 Hz, 2H), 3.47 (q, J = 6.7 Hz, 2H), 3.70 (m,
2H), 6.68 (d, J = 8.5 Hz, 1H), 6.92 (d, J = 7.5 Hz, 1H), 7.07 (d, J = 15.2 Hz, 1H), 7.12
(d, J = 8.6 Hz, 1H), 7.13 (s, 1H), 7.35 (t, J = 8.8 Hz, 1H), 7.75 (d, J = 15.3 Hz, 1H),
7.84 (d, J = 8.5 Hz, 1H); MS (ESI⁺) m/z 507 (M+H)⁺. Anal. calcd for C₂₅H₂₆N₂Cl₂SO₃
15 0.2 TFA: C, 57.53; H, 5.36; N, 5.28. Found: C, 57.68; H, 5.38; N, 5.33.




Example 419

[3-(4-Carboxypiperidin-1-yl)phenyl] [2,3-dichloro-4-(E-((4-ethylpiperazin-yl)carbonyl)ethenyl)phenyl] sulfide

The title compound was prepared by the procedures described in Example 412

- 5 substituting 4-hydroxypiperidine with 1-ethylpiperazine. ¹H NMR (300 MHz, DMSO-d₆) δ 1.22 (t, J = 7.5 Hz, 3H), 1.63 (m, 2H), 1.87 (m, 2H), 2.42 (m, 1H), 2.81 (t, J = 10.5 Hz, 2H), 3.00 (m, 2H), 3.15 (m, 2H), 3.40 (m, 1H), 3.51 (m, 2H), 3.67 (m, 2H), 4.50 (m, 2H), 6.73 (d, J = 8.8 Hz, 1H), 6.89 (d, J = 7.8 Hz, 1H), 7.11 (m, 2H), 7.28 (d, J = 15.2 Hz, 1H), 7.36 (m, 1H), 7.80 (d, J = 15.2 Hz, 1H), 7.86 (d, J = 8.5 Hz); MS (APCI⁺) *m/z* 548, 550 (M+H)⁺. Anal. calcd for C₂₇H₃₁N₃Cl₂SO₃ 2.2 TFA: C, 47.18; H, 4.19; N, 5.26. Found: C, 47.27; H, 4.27; N, 5.30.
- 10

Example 420


- 15 [3-(4-Carboxypiperidin-1-yl)phenyl] [2,3-dichloro-4-(E-((4-(aminocarbonyl)piperidin-1-yl)carbonyl)ethenyl)phenyl] sulfide

The title compound was prepared by the procedures described in Example 412 substituting 4-hydroxypiperidine with isonipecotamide. ¹H NMR (300 MHz, DMSO-d₆) δ 1.45 (m, 2H), 1.63 (m, 2H), 1.73 (m, 2H), 1.87 (m, 2H), 2.43 (m, 2H), 2.78 (m, 2H), 3.10 (m, 2H), 3.7 (m, 3H), 4.30 (m, 1H), 4.40 (m, 1H), 6.70 (d, J = 8.5 Hz, 1H),

20

6.78 (s, 1H), 6.92 (d, $J = 7.5$ Hz, 1H), 7.12 (m, 2H), 7.25 (d, $J = 15.2$ Hz, 1H), 7.27 (s, 1H), 7.35 (m, 1H), 7.73 (d, $J = 15.2$ Hz, 1H), 7.86 (d, $J = 8.6$ Hz, 1H); MS (APCI⁺) m/z 562 (M+H)⁺. Anal. calcd for C₂₇H₂₉N₃Cl₂SO₄ 0.2 TFA: C, 56.23; H, 5.03; N, 7.18. Found: C, 56.41; H, 4.96; N, 6.98.


5

Example 421

[3-(4-Carboxypiperidin-1-yl)phenyl] [2,3-dichloro-4-(E-(4-(2-(ethoxyethyl)piperidin-1-yl)carbonyl)ethenyl)phenyl] sulfide


10 The title compound was prepared by the procedures described in Example 412 substituting 4-hydroxypiperidine with 1-(2-ethoxyethyl)piperazine. ¹H NMR (300 MHz, DMSO-d₆) δ 1.15 (t, $J = 6.8$ Hz, 3H), 1.63 (m, 2H), 1.90 (m, 2H), 2.42 (m, 1H), 2.81 (t, $J = 10.2$ Hz, 2H), 3.09 (m, 2H), 3.32 (m, 2H), 3.50 (q, $J = 6.8$ Hz, 2H), 3.51 (m, 2H), 3.68 (m, 4H), 4.45 (m, 2H), 6.73 (d, $J = 8.6$ Hz, 1H), 6.91 (d, $J = 7.5$ Hz, 1H),
 15 7.11 (m, 2H), 7.26 (d, $J = 15.3$ Hz, 1H), 7.36 (m, 1H), 7.80 (d, $J = 15.2$ Hz, 1H), 7.83 (d, $J = 8.5$ Hz, 1H); MS (APCI⁺) m/z 592 (M+H)⁺. Anal. calcd for C₂₉H₃₅N₃Cl₂SO₄ 2.5 TFA: C, 46.53; H, 4.31; N, 4.79. Found: C, 46.51; H, 4.31; N, 4.77.

382

Example 422


[3-((4-Carboxymethyl)piperazin-1-yl)phenyl] [(2,3-dichloro-4-(E-(4-morpholinyl)carbonyl)ethenyl)phenyl] sulfide

- 5 The title compound (24 mg, 42%) was prepared from Example 393A as described in Example 384D, substituting morpholine with 1-((ethoxycarbonyl)methyl)piperazine, followed by hydrolysis with LiOH as described in Example 340H. ^1H NMR (300 MHz, MeOH- d_4) δ 3.54 (s, 8H), 3.69 (s, 8H), 4.11 (s, 2H), 6.77 (d, $J=8.5$ Hz, 1H), 7.06 (d, $J=15$ Hz, 1H), 7.08 (m, 1H), 7.42 (t, $J=7.5$ Hz, 1H), 7.59 (d, $J=8.25$ Hz, 1H), 7.93 (d, $J=15$ Hz, 1H); MS (ESI) m/e 536, 538 (M+H) $^+$.

Example 423


- 15 [3-(4-Carboxypiperidin-1-yl)phenyl] [2,3-bis(trifluoromethyl)-4-(E-(4-morpholino)carbonyl)ethenyl]phenyl sulfide

383

Example 423A

[3-Bromophenyl] [2,3-bis(trifluoromethyl)-4-(E-(2-carboxy)ethenyl)phenyl]sulfide


The title compound was prepared from Example 401C using the procedures
 5 described in Example 384A, followed by hydrolysis with LiOH as given in Example
 340H.

Example 423B

10 [3-(4-Carboxypiperidin-1-yl)phenyl] [2,3-bis(trifluoromethyl)-4-(E-((4-
morpholino)carbonyl)ethenyl)phenyl]sulfide

The title compound was prepared from Example 423A as described in
 Example 340G, substituting methyl isonipecotate with morpholine, followed by
 amination with ethyl isonipecotate as described in Example 396, and subsequent
 15 hydrolysis according to the procedure of Example 340H. ¹H NMR (400 MHz,
 DMSO-d₆) δ 1.5-1.64 (m, 2H), 1.67-1.75 (m, 1H), 1.87-1.96 (m, 1H), 2.49-2.57 (m,
 2H), 2.82-2.91 (m, 1H), 2.99-3.06 (m, 1H), 3.46-3.54 (m, 2H), 3.54-3.62 (m, 5H),
 3.63-3.72 (m, 3H), 6.87 (d, J=8 Hz, 1H), 7.06-7.13 (m, 2H), 7.16 (d, J=15 Hz, 1H),

7.25-7.36 (m, 2H), 7.66 (dd, $J_1=15$ Hz, $J_2=4$ Hz, 1H), 8.00 (d, $J=8$ Hz, 1H); MS (ESI) m/e 589 (M+H)⁺.




Example 424

(3-Hydroxyphenyl) [2,3-dichloro-4-(E-((4-morpholino)carbonyl)ethenyl)phenyl]sulfide

Example 310B was processed as described in Examples 310 and 405, substituting 3-methoxythiophenol for the thiol. ¹H NMR (DMSO-d₆, 300 MHz) δ


9.88 (s, 1H), 7.86 (d, $J=8.8$, 1H), 7.77 (d, $J=14.9$ Hz, 1H), 7.33 (dt, $J=7.4$ Hz, $J=1.0$ Hz, 1H), 7.24 (d, $J=14.8$ Hz, 1H), 6.96 (dd, $J=8.8$ Hz, $J=1.0$ Hz, 1H), 6.90 (dd, $J=8.8$ Hz, $J=1.0$ Hz, 1H), 6.89 (s, 1H), 6.79 (d, $J=8.5$ Hz, 1H), 3.67 (s, 2H), 3.58 (s, 6H); MS (APCI) m/z 410 (M+H)⁺, 427 (M+NH₄)⁺; 408 (M-H)⁻, 446 (M+Cl)⁻. Anal. calcd for C₁₉H₁₇Cl₂NO₃S 0.25 H₂O: C, 55.01; H, 4.25; N, 3.38. Found: C, 55.15; H, 4.25; N, 3.51.

385

Example 425

[3-(4-Butyroxyl)phenyl] [2,3-dichloro-4-(E-((4-morpholino)carbonyl)ethenyl)phenyl]sulfide


- 5 Example 424 was processed as described in Example 414 to provide the title compound as a white solid. ¹H NMR (DMSO-d₆, 300 MHz) δ 7.86 (d, *J*=8.8 Hz, 1H), 7.77 (d, *J*=15.3 Hz, 1H), 7.43 (dt, *J*=7.5 Hz, *J*=1.7 Hz, 1H), 7.24 (d, *J*=15.2 Hz, 1H), 7.11 (s, 1H), 6.79 (d, *J*=8.4 Hz, 1H), 6.73 (d, *J*=8.5 Hz, 1H), 4.02 (t, *J*=6.5 Hz, 2H), 3.67 (s, 2H), 3.58 (s, 6H), 2.37 (t, *J*=7.5 Hz, 2H), 1.95 (m, 2H); MS (APCI) *m/z* 410 (M+H⁺), 494 (M-H), 530 (M+Cl).
- 10

Example 426

(2-Hydroxyphenyl) [2,3-bis(trifluoromethyl)-4-(E-((4-morpholino)carbonyl)ethenyl)phenyl]sulfide

- 15 Example 401D was processed as described in Example 405 to provide the title compound as a white solid. ¹H NMR (DMSO-d₆, 300 MHz) δ 10.26 (s, 1H), 7.96 (d, *J*=8.5, 1H), 7.67 (m, 1H), 7.46 (dd, *J*=7.4 Hz, *J*=1.3 Hz, 1H), 7.38 (dt, *J*=7.5 Hz, *J*=1.3 Hz, 1H), 7.16 (d, *J*=15.2 Hz 1H), 7.13 (d, *J*=8.5 Hz, 1H), 7.00 (d, *J*=8.2 Hz,

1H), 6.90 (t, $J=7.4$ Hz, 2H), 3.65 (s, 2H), 3.57 (s, 6H). MS (APCI) m/z 478 (M+H)⁺, 495 (M+NH₄)⁺, 476 (M-H)⁻, 512 (M+Cl)⁻.



5


Example 427

(3-Hydroxyphenyl)[2,3-bis(trifluoromethyl)-4-(E-((4-
morpholino)carbonyl)ethenyl)phenyl]sulfide

Example 401C was processed as described in Example 401D, substituting 3-methoxythiophenol for the thiol, and in Example 405 to provide the title compound as

10 a white solid. ¹H NMR (DMSO-d₆, 300 MHz) δ 9.86 (s, 1H), 8.02 (d, $J=8.8$, 1H), 7.67 (m, 1H), 7.35 (d, $J=9.5$ Hz, 1H), 7.30 (dt, $J=7.8$ Hz, $J=0.7$ Hz, 1H), 7.19 (d, $J=15.2$ Hz 1H), 6.95 (d, $J=8.8$ Hz, 1H), 6.88 (d, $J=7.5$ Hz, 1H), 6.85 (s, 1H), 3.67 (s, 2H), 3.58 (s, 6H). MS (APCI) m/z 478 (M+H)⁺, 495 (M+NH₄)⁺, 476 (M-H)⁻, 512 (M+Cl)⁻.


15

Example 428

[3-(4-Carboxypiperidin-1-yl)phenyl] [2,3-ditrifluoromethyl-4-(E-((4-hydroxypiperidin-1-yl)carbonyl)ethenyl)phenyl] sulfide

The title compound was prepared by the procedures described in Example 412 substituting Example 384B with Example 423A. ¹H NMR (300 MHz, DMSO-d₆) δ

- 5 1.31 (m, 2H), 1.63 (m, 2H), 1.71 (m, 2H), 1.91 (m, 2H), 2.42 (m, 1H), 2.82 (t, J = 10.5 Hz, 2H), 3.16 (m, 1H), 3.31 (m, 1H), 3.70 (m, 3H), 3.93 (m, 2H), 6.88 (d, J = 7.1 Hz, 1H), 7.10 (d, J = 8.5 Hz, 1H), 7.14 (t, J = 7.1 Hz, 1H), 7.23 (d, J = 8.8 Hz, 1H), 7.28 (d, J = 8.2 Hz, 1H), 7.34 (d, J = 8.2 Hz, 1H), 7.60 (dq, J = 15.2, 4.5 Hz, 1H), 8.01 (d, J = 8.8 Hz, 1H); MS (APCI⁺) *m/z* 603 (M+H)⁺. Anal. calcd for C₂₈H₂₈F₆N₂O₄S 1.15
- 10 TFA: C, 49.60; H, 4.00; N, 3.82. Found: C, 49.65; H, 3.80; N, 3.81.



Example 429

[3-(4-Carboxypiperidin-1-yl)phenyl] [2,3-ditrifluoromethyl-4-(E-((1,2,5,6-tetrahydropyridin-1-yl)carbonyl)ethenyl)phenyl] sulfide


The title compound was prepared by the procedures described in Example 412 substituting Example 384B with Example 423A, and substituting 4-hydroxypiperidine

with 1,2,5,6-tetrahydropyridine. ¹H NMR (300 MHz, DMSO-d₆) δ 1.63 (m, 2H), 1.90 (m, 2H), 2.12 (m, 2H), 2.43 (m, 1H), 2.81 (t, J = 10.5 Hz, 2H), 3.75 (m, 4H),

- 20 4.01 ((s, 1H), 4.15 (s, 1H), 5.73 (m, 1H), 5.84 (m, 1H), 6.87 (d, J = 7.5 Hz, 1H), 7.10

(m, 2H), 7.30 (m, 3H), 7.62 (m, 1H), 8.01 (t, $J = 6.5$ Hz, 1H); MS (APCI⁺) m/z 585

(M+H)⁺. Anal. calcd for C₂₈H₂₆F₆N₂O₃S 0.1 TFA: C, 56.83; H, 4.41; N, 4.70. Found: C, 56.91; H, 4.44; N, 4.60.




Example 430

2-((4-Carboxy)butyloxy)phenyl [2,3-dichloro-4-(E-((4-
morpholino)carbonyl)ethenyl)phenyl]sulfide

Example 405 was processed as described in Example 414, substituting ethyl 5-

- 10 bromovalerate for the alkyl halide. ¹H NMR (DMSO-d₆, 300 MHz) δ 7.79 (d, $J=8.5$, 1H), 7.77 (d, $J=15.3$ Hz, 1H), 7.51 (dt, $J=8.2$ Hz, $J=1.7$ Hz, 1H), 7.48 (dd, $J=7.5$ Hz, $J=1.7$ Hz, 1H), 7.20 (dd, $J=7.5$ Hz, $J=1.7$ Hz, 1H), 7.20 (d, $J=15.6$ Hz, 1H), 7.05 (dt, $J=7.5$ Hz, $J=1.0$ Hz, 1H), 6.63 (d, $J=8.5$ Hz, 1H), 3.99 (t, $J=6.1$ Hz, 2H), 3.65 (s, 2H), 3.58 (s, 6H), 2.10 (t, $J=7.1$ Hz, 2H), 1.56 (m, 2H), 1.39 (m, 2H). MS (APCI) m/z 510
- 15 (M+H)⁺. Anal. calcd for C₂₄H₂₅Cl₂NO₃S 0.75 H₂O: C, 55.02; H, 5.10; N, 2.67. Found: C, 54.72; H, 4.82; N, 2.77.




Example 431

(2-Glycoxyphenyl) [2,3-bis(trifluoromethyl)-4-(E-((4-morpholino)carbonyl)ethenyl)phenyl]sulfide

Example 426 was processed as detailed in Example 414, substituting ethyl

- 5 bromoacetate for the alkyl bromide. ^1H NMR (DMSO- d_6 , 300 MHz) δ 7.94 (d, $J=8.4$, 1H), 7.66 (m, 1H), 7.50 (m, 1H), 7.47 (d, $J=7.4$ Hz, 1H), 7.28 (d, $J=8.5$ Hz, 1H), 7.16 (d, $J=14.9$ Hz, 1H), 7.07 (m, 2H), 4.74 (s, 2H), 3.65 (s, 2H), 3.57 (s, 6H); MS (APCI) m/z 536 (M+H) $^+$, 553 (M+NH $_4$) $^+$, 534 (M-H) $^+$. Anal. calcd for C $_{23}$ H $_{19}$ F $_6$ NO $_5$ S: C, 51.59; H, 3.58; N, 2.62. Found: C, 51.31; H, 3.63; N, 2.33.


10

Example 432

(2-(4-Butyroxyl)phenyl) [2,3-bis(trifluoromethyl)-4-(E-((4-morpholino)carbonyl)ethenyl)phenyl]sulfide


- 15 Example 426 was processed as described in Example 414 to provide the title compound as a white solid. ^1H NMR (DMSO- d_6 , 300 MHz) δ 7.92 (d, $J=8.5$, 1H), 7.65 (m, 1H), 7.59 (d, $J=7.8$ Hz, $J=1.7$ Hz, 1H), 7.50 (t, $J=8.2$ Hz, 1H), 7.15 (d, $J=8.2$ Hz, 1H), 7.12 (d, $J=15.6$ Hz, 1H), 7.09 (d, $J=8.8$ Hz, 1H), 7.07 (t, $J=7.2$ Hz, 1H), 3.92 (t, $J=6.1$ Hz, 2H), 3.65 (s, 2H), 3.57 (s, 6H), 1.99 (t, $J=7.1$ Hz, 2H), 1.63 (m, 2H); MS (APCI) m/z 562 (M-H) $^+$, 598 (M+Cl) $^+$.
- 20

390

Example 433

[3-(4-Carboxypiperidin-1-yl)phenyl] [2,3-ditrifluoromethyl-4-(E-((bis-(2-ethoxyethyl)amino)carbonyl)ethenyl)phenyl] sulfide


- 5 The title compound was prepared by the procedures described in Example 412 substituting Example 384B with Example 423A, and substituting 4-hydroxypiperidine with bis (2-ethoxyethyl)amine. ^1H NMR (300 MHz, DMSO- d_6) δ 0.99 (t, J = 6.8 Hz, 3H), 1.09 (t, J = 6.8 Hz, 3H), 1.63 (m, 2H), 1.90 (m, 2H), 2.44 (m, 1H), 2.82 (t, J = 10.8 Hz, 2H), 3.40 (m, 4H), 3.50 (m, 6H), 3.68 (m, 4H), 6.88 (d, J = 7.5 Hz, 1H), 7.11 (m, 3H), 7.32 (m, 2H), 7.62 (dq, J = 15.2, 4.5 Hz, 1H), 7.94 (d, J = 8.5 Hz, 1H); MS (APCI $^+$) m/z 663 (M+H) $^+$. Anal. calcd for $\text{C}_{31}\text{H}_{36}\text{F}_6\text{N}_2\text{O}_5\text{S}$ 0.7 TFA: C, 52.41; H, 4.98; N, 3.77. Found: C, 52.38; H, 4.94; N, 3.68.
- 10

Example 434

[3-(4-Carboxypiperidin-1-yl)phenyl] [2,3-bis-(trifluoromethyl)-4-(E-((bis-(2-hydroxypropyl)amino)carbonyl)ethenyl)phenyl] sulfide

15

The title compound was prepared by the procedures described in Example 412 substituting Example 384B with Example 423A, and substituting 4-hydroxypiperidine with diisopropanolamine. ^1H NMR (300 MHz, DMSO- d_6) δ 1.04 (m, 6H), 1.63 (m, 2H), 1.90 (m, 2H), 2.42 (m, 1H), 2.83 (t, J = 10.5 Hz, 2H), 3.04 (m, 1H), 3.26 (m, 1H), 3.45 (m, 2H), 3.67 (m, 2H), 3.75 (m, 1H), 3.90 (m, 1H), 6.90 (d, J = 7.5 Hz, 1H), 7.11 (m, 3H), 7.28 (d, J = 8.5 Hz, 1H), 7.33 (t, J = 8.2 Hz, 1H), 7.63 (m, 1H), 7.95 (dd, J = 8.5, 2.1 Hz, 1H); MS (APCI $^+$) m/z 635 (M+H) $^+$. Anal. calcd for $\text{C}_{29}\text{H}_{32}\text{F}_6\text{N}_2\text{O}_3\text{S}$ 1.5 TFA: C, 47.71; H, 4.19; N, 3.48. Found: C, 47.52; H, 4.28; N, 3.40.




Example 435

[3-(4-Carboxypiperidin-1-yl)phenyl] [2,3-bis-(trifluoromethyl)-4-(E-((piperazin-1-yl)carbonyl)ethenyl)phenyl] sulfide

The title compound was prepared by the procedures described in Example 412 substituting Example 384B with Example 423A, and substituting 4-hydroxypiperidine with 1-(1,2,3,4-tetrahydrofuroyl)piperazine. ^1H NMR (300 MHz, DMSO- d_6) δ 1.62 (m, 2H), 1.88 (m, 2H), 2.43 (m, 1H), 2.82 (t, J = 10.5 Hz, 2H), 3.15 (br s, 4H), 3.71 (m, 2H), 3.75 (m, 2H), 3.86 (m, 2H), 6.87 (d, J = 7.5 Hz, 1H), 7.10 (d, J = 8.5 Hz, 1H), 7.12 (d, J = 7.5 Hz, 1H), 7.20 (d, J = 15.2 Hz, 1H), 7.28 (d, J = 8.5 Hz, 1H), 7.35

(d, $J = 7.8$ Hz, 1H), 7.70 (dt, $J = 15.2, 4.5$ Hz, 1H), 7.98 (d, $J = 8.5$ Hz, 1H), 8.85 (br s, 1H); MS (APCI⁺) m/z 588 (M+H)⁺. Anal. calcd for C₂₇H₂₇F₆N₃O₃S 3.3 TFA: C, 41.87; H, 3.17; N, 4.36. Found: C, 41.78; H, 3.26; N, 4.43.




5

Example 436

(3-(4-Butyroxyl)phenyl)[2,3-bis(trifluoromethyl)-4-(E-((4-morpholino)carbonyl)ethenyl)phenyl]sulfide

Example 427 was processed as described in Example 414 to provide the title compound as a white solid. ¹H NMR (DMSO-d₆, 300 MHz) δ 8.02 (d, $J = 8.5$, 1H), 7.65 (m, 1H), 7.40 (t, $J = 7.8$ Hz, 1H), 7.33 (d, $J = 8.8$ Hz, 1H), 7.19 (d, $J = 14.9$ Hz, 1H), 7.09 (m, 3H), 4.02 (t, $J = 6.4$ Hz, 2H), 3.67 (s, 2H), 3.58 (s, 6H), 2.37 (t, $J = 7.1$ Hz, 2H), 1.95 (m, 2H); MS (APCI) m/z 564 (M+H)⁺, 562 (M-H)⁻, 598 (M+Cl)⁻.

10




15


Example 437

[2-(3-Carboxypiperidin-1-yl)phenyl][2,3-dichloro-4-(E-[(3-(2-pyrrolidinon-1-yl)propylaminocarbonyl)ethenyl]phenyl] sulfide

393

Example 437AN-Benzyloxycarbonyl nipecotic acid

- To a solution of nipecotic acid (10 g, 63.6 mmol) in 1 N NaOH (2.5 g in 64 ml
- 5 water, 63.6 mmol) at 0 °C was alternately added benzyloxycarbonyl chloride (10.9 mL, 76.5 mmol) in diethyl ether (50 mL) and 1 N NaOH (5 g in 128 ml water, 127.2 mmol) in five portions. The reaction mixture was stirred at 0 °C for 2 h, and at ambient temperature for 24 h. Then this was made acidic with 10% HCl and the solid
- 10 (18.9 g, 113%). MS (ESI) m/e 264 (M+H)⁺.

Example 437Btert-Butyl N-benzyloxycarbonyl nipecotate

- 15 A solution of Example A (18 g, 62 mmol) in THF (250 mL, 0.25 M) was treated with trichloroacetimidate (28 mL, 155 mmol) and BF₃·Et₂O (18 mL, 1 mL/g) at ambient temperature. After 18 h the reaction mixture was quenched with solid

- NaHCO₃ followed by water and stirred vigorously. Then the solvent was removed, and partitioned with ethyl acetate (250 mL). The organic layer was separated and washed with brine (3x80 mL), dried (Na₂SO₄) and evaporated to dryness under reduced pressure to obtain the crude product. The title compound (19.2 g, 96%) was
- 5 obtained by flash chromatography on silica gel eluting with 20% acetone:hexane. MS (ESI) *m/e* 320 (M+H)⁺.




Example 437C

10

tert-Butyl nipecotate

Example 437B (19 g, 59.5 mmol) was treated with 10% Pd on carbon (2 g, 10 wt %) in ethanol (237 mL, 0.25 M) to obtain the title compound (10.4 g, 94%). MS (ESI) *m/e* 186 (M+H)⁺.



15


Example 437D

2-Nitro-(3-(tert-butoxycarbonyl)piperidin-1-yl)benzene

To a solution of Example 437C (10.4 g, 56.1 mmol) in toluene (112 mL) was added 2-fluoronitrobenzene (6.0 ml, 56 mmol) and CsF (852 mg, 5.6 mmol). The

reaction mixture was stirred under reflux conditions for 18 h, and allowed to cool to ambient temperature. The mixture was diluted with ethyl acetate (100 ml), washed with 10% HCl (2x50 ml), followed by brine (3x100 ml), then dried (Na_2SO_4) and evaporated in vacuo to obtain the title compound (16.5 g, 94%). MS (ESI) m/e 307


5 (M+H)⁺.



Example 437E

2-Amino-3-(tert-butyloxycarbonyl)piperidin-1-ylbenzene


10 Example 437E (16.4 g, 53.5 mmol) was treated with 10% Pd on carbon (1.65 g, 10 wt %) in ethanol (215 ml, 0.5 M) at ambient temperature for 2 hours. The reaction mixture was filtered through celite and the filtrate was evaporated to dryness under reduced pressure to obtain the title compound (13.45 g, 91%). MS (ESI) m/e 277 (M+H)⁺.



Example 437F

2-Iodo-3-(tert-butyloxycarbonyl)piperidin-1-ylbenzene

- Example 437E was dissolved in 3 N H₂SO₄ (195 mL, 0.25 M), cooled to 0 °C and treated with NaNO₂ (3.35 g, 48.6 mmol) in water (20 mL). After 30 minutes at 0 °C potassium iodide (12.01 g, 72.8 mmol) and urea (583 mg, 9.7 mmol) in water (10 mL) were added and stirred for 1 h. The reaction mixture was quenched with 10% NaHCO₃ (50 mL) and partitioned with ethyl acetate (450 mL). The organic layer was separated and washed with 10% NaHCO₃ (2x100 mL), brine (2x100mL), dried (Na₂SO₄) and evaporated to dryness under reduced pressure. The title compound (17.2 g, 91%) was obtained by flash chromatography on silica gel eluting with 10% acetone:hexane. ¹H NMR (400 MHz, DMSO-d₆) δ 1.39 (s, 9H), 6.85 (tt, J₁=1.5 Hz, J₂=7.5 Hz, 1H), 7.14 (dd, J₁=1.5 Hz, J₂=7.5 Hz, 1H), 7.37 (tt, J₁=1.5 Hz, J₂=7.5 Hz, 1H), 7.84 (dd, J₁=1.5 Hz, J₂=7.5 Hz, 1H); MS (ESI) *m/e* 388 (M+H)⁺.




Example 437G

- [2-(3-*tert*-Butyloxycarbonyl)piperidin-1-yl]phenyl [2,3-dichloro-4-(*E*-(2-methoxycarbonyl)ethenyl)phenyl] sulfide

Example 437F was converted to the corresponding triisopropylsilyl thiol analogue by the method describe d for the preparation of Example 340B. Then this intermediate was reacted with Example 340E (2.94 g, 7.75 mmol) at -20 °C as


described in Example 340F to obtain the title compound (2.5 g, 63%). MS (ESI) *m/e* 522, 524 (M+H)⁺.



Example 437H

[2-(3-*tert*-Butyloxycarbonyl)piperidin-1-yl]phenyl [2,3-dichloro-4-(*E*-(2-carboxy)ethenyl)phenyl] sulfide

Using the procedure for Example 340H, Example 437H was hydrolyzed to the title compound. MS (ESI) *m/e* 508, 510 (M+H)⁺.




Example 437I

[2-(3-Carboxypiperidin-1-yl)phenyl] [2,3-dichloro-4-(*E*-[(3-(2-pyrrolidinon-1-yl)propyl)aminocarbonyl]ethenyl)phenyl] sulfide


The title compound (66 mg) was prepared from Example 437H (90 mg, 0.177 mmol), using the procedures described in Example 340G substituting methyl isonipecotate with 3-aminopropyl-2-pyrrolidinone followed by treatment with 20%

TFA in methylene chloride as described in Example 412C. ^1H NMR (500 MHz, DMSO- d_6) δ 1.27-2.43 (m, 2H), 2.60-2.72 (m, 3H), 2.84 (m, 3H), 2.17-2.30 (m, 3H), 2.62-2.73 (m, 2H), 3.08-3.23 (m, 5H), 3.29-3.38 (m, 3H), 6.62 (d, $J=15$ Hz, 1H), 6.92 (d, $J=8.5$ Hz, 1H), 7.12 (t, $J=7.5$ Hz, 1H), 7.41 (m, 1H), 7.58 (d, $J=8.75$ Hz, 1H), 7.70 (d, $J=15$ Hz, 1H), 8.21 (t, $J=5$ Hz, 1H); MS (ESI) m/e 576, 578 ($M+H$) $^+$.



Example 438

[2-(3-Carboxypiperidin-1-yl)phenyl] [2,3-bis(trifluoromethyl)-4-(E-[(3-(2-pyrrolidinon-1-yl)propylaminocarbonyl)ethenyl]phenyl) sulfide




Example 438A

[2-(3-tert-Butyloxycarbonyl)piperidin-1-yl)phenyl] [2,3-bis(trifluoromethyl)-4-(E-(2-carboxy)ethenyl)phenyl] sulfide

The title compound (445 mg, 71%) was prepared from the reaction of Example 401C (500 mg, 1.08 mmol) with Example 437F, using the procedures

described in Example 437G followed by hydrolysis as described in Example 340G.

MS (ESI) m/e 604 (M+H)⁺.




Example 438B

[2-(3-Carboxypiperidin-1-yl)phenyl] [2,3-bis(trifluoromethyl)-4-(E-[(3-(2-pyrrolidinon-1-yl)propyl)aminocarbonyl]ethenyl)phenyl] sulfide


The title compound was prepared from Example 438A (110 mg, 0.191 mmol), using the procedures described in Example 340G substituting methyl isonipecotate

- 10 with 3-aminopropyl-2-pyrrolidinone followed by treatment with 20% TFA in methylene chloride as described in Example 412C. ¹H NMR (500 MHz, DMSO-d₆)
- δ 1.11-1.21 (m, 1H), 1.28-1.38 (m, 1H), 1.60-1.70 (m, 3H), 1.79-1.86 (m, 1H), 1.87-1.94 (m, 2H), 2.05-2.12 (m, 1H), 2.00 (t, J=7.5 Hz, 2H), 2.58-2.66 (m, 2H), 2.96-3.01 (m, 1H), 3.11-3.18 (m, 2H), 3.19 (t, J=6.25 Hz, 2H), 3.26 (m, 1H), 3.32 (t, J=6.25 Hz,
- 15 2H), 6.46 (d, J=15 Hz, 1H), 7.15 (t, J=7.5 Hz, 1H), 7.33 (m, 2H), 7.43 (m, 1H), 7.62 (m, 1H), 7.75 (d, J=7.5 Hz, 1H), 8.22 (t, J=5 Hz, 1H); MS (ESI) m/e 644 (M+H)⁺.

Example 439

[2-(3-Carboxypiperidin-1-yl)phenyl] [2,3-dichloro-4-(E-((4-(2-hydroxyethyl)piperazin-1-yl)carbonyl)ethenyl)phenyl] sulfide


5 The title compound was prepared from Example 437H (90 mg, 0.177 mmol), using the procedures described in Example 340G substituting methyl isonipecotate with 1-(2-hydroxyethyl)piperazine followed by treatment with 20% TFA in methylene chloride as described in Example 412C. ¹H NMR (500 MHz, DMSO-d₆) δ 1.31-1.45 (m, 2H), 1.67-1.74 (m, 1H), 1.86-1.92 (m, 1H), 2.24-2.31 (m, 1H), 2.66 (t, J=10 Hz, 1H), 2.73 (t, J=10Hz, 1H), 3.02-3.24 (m, 5H), 3.33-3.38 (m, 1H), 3.52 (m, 3H), 3.75 (t, J=5 Hz, 2H), 4.31-4.60 (m, 3H), 6.90 (d, J=9 Hz, 1H), 7.11 (t, J=7.5 Hz, 1H), 7.21 (d, J=7.5 Hz, 1H), 7.25 (d, J=7.5 Hz, 1H), 7.30 (d, J=15 Hz, 1H), 7.42 (t, J=7.5Hz, 1H), 7.83 (d, J=15 Hz, 1H), 7.85 (d, 9 Hz, 1H); MS (ESI) *m/e* 564, 566 (M+H)⁺.

Example 440

[2-(3-Carboxypiperidin-1-yl)phenyl] [2,3-bis(trifluoromethyl)-4-(E-((1,2,5,6-tetrahydropyridin-1-yl)carbonyl)ethenyl)phenyl] sulfide

The title compound was prepared from Example 438A (110 mg, 0.191 mmol), using the procedures described in Example 340G substituting methyl isonipecotate

- 5 with 1,2,5,6-tetrahydropyridine followed by treatment with 20% TFA in methylene chloride as described in Example 412C. ¹H NMR (500 MHz, DMSO-d₆) δ 1.19-1.27 (m, 1H), 1.30-1.40 (m, 1H), 1.64-1.70 (m, 1H), 1.80-1.86 (m, 1H), 2.04-2.18 (m, 2H), 2.60 (t, J=10 Hz, 1H), 2.68 (t, J=10 Hz, 1H), 3.03 (br d, J=10 Hz, 1H), 3.23 (br d, J=10 Hz, 1H), 3.60-3.74 (m, 2H), 3.91-4.20 (m, 3H), 5.68-5.74 (m, 1H), 5.80-5.90 (m, 1H), 7.06-7.19 (m, 2H), 7.20-7.28 (m, 2H), 7.36 (m, 1H), 7.44 (t, J=7.5 Hz, 1H), 7.62-7.71 (m, 1H), 7.94-8.04 (m, 1H); MS (ESI) m/e 585 (M+H)⁺.
- 10




Example 441

- 15 [2-(3-Carboxypiperidin-1-yl)phenyl] [2,3-bis(trifluoromethyl)-4-(E-((4-(2-hydroxyethyl)piperazin-1-yl)carbonyl)ethenyl)phenyl] sulfide

The title compound was prepared from Example 438A (110 mg, 0.191 mmol), using the procedures described in Example 340G substituting methyl isonipecotate with 1-(2-hydroxyethyl)piperazine followed by treatment with 20% TFA in methylene

chloride as described in Example 412C. ¹H NMR (500 MHz, DMSO-d₆) δ 1.17-1.27 (m, 1H), 1.17-1.27 (m, 1H), 1.31-1.41 (m, 1H), 1.64-1.71 (m, 1H), 1.80-1.88 (m, 1H), 2.07-2.15 (m, 1H), 2.61 (t, J=10 Hz, 1H), 2.09 (t, J=10 Hz, 1H), 2.91-3.13 (m, 3H), 3.18-3.28 (m, 3H), 3.44-3.58 (m, 3H), 3.75 (t, J=5 Hz, 2H), 4.29-4.58 (m, 3H), 7.15 (d, J=7.5 Hz, 1H), 7.19 (d, J=10 Hz, 1H), 7.28 (d, J=7.5 Hz, 2H), 7.37 (d, J=7.5 Hz, 1H), 7.45 (t, J=7.5 Hz, 1H), 7.69-7.77 (m, 1H), 7.98 (d, J=7.5 Hz, 1H), 9.77 (br s, 1H); MS (ESI) *m/e* 632 (M+H)⁺.




Example 442

[2-(3-Carboxypiperidin-1-yl)phenyl] [2,3-bis(trifluoromethyl)-4-((E)-((4-(2-hydroxyethoxy)ethyl)piperazin-1-yl)carbonyl)ethenyl)phenyl] sulfide

The title compound was prepared from Example 438A (110 mg, 0.191 mmol), using the procedures described in Example 340G substituting methyl isonipecotate with 1-[2-(2-hydroxyethoxy)ethyl]piperazine followed by treatment with 20% TFA in methylene chloride as described in Example 412C. ¹H NMR (500 MHz, DMSO-d₆) δ 1.17-1.27 (m, 1H), 1.31-1.40 (m, 1H), 1.64-1.70 (m, 1H), 1.80-1.86 (m, 1H), 2.08-2.16 (m, 1H), 2.61 (t, J=10 Hz, 1H), 2.68 (t, J=10 Hz, 1H), 3.03 (br d J=10 Hz, 1H), 3.06-3.18 (m, 2H), 3.25 (br d, J=10 Hz, 1H), 3.33 (m, 2H), 3.42-3.51 (m, 2H), 3.53-

3.60 (m, 2H), 3.70-3.80 (m, 5H), 4.32-4.56 (m, 3H), 7.13-7.21 (m, 2H), 7.27 (d, $J=9$ Hz, 2H), 7.35 (d, $J=7.5$ Hz, 1H), 7.44 (t, $J=7.5$ Hz, 1H), 7.68-7.77 (m, 1H), 7.44-8.01 (m, 1H), 9.81 (br s, 1H); MS (ESI) m/e 676 (M+H)⁺.



Example 443

(3-(3-Propionyloxy)phenyl) [2,3-dichloro-4-(E-((4-morpholino)carbonyl)ethenyl)phenyl]sulfide

β -Propiolactone (50 μ L, 0.75 mmol) was added to a mixture of Example 405 (308 mg, 0.75 mmol), potassium *tert*-butoxide (750 mL, 1 M in THF), and THF (1.0 mL). After 18 h, the reaction was diluted with EtOAc, washed with 1 M aqueous HCl, washed with brine, dried (MgSO₄), filtered, and concentrated. Purification by preparative HPLC provided the title compound (72 mg, 20%) as a white solid. ¹H NMR (DMSO-d₆, 300 MHz) δ 7.80 (d, $J=8.4$ Hz, 1H), 7.78 (d, $J=15.8$ Hz, 1H), 7.52 (dt, $J=8.8$ Hz, $J=1.7$ Hz, 1H), 7.46 (dd, $J=7.8$ Hz, $J=17$ Hz, 1H), 7.23 (d, $J=9.1$ Hz, 1H), 7.22 (d, $J=15.3$ Hz, 1H), 7.08 (t, $J=7.4$ Hz, 1H), 6.58 (d, $J=8.5$ Hz, 1H), 4.22 (m, 2H), 4.05 (m, 2H), 3.66 (s, 2H), 3.58 (s, 6H); MS (APCI) m/z 482 (M+H)⁺; 480 (M-H)⁻.

Compounds that antagonize the interaction between ICAM-1 and LFA-1 can be identified, and their activities quantitated, using both biochemical and cell-based adhesion assays. A primary biochemical assay measures the ability of the compound in question to block the interaction between the integrin LFA-1 and its adhesion partner ICAM-1, as described below:

ICAM-1 / LFA-1 Biochemical Interaction Assay

In the biochemical assay, 100 μ L of anti-LFA-1 antibody (ICOS Corporation) at a concentration of 5 μ g/ml in Dulbecco's phosphate-buffered saline (D-PBS) is used to coat wells of a 96-well microtiter plate overnight at 4°C. The wells are then washed twice with wash buffer (D-PBS w/o Ca^{++} or Mg^{++} , 0.05% Tween 20) and blocked by addition of 200 μ L of D-PBS, 5% fish skin gelatin. Recombinant LFA-1 (100 μ L of 0.7 μ g/ml, ICOS Corporation) in D-PBS is then added to each well. Incubation continues for 1 hour at room temperature and the wells are washed twice with wash buffer. Serial dilutions of compounds being assayed as ICAM-1/LFA-1 antagonists, prepared as 10 mM stock solutions in dimethyl sulfoxide (DMSO), are diluted in D-PBS, 2mM MgCl_2 , 1% fish skin gelatin and 50 μ L of each dilution added to duplicate wells. This is followed by addition of 50 μ L of 0.8 μ g/ml biotinylated recombinant ICAM-1/Ig (ICOS Corporation) to the wells and the plates are incubated at room temperature for 1 hour. The wells are then washed twice with wash buffer and 100 μ L of Europium-labeled Streptavidin (Wallac Oy) diluted 1:100 in Delfia assay

buffer (Wallac Oy) are added to the wells. Incubation proceeds for 1 hour at room temperature. The wells are washed eight times with wash buffer and 100 μ L of enhancement solution (Wallac Oy, cat. No. 1244-105) are added to each well. Incubation proceeds for 5 minutes with constant mixing. Time-resolved fluorimetry measurements are made using the Victor 1420 Multilabel Counter (Wallac Oy) and the percent inhibition of each candidate compound is calculated using the following equation:

10

$$\% \text{ inhibition} = 100 \times \left\{ 1 - \frac{\text{average OD w/ compound minus background}}{\text{average OD w/o compound minus background}} \right\}$$

where "background" refers to wells that are not coated with anti-LFA-1 antibody.

Compounds of the present invention exhibit inhibitory activity in the above assay. % inhibition at 4 μ M was demonstrated.

Biologically relevant activity of the compounds in this invention is

- 5 confirmed using a cell-based adhesion assay, which measures their ability to block the adherence of JY-8 cells (a human EBV-transformed B cell line expressing LFA-1 on its surface) to immobilized ICAM-1, as follows:

ICAM-1 / JY-8 cell adhesion assay

- 10 For measurement of inhibitory activity in the cell-based adhesion assay, 96-well microtiter plates are coated with 70 μ L of recombinant ICAM-1/Ig (ICOS Corporation) at a concentration of 5 μ g/mL in D-PBS w/o Ca^{++} or Mg^{++} overnight at 4°C. The wells are then washed twice with D-PBS and blocked by addition of 200 μ L of D-PBS, 5% fish skin gelatin by incubation for 1 hour at room temperature.
- 15 Fluorescent tagged JY-8 cells (a human EBV-transformed B cell line expressing LFA-1 on its surface; 50 μ L at 2×10^6 cells/ml in RPMI 1640/1% fetal bovine serum) are added to the wells. For fluorescent labeling of JY-8 cells, 5×10^6 cells washed once in RPMI 1640 are resuspended in 1 mL of RPMI 1640 containing 2 μ M Calcein AM (MolecularProbes), are incubated at 37°C for 30 minutes and
- 20 washed once with RPMI-1640/ 1% fetal bovine serum. Dilutions of compounds to be assayed for ICAM-1/LFA-1 antagonistic activity are prepared in RPMI-1640/ 1% fetal bovine serum from 10mM stock solutions in DMSO and 50 μ L are added to

duplicate wells. Microtiter plates are incubated for 45 minutes at room temperature and the wells are washed gently once with RPMI-1640/ 1% fetal bovine serum.

Fluorescent intensity is measured in a fluorescent plate reader with an excitation wavelength at 485 nM and an emission wavelength at 530 nM. The percent

- 5 inhibition of a candidate compound at a given concentration is calculated using the following equation:

$$\% \text{ inhibition} = 100 \times \left\{ 1 - \frac{\text{average OD w/ compound}}{\text{average OD w/o compound}} \right\}$$

- 10 and these concentration/inhibition data are used to generate dose response curves, from which IC_{50} values are derived.

Compounds of the present invention exhibit blocking activity in the above assay. Inhibition at 4 μ M was demonstrated.

15

Compounds of the present invention have been demonstrated to act via interaction with the integrin LFA-1, specifically by binding to the interaction domain (I-domain), which is known to be critical for the adhesion of LFA-1 to a variety of cell adhesion molecules. As such, it is expected that these compounds

20 should block the interaction of LFA-1 with other CAM's. This has in fact been demonstrated for the case of ICAM-3. Compounds of the present invention may be

evaluated for their ability to block the adhesion of JY-8 cells (a human EBV-transformed B cell line expressing LFA-1 on its surface) to immobilized ICAM-3, as follows:

5 ICAM-3 / JY-8 cell adhesion assay

For measurement of inhibitory activity in the cell-based adhesion assay, 96-well microtiter plates are coated with 50 μ L of recombinant ICAM-3/Ig (ICOS Corporation) at a concentration of 10 μ g/mL in D-PBS w/o Ca^{++} or Mg^{++} overnight at 4°C. The wells are then washed twice with D-PBS, blocked by addition of 100 μ L of D-PBS, 1% bovine serum albumin (BSA) by incubation for 1 hour at room temperature, and washed once with RPMI-1640 / 5% heat-inactivated fetal bovine serum (adhesion buffer). Dilutions of compounds to be assayed for ICAM-3/LFA-1 antagonistic activity are prepared in adhesion buffer from 10 mM stock solutions in DMSO and 100 μ L are added to duplicate wells. JY-8 cells (a human EBV-transformed B cell line expressing LFA-1 on its surface; 100 μ L at 0.75×10^6 cells/ml in adhesion buffer) are then added to the wells. Microtiter plates are incubated for 30 minutes at room temperature; the adherent cells are then fixed with 50 μ L of 14% glutaraldehyde/D-PBS and incubated for an additional 90 minutes. The wells are washed gently with dH_2O ; 50 μ L of dH_2O is added, followed by 50 μ L of 1% crystal violet. After 5 minutes the plates are washed 3X with dH_2O ; 75 μ L of dH_2O and 225 μ L of 95% EtOH are added to each well to extract the crystal violet from the cells. Absorbance is measured at 570 nm in an ELISA plate reader.

The percent inhibition of a candidate compound is calculated using the following equation:

$$\% \text{ inhibition} = 100 \times \left\{ 1 - \frac{\text{average OD w/ compound}}{\text{average OD w/o compound}} \right\}$$

5

Compounds of the present invention exhibit blocking activity in the above assay. 100 % inhibition at 0.6 μM was demonstrated.

001230-56214560

Additional JY-8 (ICAM/LFA-1) Cell Adhesion Assay Protocol

Reagents

- ICAM-1/Ig, ICOS
- 5 • D-PBS, Dulbecco's w/o Ca & Mg
- D-PBS, Dulbecco's w/ Ca & Mg
- Blocking Solution: 1% non-fat dried milk in PBS w/o Ca & Mg
- RPMI 1640 media
- RPMI 1640 media with 1%FBS (RPMI-1%FBS)
- 10 • RPMI 1640 media with 50%FBS (RPMI-50%FBS)
- 1 mM Calcein AM, Molecular Probes, cat. C-1430 or C-3099
- DMSO
- JY-8 cells

15 Procedure

1. Coat plate (70ul/well) with 5ug/ml in D-PBS w/ Ca & Mg of ICAM-1/Ig. Cover and incubate overnight at 4°C.
2. Make compound and control dilutions using RPMI-1%FBS and RPMI-50%FBS as the diluents.
- 20 3. Decant ICAM-1/Ig coated plate(s), and wash 3X with D-PBS w/o Ca & Mg.
4. Block entire plate(s) with 150ul/well of Blocking solution. Cover and incubate for approximately 1 hour at room temperature.
5. Count the number of viable JY-8 cells using standard methodology. Need approximately 10-15 x10E6 cells per 96mw tray.

6. Wash cells 1X in RPMI 1640 media without serum - centrifuging for 5 minutes at approximately 1400rpms. Remove supernate and resuspend cell pellet to 5x10E6 cells per ml in RPMI 1640 media without serum.
7. Add 2ul of 1mM Calcein AM for every 1ml of cell suspension. Mix. Incubate for 30-60 minutes at 37 degrees C in a CO2 incubator (keeping cap of centrifuge tube loose for gas exchange).
8. Add approximately 10mls of RPMI-1%FBS, aliquot into two equal pools and centrifuge for 5 minutes at 1400rpms.
9. Remove supernate from each pool and resuspend each cell pellet to 2x10E6 cell per ml with RPMI-1%FBS or RPMI-50%FBS.
10. Decant blocked 96mw plate(s) and wash 3X with D-PBS w/o Ca & Mg.
11. Add 50ul/well of each compound dilution or control. Add 50ul of Calcein labeled JY-8 cells to all wells. Centrifuge plate(s) briefly (2-5 seconds) at 100-150rpms. Cover and incubate for 30-60 minutes at 37 degrees C.
12. Gently wash wells 1X with approximately 150ul per well of PBS w/Ca & Mg. Remove all liquid from wells.
13. Read absorbance using reader with an excitation of 485/20 and an emission of 530/25.
14. Calculate % inhibition using the following equation:

$$\% \text{ inhibition} = 100 \times \left\{ 1 - \frac{\text{average OD w/ compound}}{\text{average OD w/o compound}} \right\}$$

The ability of the compounds of this invention to treat arthritis can be demonstrated in a murine collagen-induced arthritis model according to the method of Kakimoto, et al., *Cell Immunol* 142: 326-337, 1992, in a rat collagen-induced arthritis model according to the method of Knoerzer, et al., *Toxicol Pathol* 25:13-19, 1997, in a rat adjuvant arthritis model according to the method of Halloran, et al., *Arthritis Rheum* 39: 810-819, 1996, in a rat streptococcal cell wall-induced arthritis model according to the method of Schimmer, et al., *J Immunol* 160: 1466-1477, 1998, or in a SCID-mouse human rheumatoid arthritis model according to the method of Oppenheimer-Marks et al., *J Clin Invest* 101: 1261-1272, 1998.

The ability of the compounds of this invention to treat Lyme arthritis can be
15 demonstrated according to the method of Gross et al., *Science* 281, 703-706, 1998.

The ability of compounds of this invention to treat asthma can be demonstrated in a murine allergic asthma model according to the method of Wegner et al., *Science* 247:456-459, 1990, or in a murine non-allergic asthma model according to the method of Bloemen et al., *Am J Respir Crit Care Med* 153:521-529, 1996.

The ability of compounds of this invention to treat inflammatory lung injury can be demonstrated in a murine oxygen-induced lung injury model according to the method of Wegner et al., *Lung* 170:267-279, 1992, in a murine immune complex-induced lung injury model according to the method of Mulligan et al., *J Immunol* 154:1350-1363, 1995, or in a murine acid-induced lung injury model according to the method of Nagase, et al., *Am J Respir Crit Care Med* 154:504-510, 1996.

The ability of compounds of this invention to treat inflammatory bowel disease can be demonstrated in a rabbit chemical-induced colitis model according to the method of Bennet et al., *J Pharmacol Exp Ther* 280:988-1000, 1997.

The ability of compounds of this invention to treat autoimmune diabetes can be demonstrated in an NOD mouse model according to the method of Hasagawa et al., *Int Immunol* 6:831-838, 1994, or in a murine streptozotocin-induced diabetes model according to the method of Herrold et al., *Cell Immunol* 157:489-500, 1994.

The ability of compounds of this invention to treat inflammatory liver injury can be demonstrated in a murine liver injury model according to the method of Tanaka et al., *J Immunol* 151:5088-5095, 1993.

The ability of compounds of this invention to treat inflammatory glomerular injury can be demonstrated in a rat nephrotoxic serum nephritis model according to the method of Kawasaki, et al., *J Immunol* 150:1074-1083, 1993.

5

10

15

20

The ability of compounds of this invention to treat stroke can be demonstrated in a rabbit cerebral embolism stroke model according the method of Bowes et al., *Exp Neurol* 119(2): 215-219, 1993, in a rat middle cerebral artery ischemia-reperfusion

model according to the method of Chopp et al., *Stroke* 25(4): 869-875, 1994, or in a rabbit reversible spinal cord ischemia model according to the method of Clark et al., *Neurosurg* 75(4): 623-627, 1991.

5 The ability of compounds of this invention to treat peripheral artery occlusion can be demonstrated in a rat skeletal muscle ischemia / reperfusion model according to the method of Gute et al., *Mol Cell Biochem* 179: 169-187, 1998.

10 The ability of compounds of this invention to treat graft rejection can be demonstrated in a murine cardiac allograft rejection model according to the method of Isobe et al., *Science* 255: 1125-1127, 1992, in a murine thyroid gland kidney capsule model according to the method of Talento et al., *Transplantation* 55: 418-422, 1993, in a cynomolgus monkey renal allograft model according to the method of Cosimi et al., *J Immunol* 144: 4604-4612, 1990, in a rat nerve allograft model according to the
15 method of Nakao et al., *Muscle Nerve* 18: 93-102, 1995, in a murine skin allograft model according to the method of Gorczynski and Wojcik, *J Immunol* 152: 2011-2019, 1994, in a murine corneal allograft model according to the method of He et al., *Ophthalmol Vis Sci* 35: 3218-3225, 1994, or in a xenogeneic pancreatic islet cell transplantation model according to the method of Zeng et al., *Transplantation* 58: 681-
20 689, 1994.

The ability of compounds of this invention to treat graft-vs.-host disease (GVHD) can be demonstrated in a murine lethal GVHD model according to the method of Harning et al., *Transplantation* 52:842-845, 1991.


- The ability of compounds of this invention to treat cancers can be
- 5 demonstrated in a human lymphoma metastasis model (in mice) according to the method of Aoudjit et al., *J Immunol* 161:2333-2338, 1998.

0054175, 033100

CLAIMS

What is claimed is:

- 5 1. A compound of formula I



I


or a pharmaceutically-acceptable salt or prodrug thereof,

10


wherein R₁, R₂, R₃, R₄, and R₅ are independently selected from

- a. hydrogen,
- b. halogen,
- 15 c. alkyl,
- d. haloalkyl,
- e. alkoxy,
- f. cyano,
- g. nitro,
- 20 h. carboxaldehyde, and

with the proviso that at least one of R_1 or R_3 is a "*cis*-cinnamide" or a "*trans*-cinnamide", defined as



"*cis*-cinnamide"



"*trans*-cinnamide",

wherein R_8 and R_9 are independently selected from

- a. hydrogen, and
- b. alkyl,
- c. carboxy alkyl,
- d. alkylaminocarbonyl alkyl, and
- e. dialkylaminocarbonyl alkyl,

and R_{10} and R_{11} are independently selected from

- a. hydrogen,
- b. alkyl,
- c. cycloalkyl,
- d. alkoxy carbonyl alkyl,
- e. hydroxy alkyl,

- f. substituted aryl,
- g. heterocyclyl,
- h. heterocyclylalkyl,
- i. heterocyclylamino,
- 5 j. substituted heterocyclyl, and
- k. substituted heterocyclylalkyl,

or where $\text{NR}_{10}\text{R}_{11}$ is heterocyclyl or substituted heterocyclyl, where
substituents are independently selected from

- 1) alkyl,
- 2) alkoxy,
- 3) alkoxyalkyl,
- 4) cycloalkyl,
- 15 5) aryl,
- 6) heterocyclyl,
- 7) heterocyclylcarbonyl,
- 8) heterocyclylalkylaminocarbonyl,
- 9) hydroxy,
- 20 10) hydroxyalkyl,
- 11) hydroxyalkoxyalkyl,
- 12) carboxy,

- 13) carboxyalkyl,
14) carboxycarbonyl,
15) carboxaldehyde,
16) alkoxycarbonyl,
5 17) arylalkoxycarbonyl,
18) aminoalkyl,
19) aminoalkanoyl,
20) carboxamido,
21) alkoxycarbonylalkyl,
10 22) carboxamidoalkyl,
23) cyano,
24) tetrazolyl,
25) substituted tetrazolyl,
26) alkanoyl,
15 27) hydroxyalkanoyl,
28) alkanoyloxy,
29) alkanoylamino,
30) alkanoyloxyalkyl,
31) alkanoylaminoalkyl,
20 32) sulfonate,
33) alkylsulfonyl,
34) alkylsulfonylaminocarbonyl,

D01E00-2621-1560

35) arylsulfonylaminocarbonyl, and

36) heterocyclylsulfonylaminocarbonyl,

and wherein **Ar** is a substituted aryl or substituted heteroaryl group, where

5 substitutions are independently selected from

a. hydrogen,

b. halogen,

c. alkyl,

10 d. aryl,

e. haloalkyl,

f. hydroxy,

g. alkoxy,

h. alkoxyalkyl,

15 i. alkoxycarbonyl,

j. alkoxyalkoxy,

k. hydroxyalkyl,

l. aminoalkyl,

m. aminocarbonyl,

20 n. alkyl(alkoxycarbonylalkyl)aminoalkyl,

o. heterocyclyl,

p. substituted heterocyclyl,

- q. heterocyclalkyl,
- r. substituted heterocyclalkyl,
- s. carboxaldehyde,
- t. carboxaldehyde hydrazone,
- 5 u. carboxamide,
- v. alkoxycarbonylalkyl,
- w. carboxy,
- x. carboxyalkyl,
- y. carboxyalkoxy,
- 10 z. carboxythioalkoxy,
- aa. carboxycycloalkoxy,
- bb. thioalkyl,
- cc. hydroxycarbonylalkyl (carboxyalkyl),
- dd. hydroxyalkylaminocarbonyl,
- 15 ee. cyano,
- ff. amino,
- gg. heterocyclalkylamino,
- hh. carboxyalkylamino,
- ii. heterocyclalkylaminocarbonyl, and
- 20 jj. "*trans*-cinnamide",

or a pharmaceutically-acceptable salt or prodrug thereof.

2. A compound according to Claim 1 wherein R₁ is a "*cis*-cinnamide" or a "*trans*-cinnamide", and R₃ is hydrogen.
- 5 3. A compound according to Claim 1 wherein R₃ is a "*cis*-cinnamide" or a "*trans*-cinnamide", and R₁ is hydrogen.
4. A compound according to Claim 1 wherein R₃ is a "*cis*-cinnamide" or a "*trans*-cinnamide", and R₁, R₈, and R₉ are hydrogen.
- 10 5. A compound according to Claim 4 wherein R₃ is a "*cis*-cinnamide".
6. A compound according to Claim 4 wherein R₃ is a "*trans*-cinnamide".
- 15 7. A compound according to Claim 1 wherein R₃ is a "*cis*-cinnamide" or a "*trans*-cinnamide", R₁, R₂, and R₄ are each independently hydrogen or alkyl; and R₅ is selected from halogen, haloalkyl, and nitro.
8. A compound according to Claim 4 wherein **Ar** is substituted aryl or substituted
- 20 heteroaryl.

9. A compound according to Claim 4 wherein R_{10} and R_{11} are each independently selected from hydrogen, alkyl, cycloalkyl, alkoxy carbonyl alkyl, hydroxy alkyl, and heterocyclalkyl.
- 5 10. A compound according to Claim 4 wherein $NR_{10}R_{11}$ is heterocyclyl or substituted heterocyclyl.
11. A compound according to Claim 8 wherein **Ar** is selected from substituted phenyl, 1,3-benzimidazol-2-one, 1,4-benzodioxane, 1,3-benzodioxole, 1-benzopyr-2-en-4-one, indole, isatin, 1,3-quinazolin-4-one, and quinoline.
- 10 12. A compound according to Claim 1 selected from the group consisting of:
 (2,4-Dichlorophenyl)[2-(*E*-((6-hydroxyhexylamino)carbonyl)ethenyl)phenyl] sulfide;
- 15 (2,4-Dichlorophenyl)[2-(*E*-((3-(1-imidazolyl)propylamino)carbonyl)ethenyl)phenyl] sulfide;
- (2,4-Dichlorophenyl)[2-chloro-4-(*E*-((2 hydroxyethylamino)carbonyl)ethenyl)phenyl] sulfide;
- (2,4-Dichlorophenyl)[2-chloro-4-(*E*-((6 hydroxyhexylamino)carbonyl)ethenyl)phenyl] sulfide;
- 20 (2,4-Dichlorophenyl)[2-chloro-4-(*E*-((bis-(2-hydroxyethyl)amino)carbonyl)ethenyl)phenyl] sulfide;

(2,4-Dichlorophenyl)[2-chloro-4-(*E*-((3-(1-pyrrolidin-2-yl)propylamino)carbonyl)ethenyl)phenyl] sulfide;

(2,4-Dichlorophenyl)[2-chloro-4-(*E*-((1-morpholinyl)carbonyl)ethenyl)phenyl] sulfide;

5 (2,4-Dichlorophenyl)[2-chloro-4-(*E*-((4-methylpiperazin-1-yl)carbonyl)ethenyl)phenyl] sulfide;

(2,4-Dichlorophenyl)[2-chloro-4-(*E*-((4-acetyl) piperazin-1-yl)carbonyl)ethenyl)phenyl] sulfide;

(2,4-Dichlorophenyl)[2-chloro-4-(*E*-((4-(2-pyridyl)piperazin-1-yl)carbonyl)

10 ethenyl)phenyl] sulfide;

(2-(Hydroxymethyl)phenyl)[2-chloro-4-(*E*-((1-morpholinyl)carbonyl)ethenyl)phenyl] sulfide;

(2-Bromophenyl)[2-chloro-4-(*E*-((1-morpholinyl)carbonyl) ethenyl)phenyl] sulfide;

(2,4-Dichlorophenyl)[2-chloro-4-(*E*-((4-(2-hydroxyethyl)piperazin-1-yl)carbonyl)

15 ethenyl)phenyl] sulfide;

(2,4-Dichlorophenyl)[2-chloro-4-(*E*-((4-(2-hydroxyethoxyethyl)piperazin-1-yl)carbonyl) ethenyl)phenyl] sulfide;

(2-Bromophenyl)[2-chloro-4-(*E*-((3-(hydroxymethyl)piperidin-1-yl)carbonyl)ethenyl)phenyl] sulfide;

20 (2-Bromophenyl)[2-chloro-4-(*E*-((2-(hydroxymethyl)piperidin-1-yl)carbonyl)ethenyl)phenyl] sulfide;

0054795.033400
00FEE0.5674560

(2-Bromophenyl)[2-chloro-4-(*E*-((3-acetamidopyrrolidin-1-yl)carbonyl)ethenyl)phenyl] sulfide;

(2-Bromophenyl)[2-chloro-4-(*E*-((4-hydroxypiperidin-1-yl)carbonyl)ethenyl)phenyl] sulfide;

- 5 (2-Bromophenyl)[2-chloro-4-(*E*-((piperidin-1-yl)carbonyl) ethenyl)phenyl] sulfide;
(2,4-Dichlorophenyl)[2-chloro-4-(*E*-((3-carboxypiperidin-1-yl)carbonyl)ethenyl)phenyl] sulfide;

(2,4-Dichlorophenyl)[2-chloro-4-(*E*-((4-carboxypiperidin-1-yl)carbonyl)ethenyl)phenyl] sulfide;

- 10 (2-Bromophenyl)[2-chloro-4-(*E*-((4-acetylhomopiperazin-1-yl)carbonyl)ethenyl)phenyl] sulfide;

(2-Bromophenyl)[2-chloro-4-(*E*-((thiomorpholin-1-yl)carbonyl)ethenyl)phenyl] sulfide;

(2-Bromophenyl)[2-chloro-4-(*E*-((4-(1-benzimidazol-2-onyl)piperidin-1-yl)carbonyl)ethenyl)phenyl] sulfide;

- 15 (2-Bromophenyl)[2-chloro-4-(*E*-((2-etrahydroisoquinoliny)carbonyl)ethenyl)phenyl] sulfide;

(2-Methylphenyl)[2-trifluoromethyl-4-(*E*-((4-acetyl)piperazin-1-yl)carbonyl)ethenyl)phenyl] sulfide;

- 20 (2-Methylphenyl)[2-trifluoromethyl-4-(*E*-((1-morpholinyl)carbonyl)ethenyl)phenyl] sulfide;

(2-Methylphenyl)[2-trifluoromethyl-4-(*E*-((2-(1-morpholinyl)ethylamino)carbonyl)

ethenyl)phenyl] sulfide;

(2-Methylphenyl)[2-trifluoromethyl-4-(*E*-((4-phenylpiperazin-1-yl)carbonyl)

ethenyl)phenyl] sulfide;

(2-Methylphenyl)[2-trifluoromethyl-4-(*E*-((3-(1-pyrrolidin-2-onyl)propylamino)

5 carbonyl) ethenyl)phenyl] sulfide;

(2-Methylphenyl)[2-trifluoromethyl-4-(*E*-

((cyclopropylamino)carbonyl)ethenyl)phenyl]

sulfide;

(2,4-Dichlorophenyl)[2-nitro-4-(*E*-((4-acetyl) piperazin-1-yl)carbonyl)ethenyl)phenyl]

10 sulfide;

(2,4-Dichlorophenyl)[2-nitro-4-(*E*-((3-(1-pyrrolidin-2-onyl)propylamino)carbonyl)

ethenyl)phenyl] sulfide;

(2,3-Dichlorophenyl)[2-nitro-4-(*E*-((4-acetyl) piperazin-1-yl)carbonyl)ethenyl)phenyl]

sulfide;

15 (4-Bromophenyl)[2-nitro-4-(*E*-((4-acetyl) piperazin-1-yl)carbonyl)ethenyl)phenyl]

sulfide;

(4-Methylphenyl)[2-nitro-4-(*E*-((4-acetyl) piperazin-1-yl)carbonyl)ethenyl)phenyl]

sulfide;

(2,4-Dichlorophenyl)[2-nitro-4-(*E*-((4-(*tert*-butoxycarbonyl) piperazin-1-yl)carbonyl)

20 ethenyl)phenyl] sulfide;

(2,4-Dichlorophenyl)[2-nitro-4-(*E*-((4-(2-furoyl)carbonyl) piperazin-1-yl)carbonyl)

ethenyl)phenyl] sulfide;

00544795.033100
00544795.033100

- (2,4-Dichlorophenyl)[2-nitro-4-(*E*-((4-(methanesulfonyl)piperazin-1-yl)carbonyl)ethenyl)phenyl] sulfide;
- (2,4-Dichlorophenyl)[2-nitro-4-(*E*-((4-(diethylaminocarbonylmethyl)piperazin-1-yl)carbonyl)ethenyl)phenyl] sulfide;
- 5 (2,4-Dichlorophenyl)[2-nitro-4-(*E*-((4-(diethylaminocarbonyl)piperazin-1-yl)carbonyl)ethenyl)phenyl] sulfide;
- (2,4-Dichlorophenyl)[2-nitro-4-(*E*-((4-(*tert*-butoxycarbonylmethyl)piperazin-1-yl)carbonyl)ethenyl)phenyl] sulfide;
- (2,4-Dichlorophenyl)[2-nitro-4-(*E*-((4-(carboxycarbonyl)piperazin-1-yl)carbonyl)ethenyl)phenyl] sulfide;
- 10 (2,4-Dichlorophenyl)[2-nitro-4-(*E*-((4-(carboxymethyl)piperazin-1-yl)carbonyl)ethenyl)phenyl] sulfide;
- (2-Methylphenyl)[2-nitro-4-(*E*-((4-(acetyl)piperazin-1-yl)carbonyl)ethenyl)phenyl] sulfide;
- 15 (2-Chlorophenyl)[2-nitro-4-(*E*-((4-(acetyl)piperazin-1-yl)carbonyl)ethenyl)phenyl] sulfide;
- (2-Aminophenyl)[2-nitro-4-(*E*-((4-(acetyl)piperazin-1-yl)carbonyl)ethenyl)phenyl] sulfide;
- (2-Hydroxymethylphenyl)[2-nitro-4-(*E*-((4-(acetyl)piperazin-1-yl)carbonyl)ethenyl)phenyl] sulfide;
- 20 (2-Ethylphenyl)[2-nitro-4-(*E*-((4-(acetyl)piperazin-1-yl)carbonyl)ethenyl)phenyl] sulfide;

(2-*iso*-Propylphenyl)[2-nitro-4-(*E*-((4-acetyl)piperazin-1-yl)carbonyl)ethenyl]phenyl sulfide;

(2-*tert*-Butylphenyl)[2-nitro-4-(*E*-((4-acetyl)piperazin-1-yl)carbonyl)ethenyl]phenyl sulfide;

- 5 (2-Chlorophenyl)[2-chloro-4-(*E*-((4-acetyl)piperazin-1-yl)carbonyl))2-propenyl]phenyl sulfide;

(2-(1-Morpholinylmethyl)phenyl)[2-chloro-4-(*E*-((1-morpholinyl)carbonyl) ethenyl] phenyl sulfide;

- 10 (2-(4-(1,3-Benzodioxolyl-5-methyl)piperazin-1-ylmethyl)phenyl)[2-chloro-4-(*E*-((1-morpholinyl)carbonyl) ethenyl)phenyl] sulfide;
- (2-(4-(*iso*-Propylaminocarbonylmethyl)piperazin-1-ylmethyl)phenyl)[2-chloro-4-(*E*-((1-morpholinyl)carbonyl) ethenyl)phenyl] sulfide;
- (2-((*N*-Ethoxycarbonylmethyl-*N*-methyl)aminomethyl)phenyl)[2-chloro-4-(*E*-((1-

- 15 morpholinyl)carbonyl) ethenyl)phenyl] sulfide;

(2-Formylphenyl)[2-chloro-4-(*E*-((1-morpholinyl)carbonyl)ethenyl)phenyl] sulfide;

(2-(4-Formylpiperazin-1-ylmethyl)phenyl)[2-chloro-4-(*E*-((1-morpholinyl)carbonyl) ethenyl)phenyl] sulfide;

- (2-(*E*-((1-Morpholinyl)carbonyl)ethenyl)phenyl)[2-chloro-4-(*E*-((1-morpholinyl) carbonyl)ethenyl)phenyl] sulfide;
- 20 (2-Formylphenyl)[2-nitro-4-(*E*-((4-acetyl)piperazin-1-yl)carbonyl)ethenyl]phenyl sulfide;

- (2-Formylphenyl)[2-chloro-4-(*E*-((1-morpholinyl)carbonyl)ethenyl)phenyl] sulfide,
N,N-dimethyl hydrazone;
- (2-((3-(1-Morpholinyl)propyl)-1-amino)phenyl)[2-chloro-4-(*E*-((1-morpholinyl)carbonyl) ethenyl)phenyl] sulfide;
- 5 (2,4-Dichlorophenyl)[2-bromo-4-(*E*-((3-(1-pyrrolidin-2-onyl)propylamino)carbonyl)ethenyl)phenyl] sulfide;
- (2,4-Dichlorophenyl)[2-formyl-4-(*E*-((1-morpholinyl)carbonyl)ethenyl)phenyl] sulfide;
- (2-Chloro-6-formylphenyl)[2-chloro-4-(*E*-((4-acetylpiperazin-1-yl)carbonyl)ethenyl)phenyl] sulfide;
- 10 (2-Cyanophenyl)[2-chloro-4-(*E*-((4-acetylpiperazin-1-yl)carbonyl)ethenyl) phenyl] sulfide;
- (2-Isopropylphenyl)[2-cyano-4-(*E*-((morpholin-1-yl)carbonyl)ethenyl) phenyl] sulfide;
- 15 (2-Bromophenyl)[2-nitro-4-(*E*-((4-acetylpiperazin-1-yl)carbonyl)ethenyl) phenyl] sulfide;
- (2-(Pyrrolidin-1-yl)phenyl)[2-chloro-4-(*E*-((morpholin-1-yl)carbonyl)ethenyl) phenyl] sulfide;
- (2-Methoxyphenyl)-[2-chloro-4(*E*-[(morpholin-1-yl)carbonyl]ethenyl)phenyl]sulfide;
- 20 (2-Isopropylphenyl)[2-nitro-4-(*E*-((3-carbomethoxypiperazin-1-yl)carbonyl)ethenyl) phenyl] sulfide;

0054795 03100

- (2-Methylphenyl)[2-nitro-4-(*E*-((3-carboxamido-4-carbobenzoxypiperazin-1-yl)carbonyl)ethenyl) phenyl] sulfide;
- (2-Isopropylphenyl)[2-nitro-4-(*E*-((2-carbomethoxy-4-*tert*-butoxycarbonylpiperazin-1-yl)carbonyl)ethenyl) phenyl] sulfide;
- 5 (2-Isopropylphenyl)[2-nitro-4-(*E*-((2-carboxy-4-*tert*-butoxycarbonylpiperazin-1-yl)carbonyl)ethenyl) phenyl] sulfide;
- (2-Isopropylphenyl)[2-trifluoromethyl-4-(*E*-((4-acetylpiperazin-1-yl)carbonyl)ethenyl) phenyl] sulfide;
- (2-Isopropylphenyl)[2-trifluoromethyl-4-(*E*-((morpholin-1-yl)carbonyl)ethenyl) phenyl] sulfide;
- 10 (2-Isopropylphenyl)[2-trifluoromethyl-4-(*E*-((3-(pyrrolidin-2-on-1-yl)prop-1-ylamino)carbonyl) ethenyl)phenyl]sulfide;
- (2-Isopropylphenyl)[2-trifluoromethyl-4-(*E*-((cyclobutylamino)carbonyl)ethenyl) phenyl] sulfide;
- 15 (2-Isopropylphenyl)[2-trifluoromethyl-4-(*E*-((cyclopentylamino)carbonyl)ethenyl) phenyl] sulfide;
- (2-Isopropylphenyl)[2-trifluoromethyl-4-(*E*-((5-hydroxypent-1-ylamino)carbonyl)ethenyl) phenyl] sulfide;
- (2-Isopropylphenyl)[2-nitro-4-(*E*-((3-carbomethoxy-4-acetylpiperazin-1-yl)carbonyl)ethenyl) phenyl] sulfide;
- 20 (2-Biphenyl)[2-chloro-4-(*E*-((morpholin-1-yl)carbonyl)ethenyl) phenyl] sulfide;

- (3,4-Dimethylphenyl)[2-nitro-4-(*E*-((4-acetylpiperazin-1-yl)carbonyl)ethenyl)phenyl]sulfide;
- (2-Bromophenyl)[2-trifluoromethyl-4-(*E*-((4-acetylpiperazin-1-yl)carbonyl)ethenyl) phenyl] sulfide;
- 5 (5-Indolyl)[2-chloro-4-(*E*-((4-acetylpiperazin-1-yl)carbonyl)ethenyl) phenyl] sulfide;
- (5-Benzodioxolyl)[2-chloro-4-(*E*-((4-acetylpiperazin-1-yl)carbonyl)ethenyl) phenyl] sulfide;
- (2-Isopropylphenyl)[2-nitro-4-(*E*-((2-carbomethoxypiperazin-1-yl)carbonyl)ethenyl) phenyl] sulfide;
- 10 (2,3-Dimethoxyphenyl)-[2-chloro-4-(*E*-[(morpholin-1-yl)carbonyl]ethenyl)phenyl] sulfide;
- (2-Fluorophenyl)[2-nitro-4-(*E*-((4-acetylpiperazin-1-yl)carbonyl)ethenyl)phenyl]sulfide;
- 15 (2-Bromophenyl)[2-trifluoromethyl-4-(*E*-((4-(*tert*-butoxycarbonyl)piperazin-1-yl)carbonyl)ethenyl) phenyl] sulfide;
- (2-(Pyrrolidin-1-yl)phenyl)[2-trifluoromethyl-4-(*E*-((4-(*tert*-butoxycarbonyl)piperazin-1-yl)carbonyl)ethenyl) phenyl] sulfide;
- (3-Carboxamidophenyl)[2-nitro-4-(*E*-((4-acetylpiperazin-1-yl)carbonyl)ethenyl) phenyl] sulfide;
- 20 (3-(Hydroxymethyl)phenyl)[2-nitro-4-(*E*-((4-acetylpiperazin-1-yl)carbonyl)ethenyl) phenyl] sulfide;

00541705.03100

- Phenyl[2-trifluoromethyl-4-(*E*-((4-(*tert*-butoxycarbonyl)piperazin-1-yl)carbonyl)ethenyl) phenyl] sulfide;
- (2-Isopropylphenyl)[2-trifluoromethyl-4-(*E*-((2-carbomethoxy-4-(*tert*-butoxycarbonyl)piperazin-1-yl)carbonyl)ethenyl) phenyl] sulfide;
- 5 (2-Isopropylphenyl)[2-nitro-4-(*E*-((3-(pyridine-4-methylaminocarbonyl)-4-*tert*-butoxycarbonyl)piperazin-1-yl)carbonyl)ethenyl) phenyl] sulfide;
- (2-Ethoxyphenyl)-[2-chloro-4(*E*-[(morpholin-1-yl)carbonyl]ethenyl)phenyl]sulfide;
- (2-Methoxyphenyl)[2-nitro-4-(*E*-((4-acetyl)piperazin-1-yl)carbonyl)ethenyl]phenyl]sulfide;
- 10 (2-(Azetidin-1-yl)phenyl)[2-trifluoromethyl-4-(*E*-((4-(*tert*-butoxycarbonyl)piperazin-1-yl)carbonyl)ethenyl) phenyl] sulfide;
- (2-(Piperidin-1-yl)phenyl)[2-trifluoromethyl-4-(*E*-((4-(*tert*-butoxycarbonyl)piperazin-1-yl)carbonyl)ethenyl) phenyl] sulfide;
- (3-Chloro-2-formylphenyl)[2-chloro-4-(*E*-((4-acetyl)piperazin-1-yl)carbonyl)
- 15 ethenyl) phenyl] sulfide;
- (2-Trifluoromethylphenyl)[2-trifluoromethyl-4-(*E*-((4-acetyl)piperazin-1-yl)carbonyl)ethenyl) phenyl] sulfide;
- (3-Bromophenyl)[2-trifluoromethyl-4-(*E*-((4-acetyl)piperazin-1-yl)carbonyl)ethenyl) phenyl] sulfide;
- 20 (3,5-Dimethylphenyl)[2-trifluoromethyl-4-(*E*-((4-acetyl)piperazin-1-yl)carbonyl)ethenyl) phenyl] sulfide;

00941705, 0094100

(2-Isopropylphenyl)[2-nitro-4-(*E*-((3-dimethylaminocarbonyl-4-(pyridine-4-carbonyl)piperazin-1-yl)carbonyl)ethenyl) phenyl] sulfide;

(2-Isopropylphenyl)[2-nitro-4-(*E*-((3-dimethylaminocarbonyl-4-carbomethoxypiperazin-1-yl)carbonyl)ethenyl) phenyl] sulfide;

- 5 (2-Isopropylphenyl)[2-nitro-4-(*E*-((3-dimethylaminocarbonyl-4-acetyl)piperazin-1-yl)carbonyl)ethenyl) phenyl] sulfide;

(2-Isopropylphenyl)[2-nitro-4-(*E*-((3-(1-morpholinocarbonyl)-4-*tert*-butoxycarbonylpiperazin-1-yl)carbonyl)ethenyl) phenyl] sulfide;

- 10 (2-Isopropylphenyl)[2-nitro-4-(*E*-((3-(pyridine-4-methylaminocarbonyl)piperazin-1-yl)carbonyl)ethenyl) phenyl] sulfide;

(2-Isopropylphenyl)[2-nitro-4-(*E*-((3-dimethylaminocarbonyl)piperazin-1-yl)carbonyl)ethenyl) phenyl] sulfide;

(2-Isopropylphenyl)[2-nitro-4-(*E*-((3-(benzylaminocarbonyl)-4-*tert*-butoxycarbonylpiperazin-1-yl)carbonyl)ethenyl) phenyl] sulfide;

- 15 (2-Isopropylphenyl)[2-nitro-4-(*E*-((3-(dimethylaminocarbonyl)-4-*tert*-butoxycarbonylpiperazin-1-yl)carbonyl)ethenyl) phenyl] sulfide;

(2-Bromophenyl)[2-chloro-4-(*E*-((3-(5*S*-hydroxymethyl-pyrrolidin-2-on-1-yl)prop-1-ylamino)carbonyl) ethenyl)phenyl]sulfide;

(2-Bromophenyl)[2-chloro-4-(*E*-((3-(pyrrolidin-2-on-1-yl)prop-1-ylamino)carbonyl) ethenyl)phenyl]sulfide;

- 20 (2-Bromophenyl)[2-chloro-4-(*E*-(*N*-methyl-*N*-(3-(pyrrolidin-2-on-1-yl)prop-1-ylamino)carbonyl) ethenyl)phenyl]sulfide;

001220-26274580

- (2-[2-Methoxy]ethoxyphenyl)-[2-chloro-4-(E-[(morpholin-1-yl)carbonyl]ethenyl)phenyl] sulfide;
- (2-Isopropylphenyl)[2-nitro-4-(E-((3-(morpholinocarbonyl)piperazin-1-yl)carbonyl)ethenyl) phenyl] sulfide;
- 5 (2-Isopropylphenyl)[2-nitro-4-(E-((4-*tert*-butoxycarbonyl)piperazin-1-yl)carbonyl)ethenyl) phenyl] sulfide;
- (2-Isopropylphenyl)[2-nitro-4-(E-((4-methoxycarbonyl)piperazin-1-yl)carbonyl)ethenyl) phenyl] sulfide;
- (2-Isopropylphenyl)[2-nitro-4-(E-(4-(pyridine-4-carbonyl)piperazin-1-yl)carbonyl)ethenyl) phenyl] sulfide;
- 10 (2-Isopropylphenyl)[2-nitro-4-(E-((3-(pyridine-3-methylaminocarbonyl)-4-*tert*-butoxycarbonyl)piperazin-1-yl)carbonyl)ethenyl) phenyl] sulfide;
- (2-Isopropylphenyl)[2-nitro-4-(E-((3-(pyridine-2-methylaminocarbonyl)piperazin-1-yl)carbonyl)ethenyl) phenyl] sulfide;
- 15 (2-Isopropylphenyl)[2-nitro-4-(E-((3-(pyridine-3-methylaminocarbonyl)piperazin-1-yl)carbonyl)ethenyl) phenyl] sulfide;
- (4-Hydroxyphenyl)[2-nitro-4-(E-((4-acetyl)piperazin-1-yl)carbonyl)ethenyl]sulfide;
- (3,5-Dichlorophenyl)[2-nitro-4-(E-((4-acetyl)piperazin-1-yl)carbonyl)ethenyl]phenyl]sulfide;
- 20 (2-Bromophenyl)[2-chloro-4-(E-((3-(5*S*-acetoxymethyl-pyrrolidin-2-on-1-yl)prop-1-ylamino)carbonyl) ethenyl)phenyl]sulfide;

00120 56274560

(2-Bromophenyl)[2-chloro-4-(*E*-((3-(5*S*-methoxymethyl-pyrrolidin-2-on-1-yl)prop-1-ylamino)carbonyl)ethenyl)phenyl]sulfide;

(2-Bromophenyl)[2-chloro-4-(*E*-((3-(4*R*-hydroxymethyl-pyrrolidin-2-on-1-yl)prop-1-ylamino)carbonyl)ethenyl)phenyl]sulfide;

- 5 Phenyl[2-nitro-4-(*E*-((4-acetyl)piperazin-1-yl)carbonyl)ethenyl)phenyl]sulfide;
(2-Dimethylaminophenyl)[2-nitro-4-(*E*-((4-acetyl)piperazin-1-yl)carbonyl)ethenyl)phenyl]sulfide;

(3-((2-Hydroxyethyl)aminocarbonyl)phenyl)[2-nitro-4-(*E*-((4-acetyl)piperazin-1-yl)carbonyl)ethenyl)phenyl]sulfide;

- 10 (3-((3-(1-Imidazolyl)propyl)aminocarbonyl)phenyl)[2-nitro-4-(*E*-((4-acetyl)piperazin-1-yl)carbonyl)ethenyl)phenyl]sulfide;

(3-((2-(1-Morpholinyl)ethyl)aminocarbonyl)phenyl)[2-nitro-4-(*E*-((4-acetyl)piperazin-1-yl)carbonyl)ethenyl)phenyl]sulfide;

(2-Isopropylphenyl)[2-nitro-4-(*E*-((3-hydroxymethyl-4-*tert*-

- 15 butoxycarbonyl)piperazin-1-yl)carbonyl)ethenyl)phenyl]sulfide;

(2-Isopropylphenyl)[2-nitro-4-(*E*-((4-formyl)piperazin-1-yl)carbonyl)ethenyl)phenyl]sulfide;

(2-Isopropylphenyl)[2-nitro-4-(*E*-((2-hydroxymethyl-4-*tert*-butoxycarbonyl)piperazin-1-yl)carbonyl)ethenyl)phenyl]sulfide;

- 20 (2-Ethoxyphenyl)-[2-chloro-4(*E*-[(3-ethoxycarbonyl)piperidin-1-yl)carbonyl]ethenyl)phenyl]sulfide;

00541705.033100

- (3- Aminophenyl)[2-nitro-4-(*E*-((4-acetylpiperazin-1-yl)carbonyl)ethenyl)phenyl]sulfide;
- (4-Aminophenyl)[2-nitro-4-(*E*-((4-acetylpiperazin-1-yl)carbonyl)ethenyl)phenyl]sulfide;
- 5 (2,4-Dimethylphenyl)[2- nitro-4-(*E*-((4-acetylpiperazin-1-yl)carbonyl)ethenyl)phenyl]sulfide;
- (2,5-Dimethylphenyl)[2- nitro-4-(*E*-((4-acetylpiperazin-1-yl)carbonyl)ethenyl)phenyl]sulfide;
- (4-Methoxyphenyl)[2-nitro-4-(*E*-((4-acetylpiperazin-1-yl)carbonyl)ethenyl)phenyl]sulfide;
- 10 (3-Chlorophenyl)[2-nitro-4-(*E*-((4-acetylpiperazin-1-yl)carbonyl)ethenyl)phenyl]sulfide;
- (2-Chloro, 4,5-diaminophenyl)[2-chloro-4-(*E*-((4-acetylpiperazin-1-yl)carbonyl)ethenyl) phenyl] sulfide;
- 15 (3,4-Diaminophenyl)[2-chloro-4-(*E*-((4-acetylpiperazin-1-yl)carbonyl)ethenyl)phenyl]
- (6-Chlorobenzimidazol-2-on-5-yl)[2-chloro-4-(*E*-((4-acetylpiperazin-1-yl)carbonyl)ethenyl) phenyl] sulfide;
- (1-Methylindol-7-yl)[2-chloro-4-(*E*-((4-acetylpiperazin-1-yl)carbonyl)ethenyl)phenyl] sulfide;
- 20 phenyl] sulfide;
- (2-Hydroxy, 4-aminophenyl)[2-chloro-4-(*E*-((4-acetylpiperazin-1-yl)carbonyl)ethenyl) phenyl] sulfide;

00544705.033100

- (2-Isopropylphenyl)[2-nitro-4-(*E*-((4-methylpiperazin-1-yl)carbonyl)ethenyl)phenyl] sulfide;
- (2-Isopropylphenyl)[2-nitro-4-(*E*-((4-(pyridine-2-carbonyl)piperazin-1-yl)carbonyl)ethenyl) phenyl] sulfide;
- 5 (2-Isopropylphenyl)[2-nitro-4-(*E*-((4-(pyridine-3-carbonyl)piperazin-1-yl)carbonyl)ethenyl) phenyl] sulfide;
- (2-Isopropylphenyl)[2-nitro-4-(*E*-((2-carbomethoxy-4-methoxycarbonyl)piperazin-1-yl)carbonyl)ethenyl) phenyl] sulfide;
- (2-Isopropylphenyl)[2-nitro-4-(*E*-((2-carboxy-4-methoxycarbonyl)piperazin-1-yl)carbonyl)ethenyl) phenyl] sulfide;
- 10 (2-Isopropylphenyl)[2-nitro-4-(*E*-((3-carbomethoxy-4-methylpiperazin-1-yl)carbonyl)ethenyl) phenyl] sulfide;
- (2-Ethoxyphenyl)-[2-chloro-4(*E*-[(3-carboxypiperidin-1-yl)carbonyl]ethenyl)phenyl] sulfide;
- 15 (2-Ethoxyphenyl)-[2-chloro-4(*E*-[(3-carboxypiperidin-1-yl)carbonyl]ethenyl)phenyl] sulfide;
- (2-Ethoxyphenyl)-[2-chloro-4(*E*-[(2-ethoxycarbonyl)piperidin-1-yl)carbonyl]ethenyl)phenyl] sulfide;
- (2-Ethoxyphenyl)[2-trifluoromethyl-4-(*E*-((1-(*tert*-butoxycarbonyl)-4-hydroxypyrrolidin-3-ylamino)carbonyl)ethenyl) phenyl] sulfide;
- 20 (2-Ethoxyphenyl)-[2-chloro-4(*E*-[(2-carboxypiperidin-1-yl)carbonyl]ethenyl)phenyl] sulfide;

(2-Ethoxyphenyl)[2-trifluoromethyl-4-(*E*-(((pyrrol-3-yl)carbonyl)ethenyl)phenyl] sulfide;

(2-Ethoxyphenyl)[2-trifluoromethyl-4-(*E*-((3-(pyrrolidin-2-on-1-yl)prop-1-ylamino)carbonyl)ethenyl)phenyl]sulfide;

- 5 (2-Ethoxyphenyl)[2-trifluoromethyl-4-(*E*-((4-acetylpiperazin-1-yl)carbonyl)ethenyl)phenyl] sulfide;

(2-Ethoxyphenyl)[2-trifluoromethyl-4-(*E*-((4-(ethoxycarbonyl)piperazin-1-yl)carbonyl)ethenyl)phenyl] sulfide;

- 10 (2-Ethoxyphenyl)[2-trifluoromethyl-4-(*E*-((4-(2-furylcarbonyl)piperazin-1-yl)carbonyl)ethenyl)phenyl] sulfide;

(2-Ethoxyphenyl)-[2-chloro-4-(*E*-[(3-ethoxycarbonylpiperidin-1-yl)carbonyl]ethenyl)phenyl]sulfide;

(2-Ethoxyphenyl)-[2-chloro-4-(*E*-[(4-carboxypiperidin-1-yl)carbonyl]ethenyl)phenyl] sulfide;

- 15 (Benzodioxan-6-yl)[2-chloro-4-(*E*-((4-acetylpiperazin-1-yl)carbonyl)ethenyl)phenyl] sulfide;

(2-Isopropylphenyl)[2-nitro-4-(*E*-((4-ethoxycarbonylpiperazin-1-yl)carbonyl)ethenyl)phenyl] sulfide;

- 20 (2-Isopropylphenyl)[2-nitro-4-(*E*-((4-isopropoxycarbonylpiperazin-1-yl)carbonyl)ethenyl)phenyl] sulfide;

(2-Isopropylphenyl)[2-nitro-4-(*E*-((4-isobutoxycarbonylpiperazin-1-yl)carbonyl)ethenyl)phenyl] sulfide;

(2-Isopropylphenyl)[2-nitro-4-(*E*-((4-((1-propen-2-oxycarbonyl)piperazin-1-yl)carbonyl)ethenyl) phenyl] sulfide;

(2-Isopropylphenyl)[2-nitro-4-(*E*-((4-propionylpiperazin-1-yl)carbonyl)ethenyl) phenyl] sulfide;

5 (2-Isopropylphenyl)[2-nitro-4-(*E*-((4-carboxamidopiperazin-1-yl)carbonyl)ethenyl) phenyl] sulfide;

(2-Isopropylphenyl)[2-nitro-4-(*E*-((4-methylaminocarbonylpiperazin-1-yl)carbonyl)ethenyl) phenyl] sulfide;

(2-Isopropylphenyl)[2-nitro-4-(*E*-((4-(pyrimidin-2-yl)piperazin-1-yl)carbonyl)ethenyl) phenyl] sulfide;

10 (2-Isopropylphenyl)[2-nitro-4-(*E*-((4-hydroxyacetyl)piperazin-1-yl)carbonyl)ethenyl) phenyl] sulfide;

(2-Isopropylphenyl)[2-nitro-4-(*E*-((4-(pyrazine-2-carbonyl)piperazin-1-yl)carbonyl)ethenyl) phenyl] sulfide;

15 (2-Isopropylphenyl)[2-trifluoromethyl-4-(*E*-((2-carboxypyrrol-3-in-1-yl)carbonyl)ethenyl) phenyl] sulfide;

(2-Isopropylphenyl)[2-nitro-4-(*E*-((3-hydroxymethyl-4-methylpiperazin-1-yl)carbonyl)ethenyl) phenyl] sulfide;

(2-Isopropylphenyl)[2-trifluoromethyl-4-(*E*-((2-carboxypyrrol-3-in-1-yl)carbonyl)ethenyl) phenyl] sulfide;

20 (2-Isopropylphenyl)[2-trifluoromethyl-4-(*E*-((2-hydroxymethylpyrrolidin-1-yl)carbonyl)ethenyl) phenyl] sulfide;

(2-Isopropylphenyl)[2-nitro-4-(*E*-((3-methylaminocarbonyl)piperazin-1-yl)carbonyl)ethenyl] phenyl] sulfide;

(2-Isopropylphenyl)[2-nitro-4-(*E*-(((3-cyclopropylaminocarbonyl)piperazin-1-yl)carbonyl)ethenyl] phenyl] sulfide;

5 (2-Isopropylphenyl)[2-nitro-4-(*E*-((3-carboxamidopiperazin-1-yl)carbonyl)ethenyl) phenyl] sulfide;

(2-Isopropylphenyl)[2-nitro-4-(*E*-((3-carbomethoxy-4-oxopiperidin-1-yl)carbonyl)ethenyl] phenyl] sulfide;

(2-Isopropylphenyl)[2-nitro-4-(*E*-((3,5-dimethylpiperazin-1-yl)carbonyl)ethenyl) phenyl] sulfide;

10 (1-Ethylindol-7-yl)[2-chloro-4-(*E*-((4-acetyl)piperazin-1-yl)carbonyl)ethenyl] phenyl] sulfide;

(3-[2-Methoxy]ethoxyphenyl)-[2-chloro-4(*E*-[(morpholin-1-yl)carbonyl]ethenyl)phenyl] sulfide;

15 (2-Bromophenyl)[2-chloro-4-(*E*-((4,4'-*S*-dioxythiomorpholin-1-yl)carbonyl) ethenyl)phenyl]sulfide;

(2-Bromophenyl)[2-chloro-4-(*E*-(*N*-carbomethoxymethyl-*N*-(3-(pyrrolidin-2-on-1-yl)prop-1-yl)amino)carbonyl) ethenyl)phenyl]sulfide;

(2-Bromophenyl)[2-chloro-4-(*E*-((4-*S*-oxythiomorpholin-1-yl)-2-

pyrrolidinone)carbonyl) ethenyl)phenyl]sulfide;

(2-Methoxy-5-chlorophenyl)[2-nitro-4-(*E*-((4-acetyl)piperazin-1-yl)carbonyl)ethenyl) phenyl] sulfide;

00541795-033100

- (2-Isopropylphenyl)[2-nitro-4-(*E*-((3-acetoxymethyl)piperazin-1-yl)carbonyl)ethenyl] phenyl] sulfide;
- (2-Isopropylphenyl)[2-nitro-4-(*E*-((3,5-dimethyl-4-acetyl)piperazin-1-yl)carbonyl)ethenyl] phenyl] sulfide;
- 5 (1-Methylindol-5-yl)[2-chloro-4-(*E*-((4-acetyl)piperazin-1-yl)carbonyl)ethenyl] phenyl] sulfide;
- (Benzodioxan-6-yl)[2-nitro-4-(*E*-((4-acetyl)piperazin-1-yl)carbonyl)ethenyl] phenyl] sulfide;
- (Benzodioxan-6-yl)[2-nitro-4-(*E*-((3-(pyrrolidin-2-on-1-yl)prop-1-ylamino)carbonyl)ethenyl)phenyl]sulfide;
- 10 (Benzodioxan-6-yl)[2-nitro-4-(*E*-((3-carboethoxypiperidin-1-yl) carbonyl)ethenyl] phenyl] sulfide;
- (Benzodioxan-6-yl)[2-nitro-4-(*E*-((4-carboethoxypiperidin-1-yl) carbonyl)ethenyl] phenyl] sulfide;
- 15 (2-Ethoxyphenyl)[2-trifluoromethyl-4-(*Z*-((4-acetyl)piperazin-1-yl)carbonyl)ethenyl] phenyl] sulfide;
- (2-Ethoxyphenyl)[2-trifluoromethyl-4-(*E*-((6-methylpyrid-2-ylamino)carbonyl)ethenyl) phenyl] sulfide;
- (2-Methyl-3-chlorophenyl)[2-nitro-4-(*E*-((4-acetyl)piperazin-1-yl)carbonyl)ethenyl] phenyl] sulfide;
- 20 (Benzodioxan-6-yl)[2-nitro-4-(*E*-((3-carboxamidopiperidin-1-yl) carbonyl)ethenyl] phenyl] sulfide;

0541395-053100

(Benzodioxan-6-yl)[2-nitro-4-(*E*-((2-carboethoxypiperidin-1-yl) carbonyl)ethenyl) phenyl] sulfide;

(Benzodioxan-6-yl)[2-nitro-4-(*E*-((4-carboxamidopiperidin-1-yl) carbonyl)ethenyl) phenyl] sulfide;

5 (Benzodioxan-6-yl)[2-nitro-4-(*E*-((4-*tert*-butoxycarbonylpiperazin-1-yl) carbonyl)ethenyl) phenyl] sulfide;

(2-Isopropylphenyl)[2-nitro-4-(*E*-((*syn*-3,5-dimethylmorpholin-1-yl) carbonyl)ethenyl) phenyl] sulfide;

(2-Isopropylphenyl)[2-nitro-4-(*E*-((*anti*-3,5-dimethylmorpholin-1-yl) carbonyl)ethenyl) phenyl] sulfide;

10 (2-Isopropylphenyl)[2-nitro-4-(*E*-((3-carboethoxypiperazin-1-yl) carbonyl)ethenyl) phenyl] sulfide;

(2-Isopropylphenyl)[2-nitro-4-(*E*-((3-isopropoxycarbonylpiperazin-1-yl) carbonyl)ethenyl) phenyl] sulfide;

15 (2-Isopropylphenyl)[2-nitro-4-(*E*-((3-(dimethylaminocarbonyl)-4-methylpiperazin-1-yl) carbonyl)ethenyl) phenyl] sulfide;

(2-Isopropylphenyl)[2-nitro-4-(*E*-((3-carbomethoxy-4-hydroxypiperidin-1-yl) carbonyl)ethenyl) phenyl] sulfide;

(2-Isopropylphenyl)[2-nitro-4-(*E*-((3-hydroxymethyl-4-hydroxypiperidin-1-yl) carbonyl)ethenyl) phenyl] sulfide;

20 (2-Ethoxyphenyl)[2-trifluoromethyl-4-(*E*-((2-carbomethoxy-4-(methoxycarbonyl)piperazin-1-yl) carbonyl)ethenyl) phenyl] sulfide;

00541735:03300

- (2-Ethoxyphenyl)[2-trifluoromethyl-4-(*E*-((2-carbomethoxy-4-methyl piperazin-1-yl)carbonyl)ethenyl) phenyl] sulfide;
- (2-Ethoxyphenyl)[2-trifluoromethyl-4-(*E*-((2-carboxy-4-(methoxycarbonyl)piperazin-1-yl)carbonyl)ethenyl) phenyl] sulfide;
- 5 (Indol-6-yl)[2-chloro-4-(*E*-((4-acetylpiperazin-1-yl)carbonyl)ethenyl) phenyl] sulfide;
- (1-Ethyl,3-(dimethylaminomethyl)indol-7-yl)[2-chloro-4-(*E*-((4-acetylpiperazin-1-yl)carbonyl)ethenyl) phenyl] sulfide;
- (5-Ethoxybenzodioxan-6-yl)[2-chloro-4-(*E*-((4-acetylpiperazin-1-yl)carbonyl)ethenyl) phenyl] sulfide;
- 10 (2-Ethyl-4-bromophenyl)[2-nitro-4-(*E*-((4-acetylpiperazin-1-yl)carbonyl)ethenyl) phenyl] sulfide;
- (Benzodioxan-6-yl)[2-nitro-4-(*E*-((2-carboxypiperidin-1-yl) carbonyl)ethenyl) phenyl] sulfide;
- (Benzodioxan-6-yl)[2-nitro-4-(*E*-((4-carboxymethylpiperazin-1-yl) carbonyl)ethenyl) phenyl] sulfide;
- 15 (3-Morpholinophenyl)[2-nitro-4-(*E*-((4-acetylpiperazin-1-yl)carbonyl)ethenyl) phenyl] sulfide;
- (5-Ethoxybenzodioxan-8-yl)[2-chloro-4-(*E*-((4-acetylpiperazin-1-yl)carbonyl)ethenyl) phenyl] sulfide;
- 20 (5-Chloro-8-ethoxyquinolin-7-yl)[2-chloro-4-(*E*-((4-acetylpiperazin-1-yl)carbonyl)ethenyl) phenyl] sulfide;

00541795, 033100

(2-Isopropylphenyl)[2-nitro-4-(*E*-((3-carboethoxypiperidin-1-yl)carbonyl)ethenyl)phenyl] sulfide;

(2-Isopropylphenyl)[2-nitro-4-(*E*-((3-carboxypiperidin-1-yl)carbonyl)ethenyl)phenyl] sulfide;

- 5 (2-Isopropylphenyl)[2-nitro-4-(*E*-(((3-ethanesulfonylamino)carbonyl)piperidin-1-yl)carbonyl)ethenyl)phenyl] sulfide;

(2-Isopropylphenyl)[2-nitro-4-(*E*-((3-(4-methylpiperazine)sulfonylamino)carbonyl)piperidin-1-yl)carbonyl)ethenyl)phenyl] sulfide;

- 10 (2-Isopropylphenyl)[2-nitro-4-(*E*-((3-*p*-toluenesulfonylamino)carbonyl)piperidin-1-yl)carbonyl)ethenyl)phenyl] sulfide;

(2-Isopropylphenyl)[2-nitro-4-(*E*-((3-methyl-4-acetylpiperazin-1-yl)carbonyl)ethenyl)phenyl] sulfide;

(2-Hydroxyphenyl)-[2-chloro-4(*E*-[(morpholin-1-yl)carbonyl]ethenyl)phenyl]sulfide
(1-(Carboxymethyl)indol-5-yl)[2-chloro-4-(*E*-((4-acetylpiperazin-1-

- 15 yl)carbonyl)ethenyl)phenyl] sulfide;

(Benzodioxan-6-yl)[2-trifluoromethyl-4-(*E*-((4-acetylpiperazin-1-yl)carbonyl)ethenyl)phenyl] sulfide;

(2-Isopropylphenyl)[2-nitro-4-(*E*-((3-(1-pyrrolidin-2-onyl)prop-1-ylamino)carbonyl)ethenyl)phenyl] sulfide;

- 20 (3-(2-Morpholinoethylamino)phenyl)[2-trifluoromethyl-4-(*E*-((4-acetylpiperazin-1-yl)carbonyl)ethenyl)phenyl] sulfide;

- (2-Pyrrolidin-1-ylphenyl)[2-nitro-4-(*E*-((4-acetylpiperazin-1-yl)carbonyl)ethenyl)phenyl] sulfide;
- (3-Bromophenyl)[2-nitro-4-(*E*-((3-carboethoxypyrrolidin-1-yl)carbonyl)ethenyl)phenyl] sulfide;
- 5 (3-Bromophenyl)[2-nitro-4-(*E*-((4-carboethoxypyrrolidin-1-yl)carbonyl)ethenyl)phenyl] sulfide;
- (2-(Hydroxymethyl)-benzodioxan-6-yl)[2-chloro-4-(*E*-((4-acetylpiperazin-1-yl)carbonyl)ethenyl)phenyl] sulfide;
- (Benzodioxan-6-yl)[2-trifluoromethyl-4-(*E*-((3-pyrrolidin-2-on-1-yl)prop-1-ylamino)carbonyl)ethenyl)phenyl]sulfide;
- 10 (3-(Dimethylaminomethyl)indol-5-yl)[2-chloro-4-(*E*-((4-acetylpiperazin-1-yl)carbonyl)ethenyl)phenyl] sulfide;
- (2-Isopropylphenyl)[2-nitro-4-(*E*-((2-carboethoxypiperidin-1-yl)carbonyl)ethenyl)phenyl] sulfide;
- 15 (2-Isopropylphenyl)[2-nitro-4-(*E*-((2-carboxypiperidin-1-yl)carbonyl)ethenyl)phenyl] sulfide;
- (2-Isopropylphenyl)[2-nitro-4-(*E*-((4-carboethoxypiperidin-1-yl)carbonyl)ethenyl)phenyl] sulfide;
- (2-Isopropylphenyl)[2-nitro-4-(*E*-((4-carboxypiperidin-1-yl)carbonyl)ethenyl)phenyl] sulfide;
- 20 (2-Isopropylphenyl)[2-nitro-4-(*E*-(((4-*p*-toluenesulfonylaminocarbonyl)piperidin-1-yl)carbonyl)ethenyl)phenyl] sulfide;

0341735-033100

- (2-Isopropylphenyl)[2-nitro-4-(*E*-((3-carboxy-4-hydroxypiperidin-1-yl)carbonyl)ethenyl) phenyl] sulfide;
- (Benzodioxan-6-yl)[2-trifluoromethyl-4-(*E*-((3-carboethoxypiperidin-1-yl)carbonyl)ethenyl) phenyl] sulfide;
- 5 (Benzodioxan-6-yl)[2-trifluoromethyl-4-(*E*-((2-carboethoxypiperidin-1-yl)carbonyl)ethenyl) phenyl] sulfide;
- (Benzodioxan-6-yl)[2-nitro-4-(*E*-((4-carboxypiperidin-1-yl) carbonyl)ethenyl) phenyl] sulfide;
- (Benzodioxan-6-yl)[2-trifluoromethyl-4-(*E*-((3-carboxypyrrolidin-1-yl)carbonyl)ethenyl) phenyl] sulfide;
- 10 (Benzodioxan-6-yl)[2-trifluoromethyl-4-(*E*-((4-carboethoxypiperidin-1-yl)carbonyl)ethenyl) phenyl] sulfide;
- (Benzodioxan-6-yl)[2-trifluoromethyl-4-(*E*-((4-carbomethoxy-4-*tert*-butoxycarbonylpiperazin-1-yl) carbonyl)ethenyl) phenyl] sulfide;
- 15 (Benzodioxan-6-yl)[2-trifluoromethyl-4-(*E*-((2-carbomethoxy-4-methoxycarbonylpiperazin-1-yl) carbonyl)ethenyl) phenyl] sulfide;
- (Benzodioxan-6-yl)[2-trifluoromethyl-4-(*E*-((2-carbomethoxypiperazin-1-yl)carbonyl)ethenyl) phenyl] sulfide;
- (2-Methyl-3-(carboethoxymethyl)indol-5-yl)[2-trifluoromethyl-4-(*E*-((morpholin-1-yl)carbonyl)ethenyl) phenyl] sulfide;
- 20 (1-(2-Methoxyethyl)indol-5-yl)[2-chloro-4-(*E*-((4-acetylpiperazin-1-yl)carbonyl)ethenyl) phenyl] sulfide;

00140 367450

- (2-Isopropylphenyl)[2-nitro-4-(*E*-((3-acetoxymethyl-4-hydroxypiperidin-1-yl)carbonyl)ethenyl) phenyl] sulfide;
- (2-Isopropylphenyl)[2-nitro-4-(*E*-((3-(dimethylaminocarbonyl)-4-hydroxypiperidin-1-yl)carbonyl)ethenyl) phenyl] sulfide;
- 5 (2-Isopropylphenyl)[2-nitro-4-(*E*-((3-cyanomorpholin-1-yl)carbonyl)ethenyl) phenyl] sulfide;
- (2-Isopropylphenyl)[2-nitro-4-(*E*-((3-carboethoxymorpholin-1-yl)carbonyl)ethenyl) phenyl] sulfide;
- (2-Isopropylphenyl)[2-nitro-4-(*E*-((3-(tetrazol-5-yl)morpholin-1-yl)carbonyl)ethenyl) phenyl] sulfide;
- 10 (Benzodioxan-6-yl)[2-trifluoromethyl-4-(*E*-((4-carboxypiperidin-1-yl)carbonyl)ethenyl) phenyl] sulfide;
- (Benzodioxan-6-yl)[2-trifluoromethyl-4-(*E*-((2-carboxypiperidin-1-yl)carbonyl)ethenyl) phenyl] sulfide;
- 15 (Benzodioxan-6-yl)[2-trifluoromethyl-4-(*E*-((4-carbomethoxypiperazin-1-yl)carbonyl)ethenyl) phenyl] sulfide;
- (Benzodioxan-6-yl)[2-trifluoromethyl-4-(*E*-((3-aza-6,9-dioxaspiro[5.4]decan-1-yl)carbonyl)ethenyl) phenyl] sulfide;
- (Benzodioxan-6-yl)[2-trifluoro-4-(*E*-((4-(benzimidazol-1-yl)piperidin-1-yl)carbonyl)ethenyl) phenyl] sulfide;
- 20 (Benzodioxan-6-yl)[2-trifluoromethyl-4-(*E*-((4-(methylaminocarbonyl)piperidin-1-yl)carbonyl)ethenyl) phenyl] sulfide;

00120-5624450

(Benzodioxan-6-yl)[2-trifluoromethyl-4-(*E*-((3-carbomethoxy-4-methoxycarbonylpiperazin-1-yl) carbonyl)ethenyl) phenyl] sulfide;

(2-Isopropylphenyl)[2-nitro-4-(*E*-((3-carboxymorpholin-1-yl)carbonyl)ethenyl) phenyl] sulfide;

- 5 (Benzodioxan-6-yl)[2-trifluoromethyl-4-(*E*-((2-carboxy-4-methoxycarbonylpiperazin-1-yl) carbonyl)ethenyl) phenyl] sulfide;

(Benzodioxan-6-yl)[2-trifluoromethyl-4-(*E*-((morpholin-1-yl)carbonyl)ethenyl) phenyl] sulfide;

- (Benzodioxan-6-yl)[2-trifluoromethyl-4-(*E*-((4-(pyrrolidin-1-yl)piperidin-1-yl)carbonyl)ethenyl) phenyl] sulfide;

- 10 (2-Isopropylphenyl)[2-nitro-4-(*E*-((3-aza-6,9-dioxaspiro[5.4]decan-1-yl)carbonyl)ethenyl) phenyl] sulfide;

(2-Isopropylphenyl)[2-nitro-4-(*E*-((2-(dimethylaminomethyl)piperidin-1-yl)carbonyl)ethenyl) phenyl] sulfide;

- 15 (2-Isopropylphenyl)[2-nitro-4-(*E*-((piperidin-1-ylamino)carbonyl)ethenyl) phenyl] sulfide;

(Benzodioxan-6-yl)[2-trifluoromethyl-4-(*E*-((3-carboxy-4-methoxycarbonylpiperazin-1-yl) carbonyl)ethenyl) phenyl] sulfide;

- (2-(Dimethylaminocarbonyl)-benzodioxan-6-yl)[2-chloro-4-(*E*-((4-acetylpiperazin-1-yl)carbonyl)ethenyl) phenyl] sulfide;;

(2-Isopropylphenyl)[2-nitro-4-(*E*-((3-(2-(methoxymethyl)tetrazol-5-yl) piperidin-1-yl)carbonyl)ethenyl) phenyl] sulfide

(2-Isopropylphenyl)[2-nitro-4-(*E*-((3-(1-(methoxymethyl)tetrazol-5-yl) piperidin-1-yl)carbonyl)ethenyl) phenyl] sulfide;

(1-Methylindol-5-yl)[2-chloro-4-(*E*-((3-(1-pyrrolidin-2-onyl)propylamino) carbonyl)ethenyl) phenyl] sulfide;

- 5 (2-Isopropylphenyl)[2-nitro-4-(*E*-((3-(tetrazol-5-yl) piperidin-1-yl)carbonyl)ethenyl) phenyl] sulfide;

(1-Methylindol-5-yl)[2-chloro-4-(*E*-((3-carboethoxypiperidin-1-yl)carbonyl)ethenyl) phenyl] sulfide

- 10 (1-Methylindol-5-yl)[2-chloro-4-(*E*-((3-carboxypiperidin-1-yl)carbonyl)ethenyl) phenyl] sulfide;

(1-Methylindol-5-yl)[2-chloro-4-(*E*-((4-carboethoxypiperidin-1-yl)carbonyl)ethenyl) phenyl] sulfide;

(1-Methylindol-5-yl)[2-chloro-4-(*E*-((3-carboxypiperidin-1-yl)carbonyl)ethenyl) phenyl] sulfide;

- 15 (2-Isopropylphenyl)[2-nitro-4-(*E*-((2-(1-methylpyrrolidin-2-yl)ethylamino)carbonyl)ethenyl) phenyl] sulfide;

(2-Isopropylphenyl)[2-nitro-4-(*E*-((4-(pyrrolidin-1-yl)piperidin-1-yl)carbonyl)ethenyl) phenyl] sulfide;

(2-Isopropylphenyl)[2-nitro-4-(*E*-((4-sulfopiperidin-1-yl)carbonyl)ethenyl) phenyl]

- 20 sulfide;

(2-Isopropylphenyl)[2-nitro-4-(*E*-((3-hydroxypiperidin-1-yl)carbonyl)ethenyl) phenyl] sulfide;

- (Benzodioxan-6-yl)[2-trifluoromethyl-4-(*E*-((3-
 ((ethanesulfonylamino)carbonyl)piperidin-1-yl) carbonyl)ethenyl) phenyl] sulfide;
 (Benzodioxan-6-yl)[2-trifluoromethyl-4-(*E*-((3-(*p*-
 toluenesulfonylamino)carbonyl)piperidin-1-yl) carbonyl)ethenyl) phenyl] sulfide;
- 5 (Benzodioxan-6-yl)[2-trifluoromethyl-4-(*E*-((4-
 ((ethanesulfonylamino)carbonyl)piperidin-1-yl) carbonyl)ethenyl) phenyl] sulfide;
 (Benzodioxan-6-yl)[2-trifluoromethyl-4-(*E*-((2-tetrazol-5-yl)morpholin-1-
 yl)carbonyl)ethenyl) phenyl] sulfide;
 (2-Isopropylphenyl)[2-nitro-4-(*E*-((2-butyl, 5-(tetrazol-5-yl)morpholin-1-
 yl)carbonyl)ethenyl) phenyl] sulfide;
- 10 (2-(and 3-)(Hydroxymethyl)-benzodioxan-6-yl)[2-nitro-4-(*E*-((4-acetyl)piperazin-1-
 yl)carbonyl)ethenyl) phenyl] sulfide;
 (2-(and 3-)(Hydroxymethyl)-benzodioxan-6-yl)[2-nitro-4-(*E*-((3-(pyrrolidin-2-on-1-
 yl)prop-1-ylamino)carbonyl) ethenyl)phenyl]sulfide;
- 15 (2-(and 3-)(Hydroxymethyl)-benzodioxan-6-yl)[2-trifluoromethyl-4-(*E*-((3-
 (pyrrolidin-2-on-1-yl)prop-1-ylamino)carbonyl) ethenyl)phenyl]sulfide;
 (3-Hydroxymethyl)-benzodioxan-6-yl)[2-nitro-4-(*E*-((3-(pyrrolidin-2-on-1-yl)prop-1-
 ylamino)carbonyl) ethenyl)phenyl]sulfide;
 (Benzodioxan-6-yl)[2-chloro-4-(*E*-((3-carboxypiperidin-1-yl)carbonyl)ethenyl)
 phenyl]
- 20 (2-(and 3-)(Aminomethyl)-benzodioxan-6-yl)[2-trifluoromethyl-4-(*E*-((3-(pyrrolidin-
 2-on-1-yl)prop-1-ylamino)carbonyl) ethenyl)phenyl]sulfide;

00541795.033100

- (2-Isopropylphenyl)[2-nitro-4-(*E*-((3-(methyaminocarbonyl)morpholin-1-yl)carbonyl)ethenyl) phenyl] sulfide;
- (2-Isopropylphenyl)[2-nitro-4-(*E*-((3-(hydroxymethyl)morpholin-1-yl)carbonyl)ethenyl) phenyl] sulfide;
- 5 (2-Isopropylphenyl)[2-nitro-4-(*E*-((3-(acetoxymethyl)morpholin-1-yl)carbonyl)ethenyl) phenyl] sulfide;
- (2-Isopropylphenyl)[2-nitro-4-(*E*-((3-(aminomethyl)morpholin-1-yl)carbonyl)ethenyl) phenyl] sulfide;
- (2-Isopropylphenyl)[2-nitro-4-(*E*-((3-(acetamidomethyl)morpholin-1-yl)carbonyl)ethenyl) phenyl] sulfide;
- 10 (Benzodioxan-6-yl)[2-chloro-4-(*E*-((3-(pyrrolidin-2-on-1-yl)prop-1-ylamino)carbonyl) ethenyl)phenyl]sulfide;
- (Benzodioxan-6-yl)[2-chloro-4-(*E*-((3-carboethoxypiperidin-1-yl) carbonyl)ethenyl) phenyl] sulfide;
- 15 (Benzodioxan-6-yl)[2-chloro-4-(*E*-((2-carboethoxypiperidin-1-yl) carbonyl)ethenyl) phenyl] sulfide
- (2-Methoxyphenyl)-[2,3-dichloro-4(*E*-[(morpholin-1-yl)carbonyl]ethenyl)phenyl] sulfide;
- (2-Methoxyphenyl)-[2,3-dimethyl-4(*E*-[(morpholin-1-yl)carbonyl]ethenyl)phenyl]
- 20 sulfide;
- (2-Isopropylphenyl)[2-nitro-4-(*E*-((indol-5-ylamino)carbonyl)ethenyl) phenyl] sulfide;

- (Benzodioxan-6-yl)[2-chloro-4-(*E*-((3-carboxypiperidin-1-yl) carbonyl)ethenyl) phenyl] sulfide;
- (Benzodioxan-6-yl)[2-chloro-4-(*E*-((3-(tetrazol-5-yl)piperidin-1-yl) carbonyl)ethenyl) phenyl] sulfide;
- 5 (Benzodioxan-6-yl)[2-chloro-4-(*E*-((4-*tert*-butoxycarbonyl)piperazin-1-yl) carbonyl)ethenyl) phenyl] sulfide;
- (Benzodioxan-6-yl)[2-chloro-4-(*E*-((2-carboxypiperidin-1-yl) carbonyl)ethenyl) phenyl] sulfide;
- (Benzodioxan-6-yl)[2-chloro-4-(*E*-((3-(tetrazol-5-yl)morpholin-1-yl) carbonyl)ethenyl) phenyl] sulfide;
- 10 (Benzodioxan-6-yl)[2-chloro-4-(*E*-((4-(methylaminocarbonyl)piperazin-1-yl) carbonyl)ethenyl) phenyl] sulfide;
- (2-Methoxyphenyl)-[2,3-dichloro-4(*E*-[(4-carboxypiperidin-1-yl)carbonyl]ethenyl) phenyl] sulfide;
- 15 (Benzodioxan-6-yl)[2-chloro-4-(*E*-((4-(tetrazol-5-yl)piperidin-1-yl) carbonyl)ethenyl) phenyl] sulfide;
- (2-Methoxyphenyl)-[3-chloro-4(*E*-[(morpholin-1-yl)carbonyl]ethenyl)phenyl]sulfide;
- (2-Isopropylphenyl)[2-nitro-4-(*E*-((4-oxopiperidin-1-yl)carbonyl)ethenyl) phenyl] sulfide;
- 20 (Benzodioxan-6-yl)[2-trifluoromethyl-4-(*E*-((3-*R*-carboethoxypiperidin-1-yl)carbonyl)ethenyl) phenyl] sulfide;

00120.5674560

(Benzodioxan-6-yl)[2-trifluoromethyl-4-(*E*-((3-*R*-carboxypiperidin-1-yl)carbonyl)ethenyl) phenyl] sulfide;

(Benzodioxan-6-yl)[2,3-dichloro-4-(*E*-((3-pyrrolidin-2-on-1-yl)prop-1-ylamino)carbonyl) ethenyl)phenyl]sulfide;

- 5 (Benzodioxan-6-yl)[2,3-dichloro-4-(*E*-((4-acetylpiperazin-1-yl) carbonyl)ethenyl) phenyl] sulfide;

(Benzodioxan-6-yl)[2,3-dichloro-4-(*E*-((3-carboethoxypiperidin-1-yl) carbonyl)ethenyl) phenyl] sulfide;

- 10 (Benzodioxan-6-yl)[2,3-dichloro-4-(*E*-((4-carboethoxypiperidin-1-yl) carbonyl)ethenyl) phenyl] sulfide;
- (Benzodioxan-6-yl)[2,3-dichloro-4-(*E*-((3-carboxypiperidin-1-yl) carbonyl)ethenyl) phenyl] sulfide;

(Benzodioxan-6-yl)[2,3-dichloro-4-(*E*-((4-carboxypiperidin-1-yl) carbonyl)ethenyl) phenyl] sulfide;

- 15 (2-Isopropylphenyl)[2,3-dichloro-4-(*E*-((3-(1-pyrrolidin-2-onyl)propylamino) carbonyl)ethenyl) phenyl] sulfide;

(2-Isopropylphenyl)[2,3-dichloro-4-(*E*-((4-acetylpiperazin-1-yl) carbonyl)ethenyl) phenyl] sulfide;

(2-Isopropylphenyl)[2,3-dichloro-4-(*E*-((3-carboethoxypiperidin-1-yl)

- 20 carbonyl)ethenyl) phenyl] sulfide;

(2-Isopropylphenyl)[2,3-dichloro-4-(*E*-((4-carboethoxypiperidin-1-yl) carbonyl)ethenyl) phenyl] sulfide;

00541795,032100

(2-Isopropylphenyl)[2,3-dichloro-4-(*E*-((3-carboxypiperidin-1-yl) carbonyl)ethenyl) phenyl] sulfide;

(2-Isopropylphenyl)[2,3-dichloro-4-(*E*-((4-carboxypiperidin-1-yl) carbonyl)ethenyl) phenyl] sulfide;

- 5 (1-Methylindol-5-yl)[2,3-dichloro-4-(*E*-((3-carboethoxypiperidin-1-yl) carbonyl)ethenyl) phenyl] sulfide;

(1-Methylindol-5-yl)[2,3-dichloro-4-(*E*-((3-carboxypiperidin-1-yl) carbonyl)ethenyl) phenyl] sulfide;

(1-Methylindol-5-yl)[2,3-dichloro-4-(*E*-((4-carboethoxypiperidin-1-yl)

- 10 carbonyl)ethenyl) phenyl] sulfide;

(1-Methylindol-5-yl)[2,3-dichloro-4-(*E*-((4-carboxypiperidin-1-yl) carbonyl)ethenyl) phenyl] sulfide;

(2-Ethoxyphenyl)-[2,3-dichloro-4(*E*-[(4-carboxypiperidin-1-yl)carbonyl]ethenyl) phenyl] sulfide;

- 15 (2-Ethoxyphenyl)-[2,3-dichloro-4(*E*-[(morpholin-1-yl)carbonyl]ethenyl)phenyl] sulfide;

(2-Ethoxyphenyl)-[2,3-dichloro-4(*E*-[(3-carboxypiperidin-1-yl)carbonyl]ethenyl) phenyl] sulfide;

(2-Isopropylphenyl)[2-nitro-4-(*E*-((3-carboethoxypyrrolidin-1-yl)carbonyl)ethenyl)

- 20 phenyl] sulfide;

(2-Isopropylphenyl)[2-nitro-4-(*E*-((3-carboxypyrrolidin-1-yl)carbonyl)ethenyl) phenyl] sulfide;

001230.5624560

(2-Isopropylphenyl)[2,3-difluoro-4-(*E*-((3-carboethoxypiperidin-1-yl)carbonyl)ethenyl) phenyl] sulfide;

(2-Isopropylphenyl)[2,3-difluoro-4-(*E*-((3-carboxypiperidin-1-yl)carbonyl)ethenyl) phenyl] sulfide;

- 5 (2-Isopropylphenyl)[2,3-difluoro-4-(*E*-((4-carboxypiperidin-1-yl)carbonyl)ethenyl) phenyl] sulfide;

(Benzodioxan-6-yl)[2-trifluoromethyl-4-(*E*-((3-ethoxycarbonylpyrrolidin-1-yl)carbonyl)ethenyl) phenyl] sulfide

- (Benzodioxan-6-yl)[2-trifluoromethyl-4-(*E*-((3-carboxypyrrolidin-1-yl)carbonyl)ethenyl) phenyl] sulfide;

- 10 (2-Methoxyphenyl)[2-chloro-3-trifluoromethyl-4-(*E*-((4-carboethoxypiperidin-1-yl)carbonyl)ethenyl) phenyl] sulfide;

(2-Methoxyphenyl)[2-chloro-3-trifluoromethyl-4-(*E*-((4-carboethoxypiperidin-1-yl)carbonyl)ethenyl) phenyl] sulfide;

- 15 (2-Methoxyphenyl)[2-chloro-3-trifluoromethyl-4-(*E*-((morpholin-1-yl)carbonyl)ethenyl) phenyl] sulfide;

(Benzodioxan-6-yl) [4-(*E*-((4-carboxypiperidin-1-yl) carbonyl)ethenyl)naphthyl] sulfide;

(2-Methoxyphenyl) [2,3-dichloro-4-(*E*-((4-(spiro-hydantoin-5-yl)-piperidin-1-

- 20 yl)carbonyl)ethenyl)phenyl] sulfide;

(2-Methoxyphenyl) [2,3-dichloro-4-(*E*-4-(2-(2-hydroxyethoxy)ethyl)piperazin-1-yl)carbonyl)ethenyl)phenyl] sulfide;

- (2-Methoxyphenyl)[2,3-dichloro-4-(*E*-((4-ethylpiperazin-1-yl)carbonyl)ethenyl)phenyl] sulfide;
- (2-Isopropylphenyl)[2,3-dichloro-4-(*E*-((4-(2-(2-hydroxyethoxy)ethyl)piperazin-1-yl)carbonyl)ethenyl)phenyl] sulfide;
- 5 (Benzodioxan-6-yl)[2,3-bis(trifluoromethyl)-4-(*E*-((4-carboxypiperidin-1-yl)carbonyl)ethenyl)phenyl]sulfide;
- (2-Methoxyphenyl) [2,3-dichloro-4-(*E*-((4-(carboxymethylamino)carbonyl-piperidin-1-yl)carbonyl)ethenyl)phenyl] sulfide;
- (2-Methoxyphenyl) [2,3-bis(trifluoromethyl)-4-(*E*-((4-carboxymethylpiperazin-1-yl)carbonyl)ethenyl)phenyl]sulfide;
- 10 (2-Methoxyphenyl) [2,3-bis(trifluoromethyl)-4-(*E*-((4-N-(2-hydroxyethyl)piperazin-1-yl)carbonyl)ethenyl)phenyl]sulfide;
- (1-Methylindol-5-yl) [2,3-dichloro-4-(*E*-((4-(carbo-2,3-dihydroxypropylamino)piperidin-1-yl)carbonyl)ethenyl)phenyl] sulfide;
- 15 (2-Methoxyphenyl) [2,3-dichloro-4-(*E*-4-(2,3-dihydroxypropionyl)piperazin-1-yl)carbonyl)ethenyl)phenyl] sulfide;
- (2-Methoxyphenyl) [2,3-dichloro-4-(*E*-4-(2,3-dihydroxy-3-carboxypropionyl)piperazin-1-yl)carbonyl)ethenyl)phenyl] sulfide;
- (1-Methylindol-5-yl) [2,3-dichloro-4-(*E*-((4-(carboxymethylamino)carbonyl-piperidin-1-yl)carbonyl)ethenyl)phenyl] sulfide;
- 20 (1-Methylindol-5-yl) [2,3-dichloro-4-(*E*-((4-sulfopiperidin-1-yl)carbonyl)ethenyl)phenyl] sulfide;

09541795-032100

- (1-Methylindol-5-yl) [2,3-dichloro-4-(*E*-(4-methylhomopiperazin-1-ylcarbonyl)ethenyl)phenyl] sulfide;
- (1-Methylindol-5-yl) [2,3-dichloro-4-(*E*-(4-tetrahydrofuroyl)piperazin-1-ylcarbonyl)ethenyl)phenyl] sulfide;
- 5 (2-Methoxyphenyl) [2,3-dichloro-4-(*E*-((4-amino-4-carboxypiperidin-1-ylcarbonyl)ethenyl)phenyl] sulfide;
- (2-Methoxyphenyl)[2,3-dichloro-4-((4-furoylpiperazin-1-ylcarbonyl)ethenyl)phenyl] sulfide;
- (1-Methylindol-5-yl) [2,3-dichloro-4-(*E*-(4-(carbo-3-sulfopropylamino)piperadin-1-ylcarbonyl)ethenyl)phenyl] sulfide;
- 10 (2-Methoxyphenyl)[2,3-dichloro-4-(*E*-(4-acetylamino-4-carboxypiperidin-1-ylcarbonyl)ethenyl)phenyl] sulfide;
- (2-Methoxyphenyl) [2,3-bis(trifluoromethyl)-4-(*E*-((4-carboxypiperidin-1-ylcarbonyl)ethenyl)phenyl]sulfide;
- 15 (2-Methoxyphenyl) 5-[8-(*E*-((4-(aminocarbonyl)piperidin-1-ylcarbonyl)ethenyl)quinoliny]sulfide;
- (2-Methoxyphenyl) [2-trifluoromethyl-4-(*E*-((4-carboxypiperidin-1-ylcarbonyl)ethenyl)phenyl]sulfide;
- (1-Methylindol-5-yl) [2,3-dichloro-4-(*E/Z*-((1*S*,4*S*)-2,5-diazabicyclo(2,2,1)heptan-2-ylcarbonyl)ethenyl)-2,3-dichlorophenyl] sulfide;
- 20 (1-Methylindol-5-yl) [2,3-dichloro-4-(*E*-(4-hydroxy-3-carboxypiperadin-1-ylcarbonyl)ethenyl)phenyl] sulfide;;

- (1-Methylindol-5-yl) [2,3-dichloro-4-(*E*-(S-oxothiomorpholin-1-ylcarbonyl)ethenyl)phenyl] sulfide;
- (2-Methoxyphenyl) [2,3-dichloro-4-(*E*-((4-sulfophenylamino)carbonyl)ethenyl)phenyl] sulfide;
- 5 (2-Methoxyphenyl) [2,3-dichloro-4-(*E*-((4-carboxyphenylamino)carbonyl)ethenyl)phenyl] sulfide;
- [3-(4-Morpholino)phenyl] [2,3-dichloro-4-(*E*-[(4-carboxypiperidin-1-yl)carbonyl]ethenyl)phenyl] sulfide;
- [3-(4-Morpholino)phenyl] [2,3-dichloro-4-(*E*-((4-carboxypiperidin-1-yl)carbonyl)ethenyl)phenyl] sulfide;
- 10 (2-Methoxyphenyl)[2,3-bis(trifluoromethyl)-4-(*E*-((4-phenylcarboxypiperidin-1-yl)carbonyl)ethenyl)phenyl]sulfide;
- (2-Methoxyphenyl) [2,3-dichloro-4-(*E*-(((4-hydroxylaminocarbonyl)piperidin-1-yl)carbonyl)ethenyl)phenyl] sulfide;
- 15 (2-Methoxyphenyl) [2,3-dichloro-4-(*E*-((N-carboxymethyl-N-phenylamino)carbonyl)ethenyl)phenyl] sulfide;

0054795-03100
DATE 09-26-2019

- (2-Methoxyphenyl) [3-chloro-6-hydroxy-4-(E-((3-carboxypiperidin-1-yl)carbonyl)ethenyl)phenyl]sulfide;
- (2-Methoxyphenyl) [2,3-dichloro-4-(E-(4-((1-(2-phenyl-1-carboxyethyl)amino)carbonyl)piperidin-1-yl)carbonyl)ethenyl)phenyl]sulfide;
- 5 (2-Methoxyphenyl) [2,3-dichloro-4-(E-(4-((1-(2-hydroxy-1-carboxyethyl)amino)carbonyl)piperidin-1-yl)carbonyl)ethenyl)phenyl]sulfide;
- (3-(1-(3-Carboxypiperidinyl)phenyl)[2,3-dichloro-4-(E-((1,2,5,6-tetrahydropyridin-1-yl)carbonyl)ethenyl)phenyl]sulfide;
- (3-(4-Pyrrolidin-1-yl)piperidin-1-yl)phenyl [2,3-dichloro-4-(E-(((3-(2-pyrrolidinon-1-yl)propylamino)carbonyl)ethenyl)phenyl]sulfide;
- 10 [3-(4-(Spiro-2,2-dioxolanyl)piperidin-1-yl)phenyl] [2,3-dichloro-4-(E-((4-morpholinyl)carbonyl)ethenyl)phenyl]sulfide;
- [3-(3-Carboxypiperidin-1-yl)phenyl] [2,3-dichloro-4-(E-[(4-carboxypiperidin-1-yl)carbonyl]ethenyl)phenyl] sulfide;
- 15 (2-(2-Carboxy)ethenyl)phenyl [2,3-dichloro-4-(E-((4-morpholinyl)carbonyl)ethenyl)phenyl]sulfide;
- [3-(4-Carboxylpiperidin-1-yl)phenyl] [2,3-dichloro-4-(E-[(1,2,3,6-tetrahydropyridine)-1-yl)carbonyl]ethenyl)phenyl] sulfide;
- [3-(4-Carboxylpiperidinyl)phenyl] [2,3-dichloro-4-(E-[(4-morpholinyl)carbonyl]ethenyl)phenyl] sulfide;
- 20 [2-(4-Acetyl)piperazin-1-yl)phenyl] [2,3-dichloro-4-(E-[(4-carboxypiperidin-1-yl)carbonyl]ethenyl)phenyl] sulfide;

00541795-032100

- 3-(3-Carboxypiperidin-1-yl)phenyl] [2,3-dichloro-4-(*E*-[(4-morpholinyl)carbonyl]ethenyl)phenyl] sulfide;
- [3-(4-Carboxypiperidin-1-yl)phenyl] [2,3-dichloro-4-(*E*-[(4-(dimethylaminosulfamoyl)piperazin-1-yl)carbonyl]ethenyl)phenyl] sulfide;
- 5 (2-Methoxyphenyl)[2,3-bis(trifluoromethyl)-4-(*E*-[(3-carboxypiperidin-1-yl)carbonyl]ethenyl)phenyl]sulfide;
- (2-Methoxyphenyl) [2,3-bis(trifluoromethyl)-4-(*E*-[(2-carboxypyrrolidin-1-yl)carbonyl]ethenyl)phenyl]sulfide;
- [3-(4-Carboxypiperidin-1-yl)phenyl] [2,3-dichloro-4-(*E*-[(4-((trifluoromethylsulfonyl)piperazin-1-yl)carbonyl]ethenyl)phenyl] sulfide;
- 10 (2-Methoxyphenyl)[2,3-dichloro-4-(*E*-(piperidin-1-ylcarbonyl)ethenyl)phenyl] sulfide;
- (2-Hydroxyphenyl) [2,3-dichloro-4-(*E*-[(4-morpholino)carbonyl]ethenyl)phenyl]sulfide;
- 15 (2-Methoxyphenyl) [2,3-dichloro-4-(*E*-[(4-carboxyphenyl)methyl]amino)carbonyl]ethenyl)phenyl]sulfide;
- (2-Methoxyphenyl) [2,3-dichloro-4-(*E*-[(4-pyrrolidin-1-yl)piperidin-1-yl)carbonyl]ethenyl)phenyl]sulfide;
- (2-Hydroxyphenyl)[2,3-dichloro-4-(*E*-[(4-carboxypiperidin-1-yl)carbonyl]ethenyl)phenyl]sulfide;
- 20 [3-(4-Carboxypiperidin-1-yl)phenyl] [2,3-dichloro-4-(*E*-[(4-(methylsulfonyl)piperazin-1-yl)carbonyl]ethenyl)phenyl] sulfide;

- (2-Aminophenyl) [2,3-dichloro-4-(*E*-((4-morpholinyl)carbonyl)ethenyl)phenyl] sulfide;
- (3-(4-carboxypiperidin-1-yl)phenyl)[2,3-dichloro-4-(*E*-((S-oxothiomorpholin-1-yl)carbonyl)ethenyl)phenyl] sulfide;
- 5 [3-(4-Carboxypiperidin-1-yl)phenyl] [2,3-dichloro-4-(*E*-((4-hydroxypiperidin-1-yl)carbonyl)ethenyl)phenyl] sulfide;
- (2-Glycoxyphenyl) [2,3-dichloro-4-(*E*-((4-morpholino)carbonyl)ethenyl)phenyl] sulfide;
- (2-(4-Butyroxyl)phenyl)[2,3-dichloro-4-(*E*-((4-morpholino)carbonyl)ethenyl)phenyl] sulfide;
- 10 [3-(4-Carboxypiperidin-1-yl)phenyl] [2,3-dichloro-4-(*E*-((4-hydroxyethyl)piperazin-1-yl)carbonyl)ethenyl)phenyl] sulfide;
- [3-(4-Carboxypiperidin-1-yl)phenyl] [2,3-dichloro-4-(*E*-((4-furoyl)piperazin-1-yl)carbonyl)ethenyl)phenyl] sulfide;
- 15 [3-(4-Carboxypiperidin-1-yl)phenyl] [2,3-dichloro-4-(*E*-((pyrrolidin-1-yl)carbonyl)ethenyl)phenyl] sulfide;
- [3-(4-Carboxypiperidin-1-yl)phenyl] [2,3-dichloro-4-(*E*-((diethylaminocarbonyl)ethenyl)phenyl] sulfide;
- [3-(4-Carboxypiperidin-1-yl)phenyl] [2,3-dichloro-4-(*E*-((4-ethylpiperazin-1-yl)carbonyl)ethenyl)phenyl] sulfide;
- 20 [3-(4-Carboxypiperidin-1-yl)phenyl] [2,3-dichloro-4-(*E*-((4-aminocarbonyl)piperidin-1-yl)carbonyl)ethenyl)phenyl] sulfide;

00544795 033100
007100 56714560

- [3-(4-Carboxypiperidin-1-yl)phenyl] [2,3-dichloro-4-(*E*-((4-(2-ethoxyethyl)piperidin-1-yl)carbonyl)ethenyl)phenyl] sulfide;
- [3-((4-Carboxymethyl)piperazin-1-yl)phenyl] [(2,3-dichloro-4-(*E*-(4-morpholinyl)carbonyl)ethenyl)phenyl] sulfide;
- 5 [3-(4-Carboxypiperidin-1-yl)phenyl] [2,3-bis(trifluoromethyl)-4-(*E*-(4-morpholino)carbonyl)ethenyl)phenyl]sulfide;
- (3-Hydroxyphenyl) [2,3-dichloro-4-(*E*-(4-morpholino)carbonyl)ethenyl)phenyl]sulfide;
- [3-(4-Butyroxyl)phenyl] [2,3-dichloro-4-(*E*-((4-morpholino)carbonyl)ethenyl)phenyl]sulfide;
- 10 (2-Hydroxyphenyl) [2,3-bis(trifluoromethyl)-4-(*E*-(4-morpholino)carbonyl)ethenyl)phenyl]sulfide;
- (3-Hydroxyphenyl) [2,3-bis(trifluoromethyl)-4-(*E*-(4-morpholino)carbonyl)ethenyl)phenyl]sulfide;
- 15 [3-(4-Carboxypiperidin-1-yl)phenyl] [2,3-ditrifluoromethyl-4-(*E*-((4-hydroxypiperidin-1-yl)carbonyl)ethenyl)phenyl] sulfide;
- [3-(4-Carboxypiperidin-1-yl)phenyl] [2,3-ditrifluoromethyl-4-(*E*-((1,2,5,6-tetrahydropyridin-1-yl)carbonyl)ethenyl)phenyl] sulfide;

001460-5674560

- [2-((4-Carboxy)butyloxy)phenyl] [2,3-dichloro-4-(E-((4-morpholino)carbonyl)ethenyl)phenyl]sulfide;
- (2-Glycoxyphenyl) [2,3-bis(trifluoromethyl)-4-(E-((4-morpholino)carbonyl)ethenyl)phenyl]sulfide;
- 5 (2-(4-Butyroxyl)phenyl)[2,3-bis(trifluoromethyl)-4-(E-((4-morpholino)carbonyl)ethenyl)phenyl]sulfide;
- [3-(4-Carboxypiperidin-1-yl)phenyl] [2,3-ditrifluoromethyl-4-(E-((bis-(2-ethoxyethyl)amino)carbonyl)ethenyl)phenyl] sulfide;
- [3-(4-Carboxypiperidin-1-yl)phenyl] [2,3-bis-(trifluoromethyl)-4-(E-((bis-(2-hydroxypropyl)amino)carbonyl)ethenyl)phenyl] sulfide;
- 10 [3-(4-Carboxypiperidin-1-yl)phenyl] [2,3-bis-(trifluoromethyl)-4-(E-(piperazin-1-yl)carbonyl)ethenyl)phenyl] sulfide;
- (3-(4-Butyroxyl)phenyl)[2,3-bis(trifluoromethyl)-4-(E-((4-morpholino)carbonyl)ethenyl)phenyl]sulfide;
- 15 [2-(3-Carboxypiperidin-1-yl)phenyl] [2,3-dichloro-4-(E-[(3-(2-pyrrolidinon-1-yl)propylaminocarbonyl)ethenyl)phenyl] sulfide;
- [2-(3-Carboxypiperidin-1-yl)phenyl] [2,3-bis(trifluoromethyl)-4-(E-[(3-(2-pyrrolidinon-1-yl)propylaminocarbonyl)ethenyl)phenyl] sulfide;
- [2-(3-Carboxypiperidin-1-yl)phenyl] [2,3-dichloro-4-(E-((4-(2-hydroxyethyl)piperazin-1-yl)carbonyl)ethenyl)phenyl] sulfide;
- 20 [2-(3-Carboxypiperidin-1-yl)phenyl] [2,3-bis(trifluoromethyl)-4-(E-((1,2,5,6-tetrahydropyridin-1-yl)carbonyl)ethenyl)phenyl] sulfide;

00541795-033100

- [2-(3-Carboxypiperidin-1-yl)phenyl] [2,3-bis(trifluoromethyl)-4-(*E*-((4-(2-hydroxyethyl)piperazin-1-yl)carbonyl)ethenyl)phenyl] sulfide;
- [2-(3-Carboxypiperidin-1-yl)phenyl] [2,3-bis(trifluoromethyl)-4-(*E*-((4-(2-(hydroxyethoxy)ethyl)piperazin-1-yl)carbonyl)ethenyl)phenyl] sulfide; and
- 5 (3-(3-Propioxy)phenyl) [2,3-dichloro-4-(*E*-((4-morpholino)carbonyl)ethenyl)phenyl] sulfide.
13. A compound according to Claim 1 selected from the group consisting of:
- 10 (2-Formylphenyl)[2-nitro-4-(*E*-((4-acetyl)piperazin-1-yl)carbonyl)ethenyl]phenyl sulfide;
- (2-Ethoxyphenyl)-[2-chloro-4-(*E*-[(morpholin-1-yl)carbonyl]ethenyl)phenyl] sulfide;
- (2-Isopropylphenyl)[2-nitro-4-(*E*-((3-dimethylaminocarbonyl-4-acetyl)piperazin-1-yl)carbonyl)ethenyl] phenyl] sulfide;
- 15 (2-Isopropylphenyl)[2-nitro-4-(*E*-((4-methoxycarbonyl)piperazin-1-yl)carbonyl)ethenyl] phenyl] sulfide;
- (2-Ethoxyphenyl)[2-trifluoromethyl-4-(*E*-((2-carboxy-4-(methoxycarbonyl)piperazin-1-yl)carbonyl)ethenyl] phenyl] sulfide;
- (2-Ethyl-4-bromophenyl)[2-nitro-4-(*E*-((4-acetyl)piperazin-1-yl)carbonyl)ethenyl] phenyl] sulfide;
- 20 (3-Morpholinophenyl)[2-nitro-4-(*E*-((4-acetyl)piperazin-1-yl)carbonyl)ethenyl] phenyl] sulfide;

(2-Isopropylphenyl)[2-nitro-4-(*E*-((3-carboethoxypiperidin-1-yl)carbonyl)ethenyl)phenyl] sulfide;

(2-Isopropylphenyl)[2-nitro-4-(*E*-((3-(1-pyrrolidin-2-onyl)prop-1-ylamino)carbonyl)ethenyl)phenyl] sulfide;

- 5 (2-(Hydroxymethyl)-benzodioxan-6-yl)[2-chloro-4-(*E*-((4-acetyl)piperazin-1-yl)carbonyl)ethenyl)phenyl] sulfide;

(Benzodioxan-6-yl)[2-trifluoromethyl-4-(*E*-((3-carboxypyrrolidin-1-yl)carbonyl)ethenyl)phenyl] sulfide;

- (Benzodioxan-6-yl)[2-trifluoromethyl-4-(*E*-((4-carboethoxypiperidin-1-yl)carbonyl)ethenyl)phenyl] sulfide;

- 10 (Benzodioxan-6-yl)[2-trifluoromethyl-4-(*E*-((2-carbomethoxy-4-methoxycarbonylpiperazin-1-yl)carbonyl)ethenyl)phenyl] sulfide;

(Benzodioxan-6-yl)[2-trifluoromethyl-4-(*E*-((2-carbomethoxypiperazin-1-yl)carbonyl)ethenyl)phenyl] sulfide;

- 15 (Benzodioxan-6-yl)[2-chloro-4-(*E*-((4-(methylaminocarbonyl)piperazin-1-yl)carbonyl)ethenyl)phenyl] sulfide;

(2-Methoxyphenyl)-[2,3-dichloro-4-(*E*-((4-carboxypiperidin-1-yl)carbonyl)ethenyl)phenyl] sulfide;

(Benzodioxan-6-yl)[2-trifluoromethyl-4-(*E*-((3-*R*-carboethoxypiperidin-1-

- 20 yl)carbonyl)ethenyl)phenyl] sulfide;

(Benzodioxan-6-yl)[2,3-dichloro-4-(*E*-((3-(pyrrolidin-2-on-1-yl)prop-1-ylamino)carbonyl)ethenyl)phenyl] sulfide;

00541795-033100

(Benzodioxan-6-yl)[2,3-dichloro-4-(*E*-((4-acetyl)piperazin-1-yl) carbonyl)ethenyl) phenyl] sulfide;

(Benzodioxan-6-yl)[2,3-dichloro-4-(*E*-((3-carboethoxypiperidin-1-yl) carbonyl)ethenyl) phenyl] sulfide;

- 5 (Benzodioxan-6-yl)[2,3-dichloro-4-(*E*-((4-carboethoxypiperidin-1-yl) carbonyl)ethenyl) phenyl] sulfide;

(Benzodioxan-6-yl)[2,3-dichloro-4-(*E*-((3-carboxypiperidin-1-yl) carbonyl)ethenyl) phenyl] sulfide;

- 10 (Benzodioxan-6-yl)[2,3-dichloro-4-(*E*-((4-carboxypiperidin-1-yl) carbonyl)ethenyl) phenyl] sulfide;

(2-Isopropylphenyl)[2,3-dichloro-4-(*E*-((4-acetyl)piperazin-1-yl) carbonyl)ethenyl) phenyl] sulfide;

(2-Isopropylphenyl)[2,3-dichloro-4-(*E*-((3-carboethoxypiperidin-1-yl) carbonyl)ethenyl) phenyl] sulfide;

- 15 (2-Isopropylphenyl)[2,3-dichloro-4-(*E*-((3-carboxypiperidin-1-yl) carbonyl)ethenyl) phenyl] sulfide;

(2-Isopropylphenyl)[2,3-dichloro-4-(*E*-((4-carboxypiperidin-1-yl) carbonyl)ethenyl) phenyl] sulfide;

(2-Methoxyphenyl) [2,3-dichloro-4-(*E*-((4-(carboxymethylamino)carbonyl-piperidin-1-yl)carbonyl)ethenyl)phenyl] sulfide;


- 20 (2-Methoxyphenyl) [2,3-bis(trifluoromethyl)-4-(*E*-((4-carboxymethyl)piperazin-1-yl) carbonyl)ethenyl)phenyl]sulfide; and

(2-Methoxyphenyl) [2,3-bis(trifluoromethyl)-4-(E-((4-N-(2-hydroxyethyl)piperazin-1-yl)carbonyl)ethenyl)phenyl]sulfide.

Insert new compounds here

- 5 14. A compound according to Claim 1 wherein **Ar** is selected from the group consisting of methoxyphenyl and isopropylphenyl.
15. A compound according to Claim 1 wherein **Ar** is benzodioxan or substituted benzodioxan.
- 10 16. A compound according to Claim 1 wherein R_3 is a "trans-cinnamide"; and **Ar** is selected from 1,3-benzimidazol-2-one, 1,4-benzodioxane, 1,3-benzodioxole, 1-benzopyr-2-en-4-one, indole, isatin, phenyl, 1,3-quinazolin-4-one, and quinoline.
- 15 17. A compound according to Claim 1 wherein R_{10} and R_{11} are independently selected from hydrogen, alkyl, cycloalkyl, alkoxycarbonylalkyl, hydroxyalkyl, and heterocyclalkyl.
18. A compound according to Claim 1 wherein $NR_{10}R_{11}$ is heterocyclyl or substituted heterocyclyl.
- 20

19. A pharmaceutical composition comprising a compound of Claim 1 in a pharmaceutically-acceptable carrier.
20. A method of inhibiting inflammation comprising the administration of a compound of Claim 1 to a mammal in need of treatment.
21. A method of inhibiting inflammation comprising the administration of a composition of Claim 12 to a mammal in need of treatment.
22. A method of suppressing immune response comprising the administration of a compound of Claim 1 to a mammal in need of treatment.
23. A method of suppressing immune response comprising the administration of a composition of Claim 12 to a mammal in need of treatment.
24. A compound of formula II




wherein R_1 and R_2 are independently selected from the group consisting of

- i. hydrogen,
- j. halogen,
- 5 k. alkyl,
- l. haloalkyl,
- m. alkoxy,
- n. cyano,
- o. nitro, and
- 10 p. carboxaldehyde.

007E0-5674560


25. A process for preparing a compound of formula II



II

which comprises:

- 5 a) reacting a compound of formula II'



II'

with lithium hydroxide, and

10

- b) cleaving the resulting methyl ether.

26. The compound of Claim 1 with the proviso that when three of R₁, R₂, R₄, and R₅ are hydrogen, then the remaining R₁, R₂, R₄, or R₅ is not carboxyl, 5-tetrazolyl, hydroxymethyl or carboxyl derivatized in the form of a pharmaceutically acceptable ester.

15

Abstract

The present invention relates to novel cinnamide compounds that are useful for treating inflammatory and immune diseases, to pharmaceutical compositions
5 containing these compounds, and to methods of inhibiting inflammation or suppressing immune response in a mammal.

001100-56714560

PATENT

IN THE UNITED STATES PATENT & TRADEMARK OFFICE

APPLICANT: J. Link, et al.] Certificate of Mailing (37 C.F.R. §1.8(a))
]]
SERIAL NO.:] I hereby certify that this paper (along with any
] paper referred to as being attached or enclosed) is
FILED:] being deposited with the United States Postal
] Service on the date shown below with sufficient
] postage as First Class Mail, in an enveloped
] addressed to:
FOR: CELL ADHESION-INHIBITING]
] ANTIINFLAMMATORY AND]
] IMMUNE-SUPPRESSIVE]
] COMPOUNDS]
] Box Missing Parts
] Assistant Commissioner for Patents
] Washington, D.C.
EXAMINER: (not yet assigned)]
] Date of Deposit: March 31, 2000
]]
CASE NO.: 6446.US.P2]
] *Katrina M Williams 3-31-00*
] Katrina Williams Date
GROUP ART UNIT: (not yet assigned)]

**DECLARATION AND POWER OF ATTORNEY
FOR A UNITED STATES PATENT APPLICATION**

As a below-named inventor, I hereby declare:

My residence, post office address and citizenship are as stated below next to my name. I believe I am an original and first and joint inventor of the subject matter which is claimed and for which a patent is sought on the invention entitled CELL ADHESION-INHIBITING ANTIINFLAMMATORY AND IMMUNE-SUPPRESSIVE COMPOUNDS, the specification of which was filed April 2, 1999.

I hereby state that I have reviewed and understand the contents of the above-mentioned specification, including the claims.

I acknowledge a duty to disclose to the Patent and Trademark Office all information known to me to be material to patentability as defined in Title 37, Code of Federal Regulations, §1.56.

Claim to benefit of foreign application(s) as follows:

I hereby claim foreign priority benefits under 35 U.S.C. §119 for the following foreign applications for patent or inventors certificate.

NONE

00541795.03300

The following foreign applications for patent or inventor's certificate have a filing date earlier than the filing date of the applications identified above.

NONE

Claim to benefit of earlier U.S. application(s) as follows:

I hereby claim the benefit under 35 U.S.C. §120 of the following earlier-filed United States patent applications. Insofar as the subject matter of each of the claims of this application is not disclosed in the prior U.S. applications in the manner required by 35 U.S.C. § 112, first paragraph, I acknowledge a duty to disclose to the Patent and Trademark Office all information known to me to be material to patentability as defined in 37 C.F.R. §1.56 which came into existence between the filing date(s) of the prior applications and the national or PCT filing date of this application.

U.S. Serial No. 09/474,517, filed December 29, 1999.

I hereby appoint the following Attorneys and/or agents to prosecute this application and any continuation or divisional applications based hereon, and to transact all business in the Patent and Trademark Office connected therewith:

Regina M. Anderson, Reg. No. 35,820
Cheryl L. Becker, Reg. No. 35,441
Thomas D. Brainard, Reg. No. 32,459
Dianne Casuto, Reg. No. 40,943
Portia Chen, Reg. No., 44,075
Steven R. Crowley, Reg. No. 31,604
Andreas M. Danckers, Reg. No. 32,652
J. Michael Dixon, Reg. No. 32,410
Mimi C. Goller, Reg. No. 39,046
Daniel J. Hulseberg, Reg. No. 36,554
Neal D. Marcus, Reg. No. 35,267
James D. McNeil, Reg. No. 26,204

Robert A. Miller, Reg. No. 26,956
Lawrence S. Pope, Reg. No. 26,791
Nicholas A. Poulos, Reg. No. 30,209
Dugal S. Sickert, Reg. No. 33,784
Gregory W. Steele, Reg. No. 33,796
Janelle D. Strode, Reg. No. 34,738
Michael J. Ward, Reg. No. 37,960
David L. Weinstein, Reg. No. 28,128
Steven F. Weinstock, Reg. No. 30,117
Brian R. Woodworth, Reg. No. 33,137
Paul D. Yasger, Reg. No. 37,477

Send Correspondence to:

Steven F. Weinstock
Abbott Laboratories
D-377/AP6D
100 Abbott Park Road
Abbott Park, IL 60064-6050

Direct telephone calls to:

Janelle D. Strode
(847) 937-4395

Name:

James Link

Post Office Address:

1630 Chicago Avenue, Apt. 1405, Evanston, IL 60201
Evanston, Illinois 60201

Residence:

Citizenship:

United States of America

Name: Gang Liu
Post Office Address: 838 Adderly Lane, Gurnee, Illinois 60031
Residence: Gurnee, Illinois 60031
Citizenship: P.R. China

Name: ZhongHua Pei
Post Office Address: 1733 Ceder Glen Dr., Libertyville, Illinois 60048
Residence: Libertyville, Illinois 60048
Citizenship: P.R. China

Name: Tom von Geldern
Post Office Address: 4209 W. Solon Road, Richmond, Illinois 60071
Residence: Richmond, Illinois 60071
Citizenship: United States of America

Name: Martin Winn
Post Office Address: 1263 Carlisle Place, Deerfield, Illinois 60015
Residence: Deerfield, Illinois 60015
Citizenship: United States of America

Name: Zhili Xin
Post Office Address: 29607 N. Birch Ave, Lake Bluff Illinois 60044
Residence: Lake Bluff, Illinois 60044
Citizenship: United States of America

Name: Steven A. Boyd
Post Office Address: 410 Lakeview Dr., Mundelein, Illinois 60060
Residence: Mundelein, IL 60060
Citizenship: United States of America

Name: Gui-Dong Zhu
Post Office Address: 1395 Alamaden Lane, Gurnee, Illinois 60030
Residence: Gurnee, Illinois 60030
Citizenship: People's Republic of China

Name: Jennifer C. Freeman
Post Office Address: 142 Cecelia Street, Grayslake, Illinois 60030
Residence: Grayslake, Illinois 60030
Citizenship: United States of America

Name: Indrani W. Gunawardana
Post Office Address: 622 Nordic Court, Libertyville, Illinois 60048
Residence: Libertyville, Illinois 60048
Citizenship: United States of America

Name: Michael A. Staeger
Post Office Address: 4919 S. 35th Street, Greenfield, Wisconsin 53221
Residence: Greenfield, Wisconsin 53221
Citizenship: United States of America

Name: Hwan-Soo Jae
Post Office Address: 477 Woodlawn Ave, Glencoe, Illinois 60022
Residence: Glencoe, Illinois 60022
Citizenship: United States of America

Name: John K. Lynch
Post Office Address: 8736 44th Ave, Kenosha, Wisconsin 53142
Residence: Kenosha, WI 53142
Citizenship: United States of America

Name: Sheldon Wang
Post Office Address: 1020 Talbot Lane, Grayslake, Illinois 60030
Residence: Grayslake, Illinois 60030
Citizenship: United States of America

I further declare that all statements made herein of my own knowledge are true and that all statements made on information and belief are believed to be true; and further that all statements made herein were made with the knowledge that willful false statements and the like so made are punishable by fine or imprisonment, or both, under Section 1001 of Title 18 of the United States Code, and that such willful false statements may jeopardize the validity of the application or any patent issuing thereon.

James Link Date

Gang Liu Date

ZhongHau Pei Date

Tom von Geldern Date

Martin Winn Date

Zhili Xin Date

Steve A. Boyd Date

Gui-Dong Zhu Date

Jennifer C. Freeman Date

Indrani W. Gunawardana Date

Michael A. Staeger Date

Hwan-Soo Jae Date

John K. Lynch Date

Sheldon Wang Date